

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law. Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Quantum work and the thermodynamic cost of quantum measurements

Sebastian Deffner,^{1,2} Juan Pablo Paz,^{3,4} and Wojciech H. Zurek¹

¹*Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

²*Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

³*Departamento de Física, FCEyN, UBA, Ciudad Universitaria Pabellón 1, 1428 Buenos Aires, Argentina*

⁴*IFIBA CONICET, FCEyN, UBA, Ciudad Universitaria Pabellón 1, 1428 Buenos Aires, Argentina*

(Received 28 March 2016; revised manuscript received 6 May 2016; published 7 July 2016)

Quantum work is usually determined from two projective measurements of the energy at the beginning and at the end of a thermodynamic process. However, this paradigm cannot be considered thermodynamically consistent as it does not account for the thermodynamic cost of these measurements. To remedy this conceptual inconsistency we introduce a paradigm that relies only on the expected change of the average energy given the initial energy eigenbasis. In particular, we completely omit quantum measurements in the definition of quantum work, and hence quantum work is identified as a thermodynamic quantity of only the system. As main results we derive a modified quantum Jarzynski equality and a sharpened maximum work theorem in terms of the information free energy. A comparison of our results with the standard approach allows one to quantify the informational cost of projective measurements.

DOI: [10.1103/PhysRevE.94.010103](https://doi.org/10.1103/PhysRevE.94.010103)

Introduction. In classical mechanics and thermodynamics work is determined by a functional of a force along a trajectory in phase space [1–3]. For quantum systems the situation is much more subtle, since trajectories simply do not exist and a Hermitian work operator cannot be defined [4,5]. Rather, quantum work is commonly determined as the difference between two energies projectively measured at two times: one at the beginning and one at the end of a thermodynamic process [6,7]. Despite the obvious limitations of this notion—no truly open systems can be described since the change of internal energy comprises only work, but no heat—the two-time energy measurement approach has proven to be practical and powerful. For instance, this approach has led to the experimental verification of quantum fluctuation theorems [8,9] and the development of heat engines at the nanoscale [10–13].

Nevertheless, quantum work has remained somewhat elusive with many peculiar features and open questions [14,15]. For instance, it has only recently been pointed out that work can be measured at a single (final) time by means of a generalized measurement [16], that its probability distribution can be interferometrically estimated [17–20], that it reduces to the classical notion of thermodynamic work in high-temperature [21] and semiclassical [22] limits, and that the paradigm of two-time energy measurements is also applicable to open systems as long as the dynamics is unital [23] (see Refs. [24–30]).

In the following we will motivate and introduce a definition of quantum work based on a proper characterization of the role of quantum measurements—a feature not present in the semiclassical limit [22]. Imagine the typical situation for which the quantum Jarzynski equality is valid [3]: at $t = 0$ the system is prepared in a Boltzmann-Gibbs distribution, $\rho_0 \propto \exp(-\beta H_0)$, where H_0 is the initial Hamiltonian and β denotes the inverse temperature. Then the density operator ρ_0 is diagonal in energy basis, and the projective measurement of the energy simply determines the thermal occupation probabilities $p(n_0) \propto \exp[-\beta \epsilon(n_0)]$. After the first measurement, the system evolves under a time-dependent Hamiltonian $H(t)$

from $t = 0$ to $t = \tau$, and the final state, ρ_τ , is typically a complicated nonequilibrium state [31–33]. Generally, ρ_τ is not diagonal in the energy basis, and therefore a projective measurement of the energy is accompanied by a back-action on the system [34]. In many realistic situations one does not have to worry too much about this back-action. Experimentally it poses a real challenge to fully isolate a quantum system from its environment and hence almost all real quantum systems eventually decohere [35,36]. Interestingly, the energy basis can be *einselected* by the environment [37], and hence ρ_τ quickly becomes close to diagonal in energy eigenbasis. Hence, we believe that the methods put forward here will also be useful in the treatment of open systems (see, e.g., also Ref. [38]). For the time being, however, we will exclusively focus on isolated systems.

For fast processes—faster than the time scales over which decoherence happens—or small systems ρ_τ is generally not diagonal, which has been seen explicitly, e.g., in the ion trap experiment by An *et al.* [9]. In such situations the back-action of the measurement on the system does play a role and results in an increase of von Neumann entropy of the system, i.e., in a change of information. Information, however, is a thermodynamic resource [39], which can be related to a free energy usable to perform additional work [40,41]. Hence, one would expect this additional (information) free energy to explicitly show up in the thermodynamic relations [42,43]. However, in the standard treatment [5] and in experiments [8,9] this informational back-action on the system from the second projective measurement has not been considered. Hence, the paradigm of two-time measurements is thermodynamically incomplete.

In our analysis we will address this issue and resolve the conceptual inconsistency arising from neglecting the informational contribution to the laws of thermodynamics. This will lead to a new notion of quantum work, which relies only on the internal energy as an average of the time-evolved energy eigenstates. We will see that this notion is in full agreement with the first law of thermodynamics—the average work is given by the change of internal energy—but that we can

also quantify the informational contribution to the free energy from projective measurements. As a main result we will derive a modified quantum Jarzynski equality and an associated maximum work theorem, in which the thermodynamic free energy is replaced by the information free energy [41,44–46].

Two-time measurements— notions and issues. We begin by briefly reviewing the paradigm of the two-time energy measurement approach, and establish notions and notations. Here and in the following we consider an isolated quantum system with time-dependent Schrödinger equation $i\hbar|\dot{\psi}_t\rangle = H_t|\psi_t\rangle$. We are interested in describing thermodynamic processes that are induced by varying an external control parameter λ_t during time τ , with $H_t = H(\lambda_t)$.

Commonly, quantum work is determined by the following, experimentally motivated protocol: After preparation of the initial state ρ_0 a projective measurement of the energy is performed; then the system is allowed to evolve under the time-dependent Schrödinger equation, before a second projective energy measurement is performed at $t = \tau$.

For the sake of simplicity and to avoid clutter in the formulas we further assume that the system is initially prepared in a Gibbs state, $\rho_0 = \exp(-\beta H_0)/Z_0$, where β is the inverse temperature and Z_0 is the partition function, $Z_0 = \text{tr}\{\exp(-\beta H_0)\}$. Strictly speaking the projective measurement at $t = 0$ is superfluous for initially thermal states, since such ρ_0 's are diagonal in the energy basis. The internal energy of the system, and therefore the full thermodynamic behavior, can simply be determined by an average over all energy eigenstates equipped with the thermal occupation probabilities [47], $p(n_0) = \exp[-\beta\epsilon(n_0, \lambda_0)]/Z_0$.

The first law of thermodynamics determines that the average work is given by the change of internal energy, $\langle W \rangle = \text{tr}\{\rho_\tau H_\tau\} - \text{tr}\{\rho_0 H_0\}$, where $\rho_\tau = U_\tau \rho_0 U_\tau^\dagger$, and U_τ is the unitary time evolution operator, $U_\tau = \mathcal{T}_> \exp(-i/\hbar \int_0^\tau dt H_t)$, and $\mathcal{T}_>$ denotes time ordering. Accordingly, for a single realization of the two-time measurement protocol the quantum work reads

$$W_{n_0 \rightarrow n_\tau} = \epsilon(n_\tau, \lambda_\tau) - \epsilon(n_0, \lambda_0), \quad (1)$$

where $|n_0\rangle$ is the initial eigenstate with eigenenergy $\epsilon(n_0, \lambda_0)$ and $|n_\tau\rangle$ with $\epsilon(n_\tau, \lambda_\tau)$ describes the final energy eigenstate.

The corresponding quantum work probability distribution (2) is then given by an average over an ensemble of realizations of this protocol, $\mathcal{P}(W) = \langle \delta(W - W_{n_0 \rightarrow n_\tau}) \rangle$, which can be written as [18,48]

$$\mathcal{P}(W) = \sum_{n_0, n_\tau} \delta(W - W_{n_0 \rightarrow n_\tau}) p(n_0; n_\tau). \quad (2)$$

In the latter equation the symbol \sum denotes a sum over the discrete part of the eigenvalues spectrum and an integral over the continuous part. In Eq. (2) $p(n_0; n_\tau)$ are the joint probabilities for detecting n_0 and n_τ in the two energy measurements [18,27,48],

$$p(n_0; n_\tau) = \text{tr}\{\Pi_{n_\tau} U_\tau \Pi_{n_0} \rho_0 \Pi_{n_0} U_\tau^\dagger\}, \quad (3)$$

where Π_n denotes the projector into the space spanned by the n th eigenstate, which becomes for nondegenerate spectra $\Pi_n = |n\rangle\langle n|$. It is then a simple exercise to show that from the definition of $\mathcal{P}(W)$ in (2) we have the quantum Jarzynski

equality [4,5],

$$\langle e^{-\beta W} \rangle = \int dW \mathcal{P}(W) e^{-\beta W} = e^{-\beta \Delta F}, \quad (4)$$

where $\Delta F = F_\tau - F_0$ and $F_t = -(1/\beta) \ln(Z_t)$.

Neglected informational cost. Generally the final state ρ_τ is a complicated nonequilibrium state. This means, in particular, that ρ_τ does not commute with the final Hamiltonian H_τ , and one has to consider the back-action on the system due to the projective measurement of the energy [34]. For a single measurement, Π_{n_τ} , the postmeasurement state is given by $\Pi_{n_\tau} \rho_\tau \Pi_{n_\tau} / p_{n_\tau}$, where $p_{n_\tau} = \text{tr}\{\Pi_{n_\tau} \rho_\tau\}$. Thus, the system can be found on average in

$$\rho_\tau^M = \sum_{n_\tau} \Pi_{n_\tau} \rho_\tau \Pi_{n_\tau}. \quad (5)$$

Accordingly, the final measurement of the energy is accompanied by a change of information, i.e., by a change of the von Neumann entropy of the system

$$\Delta \mathcal{H}^M = -\text{tr}\{\rho_\tau^M \ln(\rho_\tau^M)\} + \text{tr}\{\rho_\tau \ln(\rho_\tau)\} \geq 0. \quad (6)$$

Information, however, is physical [49] and its acquisition “costs” work. This additional work has to be paid by the external observer—the measurement device. In a fully consistent thermodynamic framework this cost has to be taken into consideration [40], in particular when calculating the efficiency of thermodynamic devices [10,11].

Quantum work without measurements. To remedy this conceptual inconsistency arising from neglecting the informational contribution of the projective measurements, we propose an alternative paradigm. For isolated systems quantum work is clearly given by the change of internal energy. As a statement of the first law of thermodynamics this holds true no matter whether the system is measured or not. Quantum measurements, however, can be understood as an interaction with a “measuring environment”. Moreover, almost any environment induces decoherence [36]. Thus, defining work with the help of the environment and ignoring the effect of decoherence is as thermodynamically inconsistent as defining work via an external measurement and neglecting the informational cost of these projective measurements.

For thermal states measuring the energy is superfluous as state and energy commute. Hence, a notion of quantum work can be formulated that is fully based on the time evolution of energy eigenstates [50]. Quantum work for a single realization is then determined by considering how much the expectation value for a single energy eigenstate changes under the unitary evolution [51]. Hence, we define

$$\tilde{W}_{n_0} \equiv \langle n_0 | U_\tau^\dagger H_\tau U_\tau | n_0 \rangle - \epsilon(n_0, \lambda_0). \quad (7)$$

We can easily verify that the so defined quantum work (7), indeed, fulfills the first law. To this end, we compute the average work $\langle W \rangle_{\tilde{\mathcal{P}}}$ for the modified quantum work distribution $\tilde{\mathcal{P}}(W)$, and we obtain

$$\begin{aligned} \langle W \rangle_{\tilde{\mathcal{P}}} &= \sum_{n_0} \langle n_0 | U_\tau^\dagger H_\tau U_\tau | n_0 \rangle p(n_0) - \text{tr}\{\rho_0 H_0\} \\ &= \text{tr}\{\rho_\tau H_\tau\} - \text{tr}\{\rho_0 H_0\} = \langle W \rangle. \end{aligned} \quad (8)$$

It is important to note that the average quantum work determined from two-time energy measurements is identical to the (expected) value given only knowledge from a single measurement at $t = 0$. Most importantly, however, in our paradigm the external observer does not have to pay a thermodynamic cost associated with the change of information due to measurements. Hence, the present paradigm can be considered thermodynamically consistent and complete.

Modified quantum Jarzynski equality. What we have seen so far is that the first law of thermodynamics is immune to whether the energy of the system is measured or not, since projective measurements of the energy do not affect the internal energy. However, the informational content of the system of interest, i.e., the entropy, crucially depends on whether the system is measured. Therefore, we expect that the statements of the second law have to be modified to reflect the informational contribution [40]. In our paradigm the modified quantum work distribution becomes

$$\tilde{\mathcal{P}}(W) = \sum_{n_0} \delta(W - \tilde{W}_{n_0}) p(n_0), \quad (9)$$

where as before $p(n_0) = \exp[-\beta \epsilon(n_0, \lambda_0)]/Z_0$. Now, we can compute the average exponentiated work,

$$\langle e^{-\beta W} \rangle_{\tilde{\mathcal{P}}} = \frac{1}{Z_0} \sum_{n_0} e^{-\beta \langle n_0 | U_\tau^\dagger H_\tau U_\tau | n_0 \rangle}. \quad (10)$$

The right side of Eq. (10) can be interpreted as the ratio of two partition functions, where Z_0 describes the initial thermal state. The second partition function $\tilde{Z}_\tau \equiv \sum_{n_0} \exp(-\beta \langle n_0 | U_\tau^\dagger H_\tau U_\tau | n_0 \rangle)$ corresponds to the best possible guess for a thermal state of the final system given only the time-evolved energy eigenbasis. This state can be written as

$$\tilde{\rho}_\tau \equiv \frac{1}{\tilde{Z}_\tau} \sum_{n_0} e^{-\beta \langle n_0 | U_\tau^\dagger H_\tau U_\tau | n_0 \rangle} U_\tau | n_0 \rangle \langle n_0 | U_\tau^\dagger, \quad (11)$$

which differs from the true thermal state, $\rho_\tau^{\text{eq}} = \exp(-\beta H_\tau)/Z_\tau$.

In information theory the “quality” of such a best possible guess is quantified by the relative entropy [52,53], which measures the distinguishability of two (quantum) states. Hence, let us consider

$$S(\tilde{\rho}_\tau || \rho_\tau^{\text{eq}}) = \text{tr}\{\tilde{\rho}_\tau \ln(\tilde{\rho}_\tau)\} - \text{tr}\{\tilde{\rho}_\tau \ln(\rho_\tau^{\text{eq}})\}, \quad (12)$$

for which we compute both terms separately. For the first term, the negentropy of $\tilde{\rho}_\tau$, we obtain

$$\begin{aligned} \text{tr}\{\tilde{\rho}_\tau \ln(\tilde{\rho}_\tau)\} &= -\ln(\tilde{Z}) - \beta \text{tr}\left\{\tilde{\rho}_\tau \sum_{m_0} \langle m_0 | U_\tau^\dagger H_\tau U_\tau | m_0 \rangle U_\tau | m_0 \rangle \langle m_0 | U_\tau^\dagger\right\} \\ &= -\ln(\tilde{Z}) - \beta \tilde{E}, \end{aligned} \quad (13)$$

where we introduced the expected value of the energy, \tilde{E} , under the time-evolved eigenstates,

$$\tilde{E} = \frac{1}{\tilde{Z}} \sum_{n_0} e^{-\beta \langle n_0 | U_\tau^\dagger H_\tau U_\tau | n_0 \rangle} \langle n_0 | U_\tau^\dagger H_\tau U_\tau | n_0 \rangle. \quad (14)$$

The second term of Eq. (12), the cross entropy of $\tilde{\rho}_\tau$ and ρ_τ^{eq} , simplifies to

$$\begin{aligned} \text{tr}\{\tilde{\rho}_\tau \ln(\rho_\tau^{\text{eq}})\} &= -\ln(Z_\tau) - \beta \text{tr}\left\{\sum_{n_0} \frac{1}{\tilde{Z}} e^{-\beta \langle n_0 | U_\tau^\dagger H_\tau U_\tau | n_0 \rangle} U | n_0 \rangle \langle n_0 | U^\dagger H_\tau\right\} \\ &= -\ln(Z_\tau) - \beta \tilde{E}. \end{aligned} \quad (15)$$

Hence, the modified quantum Jarzynski equality (10) becomes

$$\langle e^{-\beta W} \rangle_{\tilde{\mathcal{P}}} = e^{-\beta \Delta F} e^{-S(\tilde{\rho}_\tau || \rho_\tau^{\text{eq}})}, \quad (16)$$

where as before $\Delta F = -1/\beta \ln(Z_\tau/Z_0)$. Jensen’s inequality further implies,

$$\beta \langle W \rangle \geq \beta \Delta F + S(\tilde{\rho}_\tau || \rho_\tau^{\text{eq}}), \quad (17)$$

where we used $\langle W \rangle_{\tilde{\mathcal{P}}} = \langle W \rangle$ [Eq. (8)].

Equations (16) and (17) are our main results. By defining quantum work as an average over time-evolved eigenstates we obtain a modified quantum Jarzynski equality (16) and a generalized maximum work theorem (17), in which the thermodynamic cost of projective measurements becomes apparent. These results become even more transparent by noting that similar versions of the maximum work theorem have been derived in the thermodynamics of information [40,41,44,54]. In this context it has proven useful to introduce the notion of an information free energy,

$$\tilde{F}_\tau = F_\tau + S(\tilde{\rho}_\tau || \rho_\tau^{\text{eq}})/\beta. \quad (18)$$

This free energy is a true thermodynamic quantity [41,44] that accounts for the additional capacity of a thermodynamic system to perform work due to information [40]. Note that in the present context \tilde{F}_τ is computed for the fictitious thermal state $\tilde{\rho}_\tau$ [Eq. (11)], whereas one usually considers the information free energy for the nonequilibrium state ρ_τ [40,41,44,54].

We can rewrite Eq. (17) as

$$\beta \langle W \rangle \geq \beta \Delta \tilde{F}. \quad (19)$$

Equation (19) constitutes a sharper bound than the usual maximum work theorem, and it accounts for the extra free energy available to the system. Free energy, however, describes the usable, extractable work. In real-life applications one is more interested in the maximal free energy the system has available, than in the work that could be extracted by intermediate, disruptive measurements of the energy. Therefore, our treatment could be considered thermodynamically more relevant than the two-time measurement approach.

Illustrative example: Parametric harmonic oscillator. For the remainder of this discussion we will turn to an analytically solvable example, namely, the parametric harmonic oscillator,

$$H = p^2/2m + m\omega_t^2 x^2/2. \quad (20)$$

This system has been studied extensively in the literature, and it can be shown that the average work performed by changing the angular frequency from ω_0 to ω_τ is given by [21,31]

$$\langle W \rangle = \hbar/2(Q^* \omega_\tau - \omega_0) \coth(\beta/2\hbar\omega_0). \quad (21)$$

The quantity Q^* is a measure of adiabaticity [21,55], which fully encodes the dynamics. In particular, we have $Q^* = 1$ for




FIG. 1. Average work $\langle W \rangle$ [Eq. (21)] (blue, solid line), together with the change of equilibrium free energy ΔF [Eq. (22)] (red, dotted line) and the informational maximum work theorem, $\beta\Delta F + S(\tilde{\rho}_\tau || \rho_\tau^{\text{eq}})$ [Eq. (17)] (purple, dashed line) for the parametric harmonic oscillator (20) with $\hbar = 1$, $\beta = 1$, $\omega_0 = 1$, and $\omega_\tau = 2$.




FIG. 2. Average work $\langle W \rangle$ [Eq. (21)] (blue, solid line), together with the change of equilibrium free energy ΔF [Eq. (22)] (red, dotted line) and the informational maximum work theorem, $\beta\Delta F + S(\tilde{\rho}_\tau || \rho_\tau^{\text{eq}})$ [Eq. (17)] (purple, dashed line) for the parametric harmonic oscillator (20) with $\hbar = 1$, $\beta = 1$, $\omega_0 = 2$, and $\omega_\tau = 1$.

adiabatic, infinitely slow processes, and $Q^* > 1$ for finite time driving. The change in equilibrium free energy becomes [31]

$$\Delta F = \frac{1}{\beta} \ln \left(\frac{\sinh(\beta/2\hbar\omega_\tau)}{\sinh(\beta/2\hbar\omega_0)} \right). \quad (22)$$

Therefore, we merely have to compute the partition function \tilde{Z} , which can be written as

$$\tilde{Z}_\tau = \sum_{n_0} \exp \left(- \sum_{n_\tau} \beta \hbar \omega_\tau (n_\tau + 1/2) p_{n_0, n_\tau} \right), \quad (23)$$

where $p_{n_0, n_\tau} = \text{tr} \{ \Pi_{n_\tau} U_\tau \Pi_{n_0} U_\tau^\dagger \}$. Hence, \tilde{Z} is fully determined by the average final occupation number [31]

$$\langle n_\tau \rangle_{n_0} = \sum_{n_\tau} n_\tau p_{n_0, n_\tau} = (n_0 + 1/2) Q^* - 1/2, \quad (24)$$

from which we obtain

$$\begin{aligned} \tilde{Z}_\tau &= \sum_{n_0} \exp(-\beta \hbar \omega_\tau Q^* (n_0 + 1/2)) \\ &= [2 \sinh(\beta/2 Q^* \hbar \omega_\tau)]^{-1}. \end{aligned} \quad (25)$$

Accordingly, the informational correction to the maximum work theorem arising from omitting the second projective measurement becomes

$$S(\tilde{\rho}_\tau || \rho_\tau^{\text{eq}}) = \ln \left(\frac{\sinh(\beta/2 Q^* \hbar \omega_\tau)}{\sinh(\beta/2 \hbar \omega_\tau)} \right), \quad (26)$$

which is clearly non-negative and a simple function of the measure of adiabaticity Q^* . Note that for adiabatic processes, $Q^* = 1$, the information free energy (17) becomes identical to the equilibrium free energy, since we have $S(\tilde{\rho}_\tau || \rho_\tau^{\text{eq}}) = 0$.

In Figs. 1 and 2 we plot the average work $\langle W \rangle$ [Eq. (21)] together with the change of equilibrium free energy ΔF [Eq. (22)] and the modified maximum work theorem (17). We observe that the bound arising from the information free energy (18) is process dependent and sharper—for some parametrizations even tight.

Concluding remarks. A conceptually consistent and complete framework of quantum thermodynamics crucially depends on accounting for quantum features and peculiarities [56]. Indeed, one can hope that a completely quantum approach based on symmetries of entanglement [57] can help resolve outstanding problems of the classical and quantum points of view. In the present analysis we have shown that despite its success the two-time energy measurement approach to quantum work neglects the informational back-action of the projective measurements. This informational contribution to the laws of quantum thermodynamics has been highlighted by introducing a paradigm in which quantum work is fully determined by the change of internal energy as an average over the initial energy eigenstates. This approach has allowed us to derive a modified quantum Jarzynski equality and a modified maximum work theorem, in which the equilibrium free energy is replaced by the information free energy. In conclusion, we achieved several important insights: (i) we have proposed a thermodynamically consistent notion of quantum work, which does not rely on external observers and projective measurements and (ii) we have included the thermodynamic cost of information gain in the paradigm of quantum work, and hence taken an instrumental step towards a conclusive theory of quantum thermodynamics of quantum information.

Acknowledgments. It is a pleasure to thank Jordan M. Horowitz, Christopher Jarzynski, Jim Crutchfield, and Gavin Crooks for interesting and insightful discussions. S.D. acknowledges financial support by the U.S. Department of Energy through a LANL Director's Funded Fellowship, and W.H.Z. acknowledges partial support by the Foundational Questions Institute Grant No. 2015-144057 on "Physics of What Happens".

- [1] H. B. Callen, *Thermodynamics and An Introduction to Thermostatistics*, 2nd ed. (Wiley, New York, 1985).
- [2] E. Boksénbojm, B. Wynants, and C. Jarzynski, *Physica A* **389**, 4406 (2010).
- [3] S. Deffner, *Europhys. Lett.* **103**, 30001 (2013).
- [4] P. Talkner, E. Lutz, and P. Hänggi, *Phys. Rev. E* **75**, 050102 (2007).
- [5] M. Campisi, P. Hänggi, and P. Talkner, *Rev. Mod. Phys.* **83**, 771 (2011).
- [6] J. Kurchan, [arXiv:cond-mat/0007360](https://arxiv.org/abs/cond-mat/0007360).
- [7] H. Tasaki, [arXiv:cond-mat/0009244](https://arxiv.org/abs/cond-mat/0009244).
- [8] T. B. Batalhão, A. M. Souza, L. Mazzola, R. Auccaise, R. S. Sarthour, I. S. Oliveira, J. Goold, G. De Chiara, M. Paternostro, and R. M. Serra, *Phys. Rev. Lett.* **113**, 140601 (2014).
- [9] S. An, J.-N. Zhang, M. Um, D. Lv, Y. Lu, J. Zhang, Z.-Q. Yin, H. T. Quan, and K. Kim, *Nat. Phys.* **11**, 193 (2015).
- [10] O. Abah, J. Roßnagel, G. Jacob, S. Deffner, F. Schmidt-Kaler, K. Singer, and E. Lutz, *Phys. Rev. Lett.* **109**, 203006 (2012).
- [11] J. Roßnagel, O. Abah, F. Schmidt-Kaler, K. Singer, and E. Lutz, *Phys. Rev. Lett.* **112**, 030602 (2014).
- [12] A. Dechant, N. Kiesel, and E. Lutz, *Phys. Rev. Lett.* **114**, 183602 (2015).
- [13] K. Zhang, F. Bariani, and P. Meystre, *Phys. Rev. Lett.* **112**, 150602 (2014).
- [14] P. Hänggi and P. Talkner, *Nat. Phys.* **11**, 108 (2015).
- [15] P. Talkner and P. Hänggi, *Phys. Rev. E* **93**, 022131 (2016).
- [16] A. J. Roncaglia, F. Cerisola, and J. P. Paz, *Phys. Rev. Lett.* **113**, 250601 (2014).
- [17] V. Vedral, *J. Phys. A: Math. Theor.* **45**, 272001 (2012).
- [18] D. Kafri and S. Deffner, *Phys. Rev. A* **86**, 044302 (2012).
- [19] B. Prasanna Venkatesh, G. Watanabe, and P. Talkner, *New J. Phys.* **16**, 015032 (2014).
- [20] L. Fusco, S. Pigeon, T. J. G. Apollaro, A. Xuereb, L. Mazzola, M. Campisi, A. Ferraro, M. Paternostro, and G. De Chiara, *Phys. Rev. X* **4**, 031029 (2014).
- [21] S. Deffner, O. Abah, and E. Lutz, *Chem. Phys.* **375**, 200 (2010).
- [22] C. Jarzynski, H. T. Quan, and S. Rahav, *Phys. Rev. X* **5**, 031038 (2015).
- [23] A unital map is a completely positive, trace preserving map that also preserves the identity. Thermodynamically, a system evolving under unital dynamics can be understood as an open system that is so weakly coupled that it only exchanges information but no heat with the environment.
- [24] A. E. Rastegin, *J. Stat. Mech. Theory Exp.* (2013) P06016.
- [25] T. Albash, D. A. Lidar, M. Marvian, and P. Zanardi, *Phys. Rev. E* **88**, 032146 (2013).
- [26] A. E. Rastegin and K. Życzkowski, *Phys. Rev. E* **89**, 012127 (2014).
- [27] S. Deffner and A. Saxena, *Phys. Rev. Lett.* **114**, 150601 (2015).
- [28] S. Deffner and A. Saxena, *Phys. Rev. E* **92**, 032137 (2015).
- [29] B. Gardas, S. Deffner, and A. Saxena, *Sci. Rep.* **6**, 23408 (2016).
- [30] G. Manzano, J. M. Horowitz, and J. M. R. Parrondo, *Phys. Rev. E* **92**, 032129 (2015).
- [31] S. Deffner and E. Lutz, *Phys. Rev. E* **77**, 021128 (2008).
- [32] S. Deffner and E. Lutz, *Phys. Rev. Lett.* **105**, 170402 (2010).
- [33] S. Deffner and E. Lutz, *Phys. Rev. E* **87**, 022143 (2013).
- [34] M. L. Nielsen and I. L. Chuang, *Quantum Computation and Quantum Information* (Cambridge University Press, Cambridge, UK, 2000).
- [35] M. Brune, E. Hagley, J. Dreyer, X. Maître, A. Maali, C. Wunderlich, J. M. Raimond, and S. Haroche, *Phys. Rev. Lett.* **77**, 4887 (1996).
- [36] W. H. Zurek, *Rev. Mod. Phys.* **75**, 715 (2003).
- [37] J. P. Paz and W. H. Zurek, *Phys. Rev. Lett.* **82**, 5181 (1999).
- [38] M. Esposito and S. Mukamel, *Phys. Rev. E* **73**, 046129 (2006).
- [39] To the best of our knowledge, the thermodynamic consequences of measurements have only been discussed for thermodynamic averages [42,58,59], whereas the present focus is on fluctuation theorems.
- [40] S. Deffner and C. Jarzynski, *Phys. Rev. X* **3**, 041003 (2013).
- [41] J. M. R. Parrondo, J. M. Horowitz, and T. Sagawa, *Nat. Phys.* **11**, 131 (2015).
- [42] K. Jacobs, *Phys. Rev. E* **86**, 040106 (2012).
- [43] J. Yi and Y. W. Kim, *Phys. Rev. E* **88**, 032105 (2013).
- [44] S. Deffner and E. Lutz, [arXiv:1201.3888](https://arxiv.org/abs/1201.3888) [cond-mat].
- [45] D. A. Sivak and G. E. Crooks, *Phys. Rev. Lett.* **108**, 150601 (2012).
- [46] S. Still, D. A. Sivak, A. J. Bell, and G. E. Crooks, *Phys. Rev. Lett.* **109**, 120604 (2012).
- [47] F. Brito, F. Rouxinol, M. D. LaHaye, and A. O. Caldeira, *New J. Phys.* **17**, 075002 (2015).
- [48] A. Leonard and S. Deffner, *Chem. Phys.* **446**, 18 (2015).
- [49] R. Landauer, *Phys. Today* **44**(5), 23 (1991).
- [50] The thermal state is then emulated [47] by equipping each energy eigenstate with the thermal occupation probabilities, $p(n_0) = \exp[-\beta\epsilon(n_0, \lambda_0)]/Z_0$.
- [51] It is interesting to note that a similar notion of quantum work has been considered in a different context in Refs. [60,61].
- [52] S. Kullback and R. A. Leibler, *Ann. Math. Stat.* **22**, 79 (1954).
- [53] H. Umegaki, *Tohoku Math. J.* **6**, 177 (1954).
- [54] S. Vaikuntanathan and C. Jarzynski, *Europhys. Lett.* **87**, 60005 (2009).
- [55] K. Husimi, *Prog. Theor. Phys.* **9**, 381 (1953).
- [56] J. Gemmer, M. Michel, and G. Mahler, *Quantum Thermodynamics* (Springer, Berlin/Heidelberg, 2009).
- [57] S. Deffner and W. H. Zurek, *New J. Phys.* **18**, 063013 (2016).
- [58] K. Jacobs, *Phys. Rev. A* **80**, 012322 (2009).
- [59] P. Kammerlander and J. Anders, *Sci. Rep.* **6**, 22174 (2016).
- [60] A. E. Allahverdyan and T. M. Nieuwenhuizen, *Phys. Rev. E* **71**, 066102 (2005).
- [61] M. Campisi, *New J. Phys.* **15**, 115008 (2013).