

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law. Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

SCIENTIFIC REPORTS

OPEN

Non-hermitian quantum thermodynamics

Bartłomiej Gardas^{1,2}, Sebastian Deffner^{1,3} & Avadh Saxena^{1,3}

Received: 21 December 2015

Accepted: 03 March 2016

Published: 22 March 2016

Thermodynamics is the phenomenological theory of heat and work. Here we analyze to what extent quantum thermodynamic relations are immune to the underlying mathematical formulation of quantum mechanics. As a main result, we show that the Jarzynski equality holds true for all non-hermitian quantum systems with real spectrum. This equality expresses the second law of thermodynamics for isothermal processes arbitrarily far from equilibrium. In the quasistatic limit however, the second law leads to the Carnot bound which is fulfilled even if some eigenenergies are complex provided they appear in conjugate pairs. Furthermore, we propose two setups to test our predictions, namely with strongly interacting excitons and photons in a semiconductor microcavity and in the non-hermitian tight-binding model.

More and more non-hermitian systems are becoming experimentally accessible¹. Therefore, it has become evident that questions concerning foundations of quantum mechanics are no longer only of academic interest. Recent experiments have demonstrated that hermiticity may not be as fundamental as mandated by quantum mechanics^{2,3}. For instance, in⁴ a spontaneous \mathcal{PT} -symmetry breaking has been observed indicating a condition weaker than hermiticity (namely \mathcal{PT}^5) being realized in nature. Furthermore, in⁶ exceptional eigenenergies of complex value have been measured challenging the reality of the spectrum imposed by hermiticity.

Conventional quantum mechanics is built upon the Dirac-von Neumann axioms^{2,3}. These state that if \mathcal{H} is a complex Hilbert space of countable, infinite dimension, then (i) observables of a quantum system are defined as hermitian operators O on \mathcal{H} , (ii) quantum states $|\phi\rangle$ are unit vectors in \mathcal{H} , and (iii) the expectation value of an observable O in a state $|\phi\rangle$ is given by the inner product, $\langle O \rangle = \langle \phi | O \phi \rangle$. Interestingly, only axioms (ii) and (iii) are of mathematical necessity needed for a proper probabilistic, physical theory. To demand, however, that any quantum mechanical theory has to be built on hermitian operators is rather mathematically convenient than being fundamentally necessary^{5,7}.

In particular, the restriction to hermitian observables excludes the description of, for instance, quantum field theories with \mathcal{PT} -symmetry, cases where the language of quantum mechanics is used for problems within classical statistical mechanics or diffusion in biological systems, or cases where effective complex potentials are introduced to describe interactions at edges⁸. Particularly striking examples are optical systems with complex index of refraction. Imagine, for instance, polarized light in a stratified, nontransparent, biaxially anisotropic, dielectric medium warped cyclically along the propagation direction. For such systems it has been shown⁹ that not only a non-hermitian description becomes necessary, but also that physical intuition has to be invoked carefully. For instance, Berry highlighted⁹ that adiabatic intuition can be countered dramatically for systems with non-hermitian Hamiltonians.

Very recently, it has become evident that for a special class of non-hermitian systems, namely in \mathcal{PT} -symmetric quantum mechanics¹⁰, the quantum Jarzynski equality holds without modification¹¹. For isolated quantum systems evolving under unitary dynamics the so-called two-time energy measurement approach has proven to be practical and powerful. In this paradigm, quantum work is determined by projective energy measurements at the beginning and the end of a process induced by an externally controlled Hamiltonian. The Jarzynski equality¹² together with subsequent Nonequilibrium Work Theorems, such as the Crooks fluctuation theorem¹³, is undoubtedly among the most important breakthroughs in modern Statistical Physics¹⁴. Jarzynski showed that for isothermal processes the second law of thermodynamics can be formulated as an *equality*, no matter how far from equilibrium the system is driven¹², $\langle \exp(-\beta W) \rangle = \exp(-\beta \Delta F)$. Here β is the inverse temperature of the environment, and ΔF is the free energy difference, *i.e.*, the work performed during an infinitely

¹Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA. ²Institute of Physics, University of Silesia, 40-007 Katowice, Poland. ³Center for Nonlinear Studies, Los Alamos National Laboratory, Los Alamos, NM 87545, USA. Correspondence and requests for materials should be addressed to B.G. (email: bartek.gardas@gmail.com)

slow process. The angular brackets denote the average over an ensemble of finite-time realizations of the process characterized by their nonequilibrium work W .

The present study is dedicated to an even more fundamental question. In the following we will analyze to what extent quantum thermodynamic relations are immune to the underlying mathematical formulation of quantum mechanics. Contrary to different studies (see *e.g.*¹¹) conducted on a similar subject we present the *broadest* possible class of non-hermitian systems that still allows a thermodynamic theory in the “conventional” sense.

As a main result we will prove that equilibrium as well as non-equilibrium identities of quantum thermodynamics hold, without modification also for quantum systems described by pseudo-hermitian Hamiltonians¹⁵. Those systems have either entirely real spectrum or complex eigenvalues appear in complex conjugate pairs. In particular, we will show that the Carnot statement of the second law of thermodynamics holds for any such system and that the quantum Jarzynski equality is not violated as long as the eigenvalue spectrum is real. If the two-time energy measurement could be realized *e.g.* in a microcavity⁶, then the Jarzynski equality for pseudo-hermitian systems could be put into a test (see *Discussion*).

Fundamentals of Pseudo-Hermitian Quantum Mechanics

To address physical properties of recent experiments^{4,6} we start by briefly reviewing the mathematical foundations of pseudo-hermitian quantum mechanics⁸. Let H be a general, non-hermitian Hamiltonian of a physical system, and we assume for the sake of simplicity that the spectrum of H , $\{E_n\}$, is discrete (possibly degenerate). Such a Hamiltonian is of physical relevance only if it is measurable, *i.e.*, if a representation of the eigenbasis $|E_{n,\alpha}\rangle$ is experimentally accessible. Then H is diagonal in this basis. Here n is the quantum number and α counts possible degeneracy. Diagonalizability of H is *equivalent* to the existence of biorthonormal set of left, $|\phi_{n,\alpha}\rangle$, and right, $|\psi_{n,\alpha}\rangle$, eigenvectors¹⁶. In general, the energy eigenvalues are complex, and the eigenvalue problem reads¹⁵

$$H|\psi_{n,\alpha}\rangle = E_n|\psi_{n,\alpha}\rangle, \quad H^\dagger|\phi_{n,\alpha}\rangle = E_n^*|\phi_{n,\alpha}\rangle, \quad (1)$$

with $\langle\psi_{n,\alpha}|\phi_{m,\beta}\rangle = \delta_{mn}\delta_{\alpha\beta}$ and $\sum_{n,\alpha}|\psi_{n,\alpha}\rangle\langle\phi_{n,\alpha}| = \mathbb{I}$. A non-hermitian Hamiltonian such as (1) is called pseudo-hermitian if a g exists such that

$$H^\dagger = gHg^{-1} \quad \text{and} \quad g = g^\dagger. \quad (2)$$

It does exist *if and only if* either all eigenenergies are real or complex ones appear in conjugate pairs with the same degeneracy¹⁵. If none of those criteria are met H is generally non-hermitian⁸; yet it still can be useful, *e.g.* for an effective description of open quantum systems¹⁷. However, when heat is exchanged the two-time energy measurement can no longer describe the work done during a thermodynamic process. Therefore we shall not focus on such cases here. Another interesting class relates to systems that interact with environments, but do not exchange heat. This phenomenon is called dephasing (loss of information)¹⁸. For such systems, work can still be determined by the two-time energy measurement and the Jarzynski equality holds as well^{19–21}.

Condition (2) assures that H is, in fact, hermitian however with respect to a new inner product, namely

$$\langle\psi|\phi\rangle_g := \langle\psi, g\phi\rangle. \quad (3)$$

Note that g always exists such that $\langle\psi|\phi\rangle_g$ is positive-definite (this is a genuine inner product), and it can be found *if and only if* the spectrum of H is real. To make a consistent definition of work for a quantum system within the two-time energy measurement paradigm its spectrum has to be real. Therefore, unless stated otherwise, we shall always assume this to be the case. Then, Eq. (2) can be fulfilled by the following positive-definite operators (g is a proper metric operator)²²

$$g = \sum_{n,\alpha}|\phi_{n,\alpha}\rangle\langle\phi_{n,\alpha}|, \quad g^{-1} = \sum_{n,\alpha}|\psi_{n,\alpha}\rangle\langle\psi_{n,\alpha}|. \quad (4)$$

Often, g fulfilling (2) can be deduced easily from physical properties such as the parity reflection or time reversal²³. Nevertheless, only Eq. (4) assures that $\langle\psi|\psi\rangle_g > 0$ for all states $\psi \neq 0$. This means that the proper metric may reflect “symmetries” that are hidden from the observer^{24,25}. For instance, if a rotation V exists such that $V^{-1}HV$ is diagonal in an orthonormal basis, then $g = V^\dagger V$. This follows directly from Eq. (4). The last formula is especially useful in practice. It allows one to find the metric by analyzing an experimental setup (*e.g.* inspecting the orientation of the axis, etc.).

In the following we only consider cases where changes of the Hamiltonian are induced by a time-dependent thermodynamic process λ_t , that is to say $H_t = H(\lambda_t)$. If such changes occur then the metric operator satisfying Eq. (2) is time-dependent. Nevertheless, the dynamics is still governed by a time-dependent Schrödinger equation. However, a slight modification becomes necessary to preserve unitarity^{26,27},

$$i\hbar\partial_t U_t = (H_t + G_t)U_t, \quad G_t = -\frac{i\hbar}{2}g_t^{-1}\partial_t g_t. \quad (5)$$

Above, ∂_t denotes the derivative with respect to time t . The Schrödinger equation (5) can also be rewritten in the standard form, that is, with H_t being the generator. Indeed, it is sufficient to replace ∂_t with a covariant derivative $D_t := \partial_t + g_t^{-1}\partial_t g_t/2$ ²⁸. By construction the unique solution to Eq. (5) obeys the relation

$$U_t^\dagger g_t = g_0 U_t^{-1}, \quad \text{where} \quad g_0 := g_{t=0}. \quad (6)$$

This relation can be viewed as the corresponding unitarity condition similar to the “standard” one, *i.e.*, $U_t^\dagger = U_t^{-1}$. For pseudo-hermitian systems an average value of a non-hermitian observable A , $\text{tr}\{A\}$, can be computed as

$$\text{tr}\{A\} = \sum_{k,\gamma} \langle \psi_{k,\gamma} | g A | \psi_{k,\gamma} \rangle. \quad (7)$$

Formally, this suggests one to use the following Dirac correspondence between bra and ket vectors $|\psi\rangle \leftrightarrow \langle\psi|g^{16}$.

Pseudo-hermitian Jarzynski equality. Having analyzed the mathematical structure of pseudo-hermitian quantum systems, we turn to the physical description to analyze the Jarzynski equality. Without loss of generality and to simplify our notation we assume the spectrum to be non-degenerate.

For an isolated quantum system, the work done during a thermodynamic process λ_t of duration τ is commonly determined by a two-time energy measurement²⁹. At $t = 0$ a projective energy measurement is performed. Next, the system evolves unitarily under the generalized time-dependent Schrödinger equation (5) only to be measured again at $t = \tau$. By averaging over an ensemble of realizations of such processes one can reconstruct the distribution of work values^{30,31},

$$\mathcal{P}(w) = \sum_{n,m} \delta(w - w_{nm}) p_{nm}. \quad (8)$$

Above, p_{nm} denotes a probability that a specific transition $|\psi_n(\lambda_0)\rangle \rightarrow |\psi_m(\lambda_\tau)\rangle$ will occur, whereas $w_{nm} = E_m^\tau - E_n$ is the corresponding work done during this transition. It is important to stress that this work is associated with H_t rather than $H_t + G_t$ as G_t is a gauge field, and hence it can have no influence on physical observables³².

The transition probability p_{nm} can be seen as the joint probability that the first measurement will yield the energy value E_n given the system has been initially prepared in a state ρ_0 , and the probability that the outcome of the second measurement will be E_m^τ given the initial state ψ_n . Therefore,

$$p_{nm} = \text{tr}\{\Pi_n \rho_0\} \times |\langle \psi_m^\tau, g_\tau U_\tau \psi_n \rangle|^2, \quad (9)$$

where U_τ denotes the evolution operator generated by $H_t + G_t$ at time $t = \tau$, whereas $\Pi_n = \langle \psi_n, g_0 \cdot \rangle \psi_n$ is the projector into the space spanned by the n th eigenstate. Since Π_n is not hermitian the formula for probabilities p_{nm} accounts for the metric g , and hence differs from the one usually adopted for hermitian systems³¹.

Assume the system is initially in a Gibbs state, that is $\rho_0 = \exp(-\beta H_0)/Z_0$ with $Z_0 = \text{tr}\{\exp(-\beta H_0)\}$ being the partition function, then

$$p_{nm} = \frac{e^{-\beta E_n}}{Z_0} \langle U_\tau^\dagger g_\tau \psi_m^\tau, \Pi_n U_\tau^{-1} \psi_m^\tau \rangle. \quad (10)$$

To obtain the last expression for p_{nm} we have also invoked the unitarity condition (6). Now, the average exponentiated work can be expressed as

$$\langle e^{-\beta W} \rangle = \int dw \mathcal{P}(w) \exp(-\beta w) = \frac{1}{Z_0} \sum_{m,n} e^{-\beta E_m^\tau} \langle g_\tau \psi_m^\tau, U_\tau \Pi_n U_\tau^{-1} \psi_m^\tau \rangle. \quad (11)$$

Finally, summing out all projectors Π_n and taking into account that $\langle g_\tau \psi_m^\tau, \psi_m^\tau \rangle = 1$ we arrive at

$$\langle e^{-\beta W} \rangle = \frac{1}{Z_0} \sum_m e^{-\beta E_m^\tau} = \frac{Z_\tau}{Z_0} = e^{-\beta \Delta F}, \quad (12)$$

where $F = (-1/\beta) \ln(Z)$ is the system’s free energy.

The last equation shows that the Jarzynski equality holds also for non-hermitian systems that admit real spectrum. This is our first main result. Jarzynski has shown that the second law of thermodynamics for isothermal processes can be expressed as an equality arbitrarily far from equilibrium. Our analysis has shown that his result is true for *all non-hermitian systems with real spectrum*.

Carnot bound. In the preceding section we argued that if the two-time energy measurement can be performed on a non-hermitian quantum system, then the Jarzynski equality holds as long as the eigenenergies are real. Now, we will prove that the Carnot statement of the second law is also true for all pseudo-hermitian systems.

Consider a generic system that operates between two heat reservoirs with hot, T_h , and cold, T_c , temperatures, respectively. Then, the Carnot engine consists of two isothermal processes during which the system absorbs or exhausts heat and two thermodynamically adiabatic, that is, isentropic strokes while the extensive control parameter λ is varied^{33,34}. It is well established that the maximum efficiency η for classical systems, attained in the quasi-static limit, is given by the Carnot bound^{35–37}:

$$\eta = 1 - \frac{T_c}{T_h} < 1. \quad (13)$$

Recent years have witnessed an abundance of research^{38–40} investigating whether quantum correlations can be harnessed to break this limit. Recently, the Carnot limit has been proven to be universal within the usual

framework³³. This limit can be seen as yet another formulation of the second law of thermodynamics for quasistatic processes. We will show that it holds for all pseudo-hermitian systems whether their spectrum is real or not.

We begin by proving that both the energy $E = \text{tr}\{\rho H\}$ and entropy S are real in our present framework. Indeed, from (2) it immediately follows that

$$E^* = \text{tr}\{g\rho g^{-1}gHg^{-1}\} = E, \quad (14)$$

with ρ being a Gibbs thermal state. Interestingly, this result holds true even if some of the eigenvalues E_n are complex. Note, in that case g exists but is not positive definite and thus cannot be expressed like in Eq. (4).

To understand why Eq. (14) holds when complex eigenvalues appear in conjugate pairs note that $|\psi_{n,\alpha}\rangle = g^{-1}|\phi_{n,\alpha}\rangle$, and consider

$$H|\psi_{n,\alpha}\rangle = g^{-1}H^\dagger|\phi_{n,\alpha}\rangle = E_n^*|\psi_{n,\alpha}\rangle, \quad (15)$$

showing that if E_n is in the spectrum of H so is E_n^* . Moreover g^{-1} maps the subspace spanned by all eigenvectors belonging to E_n to that belonging to E_n^* . Since g^{-1} is invertible, the mapping is one-to-one, and the multiplicity of both E_n and E_n^* is the same. An interesting realization of such systems is the non-hermitian tight-binding model⁴¹.

The result (14) can also be obtained directly, that is, without invoking the metric g explicitly. Indeed, we have

$$E = \text{tr}\{\rho H\} = \frac{1}{Z} \sum_n E_n e^{-\beta E_n} = -\frac{1}{2Z} \frac{\partial}{\partial \beta} \sum_{n/2} (e^{-\beta E_n} + e^{-\beta E_n^*}) = E^*. \quad (16)$$

In the present case, the thermodynamic entropy is given by the von Neumann entropy⁴². The latter can be further simplified and it takes the well known form $S = \beta(E - F)$ ³³. Since the partition function Z is real so is the free energy F . Hence, we conclude that the entropy S is real.

According to the first law of thermodynamics⁴³, $dE = \delta Q + \delta W$, there are two forms of energy: heat δQ is the change of internal energy associated with a change of entropy, whereas work δW is the change of internal energy due to the change of an extensive parameter, *i.e.*, change of the Hamiltonian of the system. To identify those contributions we write³³

$$dE = \text{tr}\{\delta\rho H\} + \text{tr}\{\rho\delta H\}. \quad (17)$$

In the quasistatic regime, the second law of thermodynamics for isothermal processes states that $dS = \beta\delta Q$. Combining the latter with (17) proves that (i) δQ and thus δW are real and (ii) the intuitive definitions of heat and work introduced in Ref. 44 apply also to pseudo-hermitian systems.

After completing a cycle, a quantum pseudo-hermitian heat engine has performed work $\langle W \rangle = \langle Q_h \rangle - \langle Q_c \rangle$ and exhausted a portion of heat $\langle Q_c \rangle$ to the cold reservoir. Therefore, the efficiency of such a device is given by³³

$$\eta = \frac{\langle W \rangle}{\langle Q_c \rangle} = 1 - \frac{T_c}{T_h}. \quad (18)$$

In conclusion, we have shown that the Carnot bound, which expresses the second law of thermodynamics for quasistatic processes, holds for all pseudo-hermitian systems. In contrast, the second law for arbitrarily fast processes encoded in the Jarzynski equality (12), *only* holds for all non-hermitian systems with real spectrum.

Discussion

Example 1a. We begin with a model for localization effects in solid state physics⁴¹. The general form of its Hamiltonian in one dimension reads

$$H = \frac{(p - i\xi)^2}{2m} + V(x), \quad (19)$$

where $V(x)$ is a confining potential, and p and x are the momentum and position operators respectively. They obey the canonical commutation relation $[x, p] = i\hbar$. Real parameter ξ expresses an external magnetic field and m is the mass. Using the Baker-Campbell-Hausdorff formula one can verify that

$$e^{2\xi x} p e^{-2\xi x} = p + 2\xi[x, p] + 2\xi^2[x, [x, p]] + \dots = p + 2i\xi. \quad (20)$$

Therefore, since $[V(x), e^{2\xi x}] = 0$, we conclude that H is pseudo-hermitian. The metric $g = e^{2\xi x}$ is positive definite and thus the spectrum of (19) is real. Further, we assume that the corresponding classical potential $V_c(x)$ has a non-vanishing second derivative, and a minimum at $x = 0$ (*e.g.* $V_c'(0) = 0$). Then

$$V_c(x) = V_c'(0)x + \frac{1}{2}V_c''(0)x^2 + O(\delta x^3) \approx \frac{1}{2}m\omega^2 x^2, \quad (21)$$

where $V_c'(0) = \frac{1}{2}m\omega^2$ has been introduced. After quantization, the eigenvalues and eigenvectors of this non-hermitian harmonic oscillator read (for the sake of simplicity we set $m = \hbar = 1$ throughout)




Figure 1. Left panel: Average exponentiated work $\langle e^{-\beta W} \rangle$ (blue curve) as a function of the number of terms N_{\max} included in the summation (11) for the protocol (23). The function quickly converges to $e^{-\beta \Delta F}$ (red curve) showing that the Jarzynski equality (12) holds. Right panel: $\langle W_{\text{irr}} \rangle = \langle W \rangle - \Delta F$ as a function of τ which relates to the speed at which the energy is supplied to the system. The irreversible work $\langle W_{\text{irr}} \rangle \rightarrow 0$ as τ approaches the quasistatic regime. The inset (red curve) shows the irreversible work calculated for a linear protocol, $\omega(t) = \omega_i + (\omega_f - \omega_i)t/\tau$. We see that it takes longer for the system to reach its quasistatic regime. Parameters used in the numerical simulations are: $w_i = 0.2$, $w_f = 0.6$, $N\tau = 1.5$ (left panel) and $N\tau = 3$ (right panel); the remaining parameters were set to 1.

$$\psi_n(x) := \frac{1}{\sqrt{2^n n! \sqrt{\pi}}} H_n(\sqrt{\omega} x) e^{-\omega x^2 - \xi x}, \quad E_n = \omega \left(n + \frac{1}{2} \right), \quad (22)$$

where $H_n(x)$ are the Hermite polynomials.

Now we assume that the size of this harmonic trap (e.g. ω) is changed, and thus g does not depend on time. Experimentally, harmonic traps are sensitive to initial excitations resulting for a discontinuity of the protocol itself at the beginning⁴⁵. The most common way to minimize this effect, while quenching between ω_i and ω_f is to use functions smooth enough at the “edges”, for instance,

$$\omega(t) = \frac{\omega_i + \omega_f}{2} + \frac{\omega_i - \omega_f}{2} \text{erf}(t/\tau), \quad -N\tau < t < N\tau \quad (23)$$

where $\text{erf}(\cdot)$ denotes the error function, τ is a time scale, and N is an integer emulating infinity. The transition probabilities (9) can be expressed via the following integral

$$p_{nm} = \frac{\exp(-\beta \omega_i (n + 1/2))}{\sinh(\beta \omega_i / 2)} \int_{-\infty}^{\infty} e^{2\xi x} \psi_n^{N\tau}(x) \psi_m^*(x, N\tau) dx, \quad (24)$$

where the partition function $Z_0 = 1/\sinh(\beta \omega_i / 2)$ has been calculated exactly; and $\psi_m(x, N\tau) = U_{N\tau} \psi_m(x)$ is the solution of Eq. (5), with the initial condition given by (22), at $t = N\tau$. Although $\psi_m(x, N\tau)$ cannot be obtained analytically, a closed form expressed in terms of a solution to the corresponding classical equation of motion can be found (see e.g. Ref. 46).

Figure 1 (Left panel) shows the average exponentiated work $\langle e^{-\beta W} \rangle$ (blue curve) as a function of the number of terms N_{\max} included in the summation (11). This function quickly converges to $e^{-\beta \Delta F}$ proving that the Jarzynski equality (12) holds. On the right panel we have depicted the irreversible work $\langle W_{\text{irr}} \rangle = \langle W \rangle - \Delta F$ (blue curve) as a function of τ which determines the speed at which the energy is supplied to the system. When $\tau \rightarrow \infty$ the system enters its quasistatic regime and the irreversible work becomes negligible, that is $\langle W_{\text{irr}} \rangle \rightarrow 0$ ^{47,48}. The inset (red curve) shows the irreversible work calculated for a linear protocol, $\omega(t) = \omega_i + (\omega_f - \omega_i)t/\tau$. As we can see, it takes longer for the system to reach its quasistatic regime. Moreover, the oscillatory behavior is a signature of the initial excitation which dominates for fast quenches (small τ).

Example 1b. Another class of systems that is used to explain localization effects relates to non-hermitian tight-binding models^{49,50}. For example

$$H = -\frac{t}{2} \sum_{\mathbf{x}} \sum_{\nu=1}^d e^{\alpha \cdot \mathbf{e}_\nu} a_{\mathbf{x}+\mathbf{e}_\nu}^\dagger a_{\mathbf{x}} + e^{-\alpha \cdot \mathbf{e}_\nu} a_{\mathbf{x}}^\dagger a_{\mathbf{x}+\mathbf{e}_\nu} + \sum_{\mathbf{x}} V_{\mathbf{x}} a_{\mathbf{x}}^\dagger a_{\mathbf{x}}, \quad (25)$$

where, $a_{\mathbf{x}}^\dagger$ and $a_{\mathbf{x}}$ are bosonic creation and annihilation operators respectively, \mathbf{e}_ν are the unit lattice vectors, and t is the hopping parameter, and $V_{\mathbf{x}}$ denotes the on-site potential. Interestingly, the complex eigenvectors appear in conjugate pairs (see Eq. (2) in Ref. 41 and the discussion that follows). Therefore, this model provides another example for a building block of a non-hermitian Carnot engine.

Example 2. The remainder of the present work is dedicated to a careful study of a second, experimentally relevant example⁶. Consider a two level system described by the Hamiltonian




Figure 2. Left panel: Relaxation time $T_r = |E_1^\tau - E_2^\tau|^{-1}$, as a function of the final value λ_f for the linear quench $\lambda_t = \lambda_i + (\lambda_f - \lambda_i)t/\tau$. Parameters are $\lambda_i = 0$, $\beta = \hbar = \tau = 1$. Inset: numerical confirmation of the Jarzynski equality (12). Right panel: In the broken regime quantum work can no longer be determined by the two-time energy measurement as $\langle \psi, g\psi \rangle$ can be both positive and negative. To construct the plot we set $g = \sigma_x$. States $\psi(n)$ have been chosen randomly; and n is an integer that has been assigned to them.

$$H_t = \lambda_t \sigma_+ \sigma_- + \lambda_t^* \sigma_- \sigma_+ + \gamma \sigma_+ + \gamma^* \sigma_- \quad (26)$$

where λ_t is a complex control parameter, and γ is a complex constant, whereas σ_+ and σ_- are the raising and lowering fermionic operators. This simple model (26) has been extensively studied in the literature^{11,51,52}, and it has been also realized experimentally both in optics⁴ and semiconductor microcavities⁶.

To make the spectrum of (26) real we set λ_t to be purely imaginary ($\lambda_t \rightarrow i\lambda_t$); and without any loss of generality we choose $\gamma = 1$. This corresponds to the following parameters $E_{1,2} = 0$, $\Gamma_{1,2} = \pm \lambda_t$, and $q = \gamma = 1$ for the hybrid light-matter system of quasiparticles investigated in Ref. 6. Such systems are formed as a result of a strong interaction between excitons and photons in a semiconductor microcavity⁵³. They are commonly referred to as exciton-polaritons⁵⁴.

A simple calculation shows that $H_t^\dagger = \sigma_x H_t \sigma_x$, where σ_x is the Pauli matrix in x direction. Thus H_t is indeed pseudo-hermitian. However, the corresponding σ_x is not a metric. For instance $\langle e_1, \sigma_x e_1 \rangle = 0$, where $|e_1\rangle = (1, 0)^t$. Nevertheless, we can easily find one by rewriting H_t in its diagonal form,

$$V_t^{-1} H_t V_t = \begin{pmatrix} E_1^t & 0 \\ 0 & E_2^t \end{pmatrix}, \quad E_{1,2}^t = \pm \sqrt{1 - \lambda_t^2}. \quad (27)$$

Note, both $E_{1,2}^t$ are real as long as $\lambda_t \leq 1$, otherwise $E_1^t = E_2^{t*}$. Therefore, the Carnot bound (13) holds in both these regimes, whereas the Jarzynski equality (12) only in the first one. Now, the proper metric can be defined via the similarity transformation V_t

$$g_t = V_t^\dagger V_t = 2 \begin{pmatrix} 1 & -i\lambda_t \\ i\lambda_t & 1 \end{pmatrix}. \quad (28)$$

To investigate the dynamics of (26) we assume that λ_t changes on a time scale τ in a linear manner, that is $\lambda_t = \lambda_i + (\lambda_f - \lambda_i)t/\tau$. The linearity does not pose any restriction on our analysis as the Jarzynski equality holds for all protocols λ_t ¹¹. Figure 2 (Left panel) depicts the relaxation time $T_r = \Delta^{-1}$, where $\Delta = |E_1^\tau - E_2^\tau|$, as a function of the final value λ_f ⁵⁵. The relaxation time diverges as λ_f approaches the critical point at $\lambda = 1$. Similar behavior has been observed for the irreversible work $\langle W_{\text{irr}} \rangle := \langle W \rangle - \Delta F$ in \mathcal{PT} -symmetric systems¹¹. The critical point separates the unbroken domain, where energies are real, from the broken one characterized by complex energy values. The energetic cost associated with a potential crossover between those two regimes becomes infinite, and the system “freezes out” before even having a chance to cross to the other regime^{56,57}.

In the broken regime, Eq. (28) no longer reflects pseudo-hermiticity of the system, that is V_t does not fulfill Eq. (4). In fact, all operators g for which the latter equation is true, σ_x being an example (see Fig. 2, Right panel), lead to indefinite inner product spaces. Note that in Fig. 2 (Right panel) the norm can be both positive and negative. Therefore, the evolution within those spaces cannot be unitary and the two-time energy measurement paradigm can no longer be applied⁵⁸. In the quasistatic limit, however, quantum work can still be defined, and we have shown that the second law still holds for all pseudo-hermitian systems.

Conclusions

In summary, we have carefully studied thermodynamic properties of quantum systems that do not satisfy one of the basic requirements imposed on them by the axiom of quantum mechanics - hermiticity. We have shown that if quantum work can be determined by the two-time projective energy measurements, then the Jarzynski equality still holds for non-hermitian systems with real spectrum. Note, this equality expresses the second law of thermodynamics for isothermal processes arbitrarily far from equilibrium.

We have also argued that the Carnot bound is attained for all pseudo-hermitian systems in the quasistatic limit. Furthermore, we have also proposed an experimental setup to test our predictions. As elaborated in the previous section, the system in question consists of strongly interacting excitons and photons in a semiconductor microcavity⁶. Moreover, we have investigated two non-hermitian models that were originally introduced to explain localization effects in solid state physics⁴¹. The first one, a non-hermitian harmonic oscillator that admits real spectrum was used to demonstrate the Jarzynski equality. The second one, the so called non-hermitian tight-binding model was given as an example of a quantum system having complex eigenenergies that appear in conjugate pairs. This model provides another example of a building block of a non-hermitian Carnot engine.

References

- Longhi, S. Optical realization of relativistic non-hermitian quantum mechanics. *Phys. Rev. Lett.* **105**, 013903 (2010).
- Dirac, P. A. M. A new notation for quantum mechanics. *Math. Proc. Cambridge Philos. Soc.* **35**, 416 (1939).
- von Neumann, J. *Mathematical Foundations of Quantum Mechanics* (Princeton University Press, 1955).
- Rüter, E. C. *et al.* Observation of parity-time symmetry in optics. *Nat. Phys.* **6**, 192 (2010).
- Bender, C. M. & Boettcher, S. Real spectra in non-hermitian hamiltonians having \mathcal{PT} symmetry. *Phys. Rev. Lett.* **80**, 5243 (1998).
- Gao, T. *et al.* Observation of non-hermitian degeneracies in a chaotic exciton-polariton billiard. *Nature* **526**, 554558 (2015).
- Meng-Jun Hu, X.-M. H. & Zhang, Y.-S. Are observables necessarily Hermitian? Preprint at arXiv:1601.04287v1 (2015).
- Moiseyev, N. *Non-Hermitian Quantum Mechanics* (Cambridge University Press, 2011).
- Berry, M. V. Optical polarization evolution near a non-hermitian degeneracy. *J. Opt.* **13**, 115701 (2011).
- Brody, D. C. Consistency of \mathcal{PT} -symmetric quantum mechanics. Preprint at arXiv:1508.02190 (2015).
- Deffner, S. & Saxena, A. Jarzynski equality in \mathcal{PT} -symmetric quantum mechanics. *Phys. Rev. Lett.* **114**, 150601 (2015).
- Jarzynski, C. Nonequilibrium equality for free energy differences. *Phys. Rev. Lett.* **78**, 2690 (1997).
- Crooks, G. E. Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences. *Phys. Rev. E* **60**, 2721 (1999).
- Ortiz de Zárate, J. M. Interview with Michael E. Fisher. *Europhys. News* **42**, 14 (2011).
- Mostafazadeh, A. Pseudo-hermiticity versus \mathcal{PT} -symmetry III: Equivalence of pseudo-hermiticity and the presence of antilinear symmetries. *J. Math. Phys.* **43**, 3944 (2002).
- Brody, D. C. Biorthogonal quantum mechanics. *J. Phys. A: Math. Theor.* **47**, 035305 (2014).
- Rivas, A. & Huelga, S. F. *Open Quantum Systems: An Introduction* (SpringerBriefs in Physics, 2012).
- Alicki, R. Pure decoherence in quantum systems. *Open Syst. Inf. Dyn.* **11**, 53 (2004).
- An, S. *et al.* Experimental test of the quantum Jarzynski equality with a trapped-ion system. *Nat. Phys.* **11**, 193199 (2015).
- Kafri, D. & Deffner, S. Holevo's bound from a general quantum fluctuation theorem. *Phys. Rev. A* **86**, 044302 (2012).
- Rastegin, A. E. Non-equilibrium equalities with unital quantum channels. *J. Stat. Mech. Theor. Exp.* **2013**, P06016 (2013).
- Mostafazadeh, A. Time-dependent pseudo-hermitian hamiltonians defining a unitary quantum system and uniqueness of the metric operator. *Phys. Lett. B* **650**, 208 (2007).
- Cho, J.-H. Understanding the complex position in a \mathcal{PT} -symmetric oscillator. Preprint at arXiv:1509.03653 (2015).
- Van den Broeck, C. & Toral, R. Stochastic thermodynamics for linear kinetic equations. *Phys. Rev. E* **92**, 012127 (2015).
- Yeo, J., Kwon, C., Lee, H. K. & Park, H. Housekeeping entropy in continuous stochastic dynamics with odd-parity variables. Preprint at arXiv:1511.04353 (2015).
- Znojil, M. Non-hermitian Heisenberg representation. *Phys. Lett. A* **379**, 2013 (2015).
- Gong, J. & hai Wang, Q. Time-dependent \mathcal{PT} -symmetric quantum mechanics. *J. Phys. A: Math. Theor.* **46**, 485302 (2013).
- Thiffeault, J.-L. Covariant time derivatives for dynamical systems. *J. Phys. A: Math. Gen.* **34**, 5875 (2001).
- Aharonov, Y., Massar, S. & Popescu, S. Measuring energy, estimating Hamiltonians, and the time-energy uncertainty relation. *Phys. Rev. A* **66**, 052107 (2002).
- Leonard, A. & Deffner, S. Quantum work distribution for a driven diatomic molecule. *Chem. Phys.* **446**, 18 (2015).
- Kafri, D. & Deffner, S. Holevo's bound from a general quantum fluctuation theorem. *Phys. Rev. A* **86**, 044302 (2012).
- Lewenstein, M. Quantum mechanics: No more fields. *Nat. Phys.* **11**, 211 (2015).
- Gardas, B. & Deffner, S. Thermodynamic universality of quantum Carnot engines. *Phys. Rev. E* **92**, 042126 (2015).
- Xiao, G. & Gong, J. Construction and optimization of a quantum analog of the Carnot cycle. *Phys. Rev. E* **92**, 012118 (2015).
- Carnot, S. *Réflexions sur la Puissance Motrice de feu et sur les Machines Propres à développer Cette Puissance* (Gauthier-Villars, 1824).
- Huang, X. L., Wang, T. & Yi, X. X. Effects of reservoir squeezing on quantum systems and work extraction. *Phys. Rev. E* **86**, 051105 (2012).
- Long, R. & Liu, W. Performance of quantum Otto refrigerators with squeezing. *Phys. Rev. E* **91**, 062137 (2015).
- Geusic, J. E., Schulz-DuBios, E. O. & Scovil, H. E. D. Quantum equivalent of the Carnot cycle. *Phys. Rev.* **156**, 343 (1967).
- Roßnagel, J., Abah, O., Schmidt-Kaler, F., Singer, K. & Lutz, E. Nanoscale heat engine beyond the Carnot limit. *Phys. Rev. Lett.* **112**, 030602 (2014).
- Scully, M. O., Zubairy, M. S., Agarwal, G. S. & Walther, H. Extracting work from a single heat bath via vanishing quantum coherence. *Science* **299**, 862 (2003).
- Hatano, N. & Nelson, D. R. Localization transitions in non-hermitian quantum mechanics. *Phys. Rev. Lett.* **77**, 570 (1996).
- Parrondo, J. M. R., Horowitz, J. M. & Sagawa, T. Thermodynamics of information. *Nat. Phys.* **11**, 131 (2015).
- Callen, H. B. *Thermodynamics and an Introduction to Thermostatistics* (John Wiley & Sons, 1985).
- Alicki, R. The quantum open system as a model of the heat engine. *J. Phys. A: Math. Theor.* **12**, L103 (1979).
- Ulm, S. *et al.* Observation of the Kibble-Zurek scaling law for defect formation in ion crystals. *Nat. Commun.* **4**, 2290 (2013).
- Jingbo, X. & Youhong, Y. Time evolution of the time-dependent harmonic oscillator. *Commun. in Theor. Phys.* **29**, 385 (1998).
- Deffner, S. & Lutz, E. Nonequilibrium work distribution of a quantum harmonic oscillator. *Phys. Rev. E* **77**, 021128 (2008).
- Galve, F. & Lutz, E. Nonequilibrium thermodynamic analysis of squeezing. *Phys. Rev. A* **79**, 055804 (2009).
- Longhi, S. Invisibility in non-hermitian tight-binding lattices. *Phys. Rev. A* **82**, 032111 (2010).
- Longhi, S. Convective and absolute \mathcal{PT} -symmetry breaking in tight-binding lattices. *Phys. Rev. A* **88**, 052102 (2013).
- Bender, C. M., Brody, D. C. & Jones, H. F. Complex extension of quantum mechanics. *Phys. Rev. Lett.* **89**, 270401 (2002).
- Bender, C. M., Brody, D. C. & Jones, H. F. Must a hamiltonian be hermitian? *Am. J. Phys.* **71**, 1095 (2003).
- Carusotto, I. & Ciuti, C. Quantum fluids of light. *Rev. Mod. Phys.* **85**, 299 (2013).
- Deng, H., Haug, H. & Yamamoto, Y. Exciton-polariton Bose-Einstein condensation. *Rev. Mod. Phys.* **82**, 1489 (2010).
- Eisert, J., Friesdorf, M. & Gogolin, C. Quantum many-body systems out of equilibrium. *Nat. Phys.* **11**, 124 (2015).
- Kibble, T. W. B. Topology of cosmic domains and strings. *J. Phys. A: Math. Gen.* **9**, 1387 (1976).
- Zurek, W. H. Cosmological experiments in superfluid helium? *Nature* **317**, 505 (1985).
- Albash, T., Lidar, D. A., Marvian, M. & Zanardi, P. Fluctuation theorems for quantum processes. *Phys. Rev. E* **88**, 032146 (2013).

Acknowledgements

This work was supported by the Polish Ministry of Science and Higher Education under project Mobility Plus 1060/MOB/2013/0 (B.G.); S.D. acknowledges financial support from the U.S. Department of Energy through a LANL Director's Funded Fellowship.

Author Contributions

B.G., S.D. and A.S. developed ideas and derived the main results. B.G. prepared Figures 1 and 2. B.G., S.D. and A.S. wrote and reviewed the manuscript.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Gardas, B. *et al.* Non-hermitian quantum thermodynamics. *Sci. Rep.* **6**, 23408; doi: 10.1038/srep23408 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>