Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) <https://creativecommons.org/licenses/by-nc-nd/4.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

ITRF2020: The International Laser Ranging Service (ILRS) Contribution

E. C. Pavlis, M. Kuzmich-Cieslak, K. Evans
JCET/UMBC – Baltimore, MD

V. Luceri, A. Basoni, D. Sarrocco
e-GEOS S.p.A., ASI/CGS - Matera

G. Bianco
Agenzia Spaziale Italiana, CGS - Matera

Based on the ITRF2020P results, the new ILRS contribution limits the systematic scale difference **VLBI - SLR** to ≈ 0.22 ppb (~ 1.4 mm) ☀

☀ Zuheir Altamimi, IERS DB#73

- The ILRS ASC established a new analysis approach for its contribution to ITRF2020;
- It will be implemented in the operational series after adoption of ITRF2020;
- The complete SLR series for the **38-year** period **1983 – 2021** was submitted to ITRS in early November 2021;
- Based on the initial ITRF2020P combination ITRS has corroborated the ILRS results on the scale difference between SLR & VLBI.

Contact us at:

Cinzia Luceri

cinzia.luceri@e-geos.it

Erricos C. Pavlis

epavlis@umbc.edu