

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>


Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Concise Communication

Documentation of acute change in mental status in nursing homes highlights opportunity to augment infection surveillance criteria

Austin R. Penna MPH¹ , Christina L. Sancken MPH¹, Nimalie D. Stone MD¹, Taniece R. Eure MPH¹, Wendy Bamberg MD², Grant Barney MPH³, Devra Barter MS², Stacy Carswell MPH⁴, Paula Clogher MPH⁵, Ghinwa Dumyati MD^{3,6}, Christina B. Felsen MPH^{3,6}, Linda Frank RN⁷, Deborah Godine RN⁷, Helen Johnston MPH², Marion A. Kainer MBBS⁸, Linda Li MPH⁹, Ruth Lynfield MD¹⁰, J. P. Mahoehey RN¹⁰, Joelle Nadle MPH⁷, Rebecca Pierce PhD¹¹, Susan M. Ray MD^{4,12}, Sarah Shrum Davis MPH¹³, Marla Sievers MPH¹³, Lucy E. Wilson MD⁹, Alexia Y. Zhang MPH¹¹, Shelley S. Magill MD¹ and Nicola D. Thompson PhD¹

¹Division of Healthcare Quality Promotion, Centers for Disease Control and Prevention, Atlanta, Georgia, ²Colorado Department of Public Health and Environment, Denver, Colorado, ³New York Emerging Infections Program, Rochester, New York, ⁴Georgia Emerging Infections Program, Atlanta, Georgia, ⁵Connecticut Emerging Infections Program and the Yale School of Public Health, New Haven, Connecticut, ⁶University of Rochester Medical Center, Rochester, New York, ⁷California Emerging Infections Program, Oakland, California, ⁸Tennessee Department of Health, Nashville, Tennessee, ⁹Maryland Department of Health, Baltimore, Maryland, ¹⁰Minnesota Department of Health, St Paul, Minnesota, ¹¹Oregon Health Authority, Portland, Oregon, ¹²Emory University, Atlanta, Georgia and ¹³New Mexico Department of Health, Santa Fe, New Mexico

Abstract

Acute change in mental status (ACMS), defined by the Confusion Assessment Method, is used to identify infections in nursing home residents. A medical record review revealed that none of 15,276 residents had an ACMS documented. Using the revised McGeer criteria with a possible ACMS definition, we identified 296 residents and 21 additional infections. The use of a possible ACMS definition should be considered for retrospective nursing home infection surveillance.

(Received 3 January 2020; accepted 22 February 2020; electronically published 28 April 2020)

Delirium is a serious condition in older adults in nursing homes; it leads to higher risk for adverse health outcomes and increased healthcare costs.^{1,2} Although its clinical features may go unrecognized, factors that contribute to delirium include systemic infection, fever, dehydration, adverse reaction to medications, and other forms of acute illness.¹ Various tools for delirium assessment exist; however, the validated Confusion Assessment Method (CAM)³ and its adaptations are among the most widely used.⁴ The CAM approach to assessing delirium is approximated in the CMS minimum dataset (MDS) version 3.0, and it is used in the revised McGeer Criteria for healthcare-associated infection (HAI) surveillance in nursing homes to define ACMS as a constitutional criterion.⁵

At its core, the CAM assesses the presence of confirmed delirium using 4 elements: (1) acute onset with fluctuating course; (2) inattention; and either (3) disorganized thinking or (4) altered level of consciousness.³ Despite this standardized algorithm,³ nursing staff often lack formal education or training in the recognition of

CAM elements,^{6,7} or they may fail to appropriately document resident symptoms.⁸ Given these challenges, we assessed the documentation of the 4 CAM elements in a large sample of US nursing home residents, and we evaluated the impact of using a more sensitive, modified ACMS criterion in infection surveillance definitions.

Methods

In 2017, the Centers for Disease Control and Prevention (CDC) Emerging Infections Program (EIP) conducted a nursing home prevalence survey of HAIs in 10 states: California, Colorado, Connecticut, Georgia, Maryland, Minnesota, New Mexico, New York, Oregon, and Tennessee.⁹ Trained EIP staff reviewed resident medical records for any possible signs or symptoms of infection present on the survey date or the day prior. If present, a more detailed review of documentation was conducted to identify all infection criteria present, including for any ACMS that were new or different from baseline, on the survey date and in the 6 days prior. The revised McGeer Criteria were used to identify HAIs and included the CAM-defined constitutional criterion, confirmed ACMS (ie, documentation of acute onset with fluctuating course, inattention, and either disorganized thinking or altered level of consciousness), alongside specific signs or symptoms in the following infection surveillance definitions: catheter-associated urinary tract infection (CAUTI), lower respiratory tract infection (LRTI), pneumonia (PNEU), and cellulitis and/or soft tissue and/or wound

Author for correspondence: Austin R. Penna, E-mail: apenna@cdc.gov

PREVIOUS PRESENTATION: Selected data included in this manuscript were presented in an abstract at the Society for Healthcare Epidemiology of America Spring Conference, April 26, 2019, in Boston, Massachusetts.

Cite this article: Penna AR, *et al.* (2020). Documentation of acute change in mental status in nursing homes highlights opportunity to augment infection surveillance criteria. *Infection Control & Hospital Epidemiology*, 41: 848–850, <https://doi.org/10.1017/ice.2020.77>

Table 1. Confusion Assessment Method (CAM) Element Documentation Among Nursing Home Residents with Possible Acute Change in Mental Status (ACMS) (N = 296)

Documented CAM Element Combinations ^a	No.	%
Disorganized thinking	127	42.9
Altered level of consciousness	81	27.3
Fluctuating behavior	37	12.5
Disorganized thinking and altered level of consciousness	20	6.7
Fluctuating behavior and disorganized thinking	8	2.7
Fluctuating behavior and altered level of consciousness	7	2.4
Fluctuating behavior and disorganized thinking and altered level of consciousness	7	2.4
Inattention	5	1.7
Inattention and disorganized thinking	2	0.7
Inattention and altered level of consciousness	2	0.7

^aElement combinations are mutually exclusive.

infection (SSTI).⁵ Standardized forms and written guidance for these definitions, including confirmed ACMS, were provided to EIP staff to aid data collection.

We calculated the prevalence of each CAM element, combinations of elements, and confirmed ACMS. We defined possible ACMS as the documentation of any of the 4 CAM elements. The number of residents meeting CAUTI, LRTI, PNEU, and SSTI definitions using the confirmed and possible ACMS constitutional criteria were calculated and compared. The χ^2 test and Wilcoxon's rank-sum test were used to compare differences between residents with possible ACMS and those with no CAM elements documented for selected characteristics. *P* values <.05 were considered statistically significant. All data were analyzed using SAS version 9.4 software (SAS Institute, Cary, NC).

Results

Among 15,276 eligible residents in 161 participating nursing homes, no resident had the required documentation to meet the confirmed ACMS criterion. The most frequently documented CAM element was disorganized thinking (*n* = 164, 1.1%), followed by altered level of consciousness (*n* = 117, 0.8%), fluctuating behavior (*n* = 59, 0.4%), and inattention (*n* = 9, 0.1%). In total, 296 (1.9%) residents had at least 1 CAM element documented, meeting the possible ACMS criterion. Disorganized thinking and altered level of consciousness were the most common elements in combination (Table 1). Residents meeting the possible ACMS criterion were older than those with no CAM elements documented: median age, 82 years (interquartile range [IQR], 14) versus 80 years (IQR, 19; *P* = .009). They also had shorter nursing home stay duration than those with no CAM elements documented: median days from nursing home admission date to survey date: 132 days (IQR, 585) versus 334 days (IQR, 891; *P* < .0001). Compared with confirmed ACMS, the use of the possible ACMS criterion resulted in the detection of 21 additional HAIs (5.6% overall increase) (Fig 1.). The greatest increases were for LRTI (*n* = 9, 37.5%) and PNEU (*n* = 8, 29.6%).

Discussion

Among a large number of nursing home residents, no resident had the required documentation to meet the revised McGeer

confirmed ACMS criterion, as defined using the CAM. This finding was surprising given that other long-term care and post-acute care settings have reported estimates of CAM-defined delirium ranging from 16% to 60%.^{1,2,6} Documentation of the individual core CAM elements was also lower than expected when compared with a study of CAM documentation in Canadian nursing homes, where the frequency of delirium symptoms ranged from 1.9% for inattention to 13.0% for altered level of consciousness.⁸ Notably, inattention was also the least frequently documented element (0.1%) in our survey. Because inattention is a required element in the CAM-based delirium definition, its absence had a large effect on the prevalence of residents meeting the confirmed ACMS criterion.

Although it is unclear why so few of the CAM elements were documented in our survey, explanations proposed in the literature include high resident-to-nurse staffing ratios, a preference of nursing staff to provide verbal reports of resident status, and time constraints for completing documentation.⁶⁻⁸ One study comparing delirium assessment between trained researchers and untrained clinical staff showed that due to the complexity of bedside assessments and the presence of resident factors such as dementia, specialized training is required for nursing home staff to adequately assess delirium.⁶ Additional explanations may include failure of clinical staff to perform formal CAM assessments or failure of staff to appreciate the severity of delirium symptoms.^{7,8}

When creating the possible ACMS criteria, we evaluated all combinations of the CAM elements and ultimately selected the most sensitive—any 1 of 4 CAM elements. Our proposal for use of a possible ACMS criterion in infection surveillance is supported by the findings of Inouye et al,¹⁰ who validated a chart-based delirium detection method among older persons in acute care settings. In their study, chart documentation of any key descriptive CAM term yielded good sensitivity (74%) and specificity (83%) compared to direct delirium assessment using the formal CAM. These authors recommended use of this more sensitive, chart-based method for patient safety or quality improvement initiatives but not for clinical assessment or establishing a diagnosis of delirium.¹⁰ In a separate study, chart-based detection was also recommended for use in research settings to augment formal assessments and to identify additional persons with possible delirium.¹¹

Using a possible ACMS criterion, the prevalence of ACMS in our survey increased from 0 to 1.9%. This change resulted in the identification of 21 additional HAIs (5.6% increase in HAIs), and most were respiratory infections (*n* = 17, 81.0%). Because other diagnostic or localizing criteria must also be present to meet the revised McGeer infection surveillance definitions,⁵ misclassifications of residents with HAIs were not anticipated. We also found that residents meeting the possible ACMS criterion were more recently admitted to nursing homes and were significantly older than those with no CAM elements documented. Characteristics associated with the prevalence of delirium symptoms in nursing home residents include recent hospitalization and increasing age.^{1,2} This finding increases our confidence that the possible ACMS criterion identifies a resident population of interest for delirium symptoms.

Notably, the assessment of ACMS documentation was not the primary objective of the prevalence survey, and our findings may be limited by incomplete medical record documentation in nursing homes.⁸ However, given the need for specialized CAM assessment training and evidence of suboptimal documentation of the core CAM elements in nursing homes,⁶⁻⁸ our findings suggest that use of the CAM to define confirmed ACMS in infection surveillance definitions is likely too stringent. Although investigators have




Fig 1. Number and percent increase in cellulitis and/or soft-tissue and/or wound (SSTI), catheter-associated urinary tract (CAUTI), pneumonia (PNEU), and lower respiratory tract (LRTI) healthcare-associated infections (HAIs) identified among nursing home residents using the confirmed acute change in mental status (ACMS) and possible ACMS criteria.

emphasized the importance of formal CAM assessments to clinically diagnose delirium,¹⁰ a possible ACMS criterion may be better suited for retrospective infection surveillance. Additional research to assess barriers associated with incorporating CAM assessment and documentation practices in nursing homes is needed when evaluating residents suspected of infection to improve infection surveillance and resident safety.

Acknowledgments. We thank the staff and nursing homes that participated in the 2017 Centers for Disease Control and Prevention (CDC)/Emerging Infections Program (EIP) Nursing Home Point Prevalence Survey. We also thank EIP and CDC staff for their contributions to survey coordination, data collection, data entry, and data management. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position of the Centers for Disease Control and Prevention /the Agency for Toxic Substances and Disease Registry, or the Department of Veterans' Affairs.

Financial support. This work was supported through a cooperative agreement by the Centers for Disease Control and Prevention (CDC) and the Emerging Infections Program (EIP).

Conflicts of interest. M.A.K. reports nonfinancial support from Council of State and Territorial Epidemiologists (CSTE), nonfinancial support from Society for Healthcare Epidemiology of America (SHEA), personal fees and nonfinancial support from Infectious Disease Consulting Corporation (IDCC), personal fees and nonfinancial support from WebMD, nonfinancial support from Association for Professionals in Infection Control and Epidemiology (APIC), from American Society for Microbiology, personal fees and nonfinancial support from Pfizer, outside the submitted work. R.L. reports being a coeditor for a book on surveillance and an associate editor for the American Academy of Pediatrics *Red Book* ("Report of the Committee on Infectious Diseases"); royalties from the textbook and compensation from *Red Book* have been donated to the Minnesota Department of Health.

References

1. Inouye SK. Delirium in older persons. *N Engl J Med* 2006;354:1157–1165.
2. Kiely DK, Bergmann MA, Murphy KM, Jones RN, Orav EJ, Marcantonio ER. Delirium among newly admitted postacute facility patients: prevalence, symptoms, and severity. *J Gerontol A Biol Sci Med Sci* 2003;58:441–445.
3. Inouye SK, van Dyck CH, Alessi CA, Balkin S, Siegel AP, Horwitz RI. Clarifying confusion: the confusion assessment method. A new method for detection of delirium. *Ann Intern Med* 1990;113:941–948.
4. Wei LA, Fearing MA, Sternberg EJ, Inouye SK. The Confusion Assessment Method: a systematic review of current usage. *J Am Geriatr Soc* 2008;56:823–830.
5. Stone ND, Ashraf MS, Calder J, *et al*. Surveillance definitions of infections in long-term care facilities: revisiting the McGeer criteria. *Infect Control Hosp Epidemiol* 2012;33:965–977.
6. Voyer P, Richard S, McCusker J, *et al*. Detection of delirium and its symptoms by nurses working in a long term care facility. *J Am Med Dir Assoc* 2012;13:264–271.
7. Inouye SK, Foreman MD, Mion LC, Katz KH, Cooney LM Jr. Nurses' recognition of delirium and its symptoms: comparison of nurse and researcher ratings. *Arch Intern Med* 2001;161:2467–2473.
8. Voyer P, McCusker J, Cole MG, *et al*. Nursing documentation in long-term care settings: new empirical evidence demands changes be made. *Clin Nurs Res* 2014;23:442–461.
9. Thompson ND, Penna A, Eure TR, *et al*. Epidemiology of antibiotic use for urinary tract infection in nursing home residents. *J Am Med Dir Assoc* 2020;21:91–96.
10. Inouye SK, Leo-Summers L, Zhang Y, Bogardus ST Jr, Leslie DL, Agostini JV. A chart-based method for identification of delirium: validation compared with interviewer ratings using the confusion assessment method. *J Am Geriatr Soc* 2005;53:312–318.
11. Saczynski JS, Kosar CM, Xu G, *et al*. A tale of two methods: chart and interview methods for identifying delirium. *J Am Geriatr Soc* 2014;62:518–524.