

This is a pre-copyedited, author-produced PDF of an article accepted for publication in Monthly Notices of the Royal Astronomical Society following peer review. The version of record E Aydi and others, Catching a nova X-ray/UV flash in the visible? Early spectroscopy of the very slow Nova Velorum 2022 (Gaia22alz), Monthly Notices of the Royal Astronomical Society, 2023;, stad1914, <https://doi.org/10.1093/mnras/stad1914> is available online at: <https://doi-org.proxy-bc.researchport.umd.edu/10.1093/mnras/stad1914>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

**Please provide feedback**

Please support the ScholarWorks@UMBC repository by emailing [scholarworks-group@umbc.edu](mailto:scholarworks-group@umbc.edu) and telling us what having access to this work means to you and why it's important to you. Thank you.

# Catching a nova X-ray/UV flash in the visible? Early spectroscopy of the very slow Nova Velorum 2022 (Gaia22alz)

E. Aydi<sup>1</sup>,<sup>1</sup>★† L. Chomiuk,<sup>1</sup> J. Mikołajewska,<sup>2</sup> J. Brink,<sup>3,4</sup> B. D. Metzger,<sup>5,6</sup> J. Strader,<sup>1</sup> D.A.H. Buckley<sup>3,4</sup>, E.J. Harvey<sup>7</sup>, T.W.-S. Holoien<sup>8</sup>, L. Izzo,<sup>9</sup> A. Kawash,<sup>1</sup> J.D. Linford<sup>10</sup>, P. Molaro<sup>11,12</sup>, I. Molina,<sup>1</sup> P. Mróz,<sup>13</sup> K. Mukai<sup>14,15</sup>, M. Orío<sup>16,17</sup>, T. Panurach,<sup>1</sup> P. Senchyna<sup>8</sup>, B. J. Shappee,<sup>18</sup> K. J. Shen,<sup>19</sup> J. L. Sokoloski,<sup>20</sup> K. V. Sokolovsky<sup>21,22</sup>, R. Urquhart<sup>1</sup> and R. E. Williams<sup>23,24</sup>

*Affiliations are listed at the end of the paper*

Accepted 2023 June 21. Received 2023 June 20; in original form 2023 April 12

## ABSTRACT

We present early spectral observations of the very slow Galactic nova Gaia22alz, over its gradual rise to peak brightness that lasted 180 d. During the first 50 d, when the nova was only 3–4 mag above its normal brightness, the spectra showed narrow (FWHM  $\approx 400 \text{ km s}^{-1}$ ) emission lines of H Balmer, HeI, HeII, and CIV but no P Cygni absorption. A few weeks later, the high-excitation HeII and CIV lines disappeared, and P Cygni profiles of Balmer, HeI, and eventually FeII lines emerged, yielding a spectrum typical of classical novae before peak. We propose that the early (first 50 d) spectra of Gaia22alz, particularly the emission lines with no P Cygni profiles, are produced in the white dwarf’s optically thin envelope or accretion disc, reprocessing ultraviolet and potentially X-ray emission from the white dwarf after a dramatic increase in the rate of thermonuclear reactions, during a phase known as the ‘early X-ray/UV flash’. If true, this would be one of the rare times that the optical signature of the early X-ray/UV flash has been detected. While this phase might last only a few hours in other novae and thus be easily missed, it was possible to detect in Gaia22alz due to its very slow and gradual rise and thanks to the efficiency of new all-sky surveys in detecting transients on their rise. We also consider alternative scenarios that could explain the early spectral features of Gaia22alz and its gradual rise.

**Key words:** transients: novae – stars: white dwarfs.

## 1 INTRODUCTION

Classical novae are thermonuclear runaway events taking place on the surface of accreting white dwarfs in interacting binary star systems (for reviews, see e.g. Bode & Evans 2008; Della Valle & Izzo 2020; Chomiuk, Metzger & Shen 2021). The eruption leads to an increase in the visible brightness of the system by 8–15 mag over a few hours to days. While most classical novae rapidly ascend to peak brightness, some exhibit a prolonged brightening lasting for several weeks (e.g. Chochol et al. 2003; Strope, Schaefer & Henden 2010; Tanaka et al. 2011a, b; Aydi et al. 2019, 2020a).

During the rise to peak, the spectra of novae are typically dominated by P Cygni profiles of Balmer, FeII, HeI, OI, and NaI, characterized by velocities ranging between a few hundreds to  $\approx 1000 \text{ km s}^{-1}$ . After peak, faster emission components emerge with velocities of a few  $1000 \text{ km s}^{-1}$  (McLaughlin 1944; McLaughlin 1947; Gaposchkin 1957; Aydi et al. 2020b). It is challenging to observe novae before they reach optical peak, mainly due to their

typically quick rise and their transient nature. However, slowly rising novae give us a chance to monitor them during the days – weeks leading to their visible peak. Nevertheless, even the slowest novae are discovered when they are already several magnitudes ( $>5 \text{ mag}$ ) above quiescent brightness (e.g. Hounsell et al. 2010; Strope et al. 2010; Hounsell et al. 2016), with the exception of just a few (e.g. HR Del; Friedjung 1992 and T Pyx; Arai et al. 2015). Therefore, the early times of the eruption are often missed, and spectral observations during these early phases are scarce.

Gaia22alz (AT2022bpg; Nova Velorum 2022) was reported by Gaia science alerts<sup>1</sup> on 2022 February 04.25 UT as an optical transient at  $G = 16.87 \text{ mag}$ , towards the constellation Vela (Hodgkin et al. 2022). The transient is located at equatorial coordinates of  $(\alpha, \delta) = (09^{\text{h}}29^{\text{m}}53^{\text{s}}.06, -56^{\circ}17'25''.94)$  and Galactic coordinates of  $(l, b) = (27^{\circ}.563, -3^{\circ}.692)$ . There is a Gaia EDR3 (Gaia Collaboration 2021) source (ID 5307057922488654848) matching the transient position with an average  $G$  magnitude of 18.3. Archival photometry from the All-Sky Automated Survey for SNe (ASAS-SN; Shappee et al. 2014; Kochanek et al. 2017) showed that the eruption started

\* E-mail: aydielia@msu.edu

† NFHP Hubble Fellow.

<sup>1</sup><http://gsaweb.ast.cam.ac.uk/alerts>

as early as 2022 January 25.02 UT (HJD 2459604.522; Aydi et al. 2022b); we will consider this date as  $t_0$  in the remainder of the paper. In archival data from the Dark Energy Camera Plane Survey (DECaPS; Schlafly et al. 2018), the source has median pre-eruption magnitude of  $g \simeq 19.0$ .

Brink et al. (2022) reported spectroscopic observations taken around 45 d after  $t_0$  (still during the early rise to peak), showing narrow emission features of H I Balmer, He I, He II, and C IV. Based on these features and the then-low-amplitude outburst, Brink et al. (2022) suggested that the transient might be an unusual disc instability (dwarf nova) event. Later, observations, taken as the transient climbed to peak, were more consistent with a classical nova eruption, showing typical P Cygni profiles of Balmer, He I, Fe II, and O I (Aydi et al. 2022b, c). This suggests that the early spectra of Gaia22alz are unique, covering a phase of a nova eruption that is usually missed.

Here, we present these early observations of nova Gaia22alz, focusing on its unusual spectral evolution. In Section 2, we present the observations and data reduction. In Section 3, we show our results, while in Section 4, we offer discussion about the nature of the eruption and the origin of the early spectral features. Our conclusions are presented in Section 5.

## 2 OBSERVATIONS AND DATA REDUCTION

We make use of publicly available photometry from the ASAS-SN survey. These data are taken as part of the all-sky patrol in the  $g$ -band. The data can be found in the supplementary online material.

We obtained optical spectroscopic observations of Gaia22alz between d 45 and 278 using a variety of telescopes and instruments. A log of the spectral observations is presented in Table A1. Below, we present details of these observations.

We obtained spectra using the High Resolution Spectrograph (HRS; Barnes et al. 2008; Bramall et al. 2010, 2012; Crause et al. 2014) and the Robert Stobie Spectrograph (RSS; Burgh et al. 2003; Kobulnicky et al. 2003) mounted on the Southern African Large Telescope (SALT; Buckley, Swart & Meiring 2006; O'Donoghue et al. 2006) in Sutherland, South Africa. We used HRS in the low resolution LR mode, yielding a resolving power  $R = \lambda/\Delta\lambda \approx 14\,000$  over the range 4000–9000 Å. The primary reduction of the HRS spectroscopy was conducted using the SALT science pipeline (Crawford et al. 2010), which includes over-scan correction, bias subtraction, and gain correction. The rest of the reduction was done using the MIDAS FEROS (Stahl, Kaufer & Tubbesing 1999) and ECHELLE (Ballester 1992) packages. The reduction procedure is described by Kniazev, Gvaramadze & Berdnikov (2016). RSS was used in long-slit mode with the 1.5 arcsec slit and the PG900 grating, resulting in a resolving power  $R \approx 1500$ . The spectra were first reduced using the PYSALT pipeline (Crawford et al. 2010), which involves bias subtraction, cross-talk correction, scattered light removal, bad pixel masking, and flat-fielding. The wavelength calibration, background subtraction, and spectral extraction were done using the Image Reduction and Analysis Facility (IRAF; Tody 1986).

We also carried out low- and medium-resolution optical spectroscopy using the Goodman spectrograph (Clemens, Crain & Anderson 2004) on the 4.1 m Southern Astrophysical Research (SOAR) telescope located on Cerro Pachón, Chile. The observations were carried out in two setups: one setup using the 400 lmm<sup>-1</sup> grating and a 1.2 arcsec slit, yielding a resolving power  $R \approx 1100$  over the wavelength range 3820–7850 Å. Another setup utilized a 2100 lmm<sup>-1</sup> grating and a 1.2 arcsec slit, yielding a resolving power

$R \approx 4400$  in a 570 Å wide region centred on H $\beta$ . The spectra were reduced and optimally extracted using the APALL package in IRAF.

We also used the Inamori-Magellan Areal Camera & Spectrograph (IMACS) on the Magellan-Baade 6.5 m telescope at Las Campanas Observatory. On the night of 2022 July 21, we obtained a spectrum in longslit mode with the 600 lmm<sup>-1</sup> grating and the f/4 camera through a 0.9 arcsec slit, yielding a spectral resolution of  $R \approx 1500$  over 3800–6600 Å. The spectra were reduced and optimally extracted using the APALL package in IRAF.

## 3 RESULTS

### 3.1 Photometric evolution

In Fig. 1, we present photometry of Gaia22alz from the ASAS-SN survey, focusing on the first 300 d of the eruption – as the focus of this work is on the early spectral evolution of the nova. A complete light curve is presented in Fig. A1, showing the ASAS-SN photometry up until the time of writing of this paper. Solar conjunction prevented ASAS-SN from obtaining photometric observations between d  $\sim$ 185–240 after  $t_0$ . Due to this gap in the monitoring, it is neither possible to definitively constrain the date of optical peak in the light curve nor its magnitude. However, it is possible that the peak was reached around d 177 at  $g \lesssim 11.2$  mag, implying an eruption amplitude of  $\gtrsim 7.8$  mag. This is at the lower end of eruption amplitudes distribution found in novae ( $\sim$ 8–15 mag; Vogt 1990; Warner 2008; Kawash et al. 2021).

The rise of nova Gaia22alz is slow and gradual, taking more than 180 d to reach peak brightness. While other novae have taken a similarly long time to reach peak (even more in some cases; see fig. 6 of Kato & Hachisu 2011), their rises are not as gradual as observed in Gaia22alz; these other eruptions typically start with a rapid rise in brightness by 5–6 mag, then at some point slow down. It takes a typical slow nova several weeks or months to climb the remaining 3–4 mag to its peak (e.g. V723 Ca: Munari et al. 1996; Hachisu & Kato 2004; Goranskij et al. 2007; V1548 Aql: Kato & Takamizawa 2001; V1280 Sco: Hounsell et al. 2016; and the recent example of V1405 Cas).

In contrast, Gaia22alz was detected less than a magnitude above its quiescence level, showing a slow-paced rise to optical peak. While for many slow novae the duration of their initial fast rise are well constrained, it cannot be excluded that some historical slow novae also could have shown behaviour similar to that of Gaia22alz (e.g. HR Del: Strobe et al. 2010). As deep high-cadence all-sky photometric and transient surveys like ASAS-SN and Gaia have become operational only recently, the future is bright for discovering more slow novae at a stage when they are just above their quiescent brightness.

Some time around or during solar conjunction (d  $\sim$ 180–240), the optical light curve of Gaia22alz peaked, and this peak was accompanied by at least one additional maximum/flare (e.g. the secondary maximum on day 255; see Fig. 1). Many novae have been observed to show multiple maxima/flares after peaking (e.g. Pejcha 2009; Strobe et al. 2010; Tanaka et al. 2011a, b; Walter 2016; Shore et al. 2018; Aydi et al. 2019, 2020a), particularly amongst those with slowly evolving light curves. Novae are divided into different speed classes based on the rate of decline of the light curve by 2 mag from peak, known as the  $t_2$  time-scale. We consider  $t_2$  as the last time the light curve dropped 2 mag below peak brightness (Strobe et al. 2010). Assuming that we captured the light-curve peak on d 177, we derive  $t_2 \approx 212$  d (Fig. A1), meaning that Gaia22alz belongs to the

class of very slow novae (Gaposchkin 1957). However, at the time of writing of this paper, the nova was still showing flares, and it might still rise above 2 mag below peak (see Fig. A1).


Photometric measurements taken from the American Association of Variable Star Observers (AAVSO; Kafka 2020) on d 177 (near optical peak) place the nova at  $V = 10.8$  mag. Based on this value and assuming a reddening  $A_V = 3.5$  mag and a distance  $\geq 4.5$  kpc towards Gaia22alz (see Section 3.3), we derive a peak absolute magnitude  $M_V < -5.9$ , which is within the typical range of peak absolute magnitudes of classical novae:  $M_V = -5$  to  $-11$  mag (Shafter 2017; Schaefer 2022). Moreover, the absolute magnitude of the nova and its light-curve morphology is comparable to other slowly evolving novae (e.g. V732 Cas, HR Del, and V5558: Sgr Kato, Hachisu & Henze 2013).

### 3.2 Spectral evolution

In Fig. 2, we present the overall spectral evolution of Gaia22alz throughout the eruption. Based on the spectral features, we divide the spectral evolution into four chronological stages. These stages are highlighted with different colours in Figs 1 and 2: (1) early-rise (highlighted in blue); (2) mid-rise (highlighted in red); (3) late-rise (highlighted in green); (4) post-optical-peak (highlighted in black). A complete spectral evolution is presented in Figs A2–A4.

The first two spectra obtained on d 45 and 49 during the early-rise are not typical of novae. The spectra are dominated by relatively narrow (FWHM  $\approx 400$  km s $^{-1}$ ) emission lines of H Balmer and HeI, along with HeII 4686 and 5412 Å, CIV, and the Bowen blend. The Balmer lines are characterized by broad wings extending to around 2500 km s $^{-1}$  (see Fig. 3). We measure a flux ratio  $F_{\text{HeII}}/F_{\text{H}\beta} \approx 2$  from both spectra, highlighting the strength of the HeII emission line at 4686 Å, which indicates the presence of a hot emitting source ( $T \approx$  a few times  $10^5$  K; e.g. Morisset & Pequignot 1996; Bonning et al. 2013). The narrow lines and the absence of P Cygni profiles are not consistent with optically thick ejecta typically observed in novae before peak (Aydi et al. 2020b). The spectra at this stage are reminiscent of the early spectra of the 2011 eruption of nova T Pyx, which was obtained by Arai et al. (2015) only 4.4 h after the eruption discovery. The early spectra of T Pyx were also dominated by Balmer and high-excitation lines of HeII, NIII, and CIV with no P Cygni profiles. However, the lines were characterized by high velocities, typical of classical novae (FWHM  $\approx 1100$ – $1500$  km s $^{-1}$ ). Arai et al. (2015) suggested that at this early stages, the spectrum implies that the envelope of the WD would be optically thin and the optically thick wind/outflow is still weak. In the high resolution spectrum of d 49 shows a weak emission at around 6703 Å. This might be LiI 6708 Å indicative of TNR reactions, but a definitive line classification is not possible.


On d 68, the spectra of Gaia22alz show dramatic changes, with the HeII 4686 Å emission line almost completely disappearing (see Fig. 4 for a zoom-in on the region around HeII 4686 Å and H $\beta$ ), and the emergence of P Cygni absorption components in the Balmer and HeI lines at blueshifted velocities of around  $-1700$  km s $^{-1}$  (see the line profiles in Fig. 3). Between d 68 and 96 (the mid-rise phase), the Balmer and HeI P Cygni absorptions become more prominent, while showing a decrease in the radial velocities of the P Cygni absorptions to  $-1200$  km s $^{-1}$ . At this stage, the spectra are more consistent with a classical nova eruption before peak brightness. Aydi et al. (2020b) discussed the spectral evolution of novae before and after they reach optical peak and suggested that the deceleration observed in the P Cygni profiles during the rise to peak is an optical depth effect.



**Figure 1.** The ASAS-SN optical g-band light curve of nova Gaia22alz during the first 300 d of the eruption, focusing on the rise to peak brightness. The dashed lines represent times of spectroscopic observations; blue, red, green, and black lines represent different stages in the spectral evolution, namely the early-rise, mid-rise, late-rise, and post-optical-peak, respectively (See text for more details).

During the late-rise phase, between d 135 and 178, the spectra show more dramatic changes with the HeI emission lines fading, and the FeII (42), (48), and (49) multiplets taking over. At this stage, the spectra are characteristic of classical novae before peak, originating in optically thick ejecta during the so-called iron curtain phase (Shore 2014). This evolution from a spectrum dominated by Helium to FeII emission lines during the rise to peak has been observed in a few other novae (e.g. V5558 Sgr and T Pyx; Tanaka et al. 2011a; Izzo et al. 2012; Ederoclitte 2014) and is suggested to be due to a change in the optical depth of the ejecta (e.g. Shore 2014; Hachisu & Kato 2022). The origin of the so-called He/N and FeII evolution/spectroscopic classes (Williams 1992, 2012; Shore 2014) is not the focus of this paper, and it will be the topic of a future work (Aydi et al. 2023 in prep.). Instead, we will mainly focus on the spectral evolution during the early- and mid-rise phases. The overall spectral evolution of Gaia22alz during these early phases is comparable to that of the 2011 eruption of T Pyx (e.g. fig. 3 of Arai et al. 2015) but on a much slower time-scale.

After the nova peaked (d  $\sim 180$  onwards), the spectral lines show multiple absorption features at different velocities (e.g. d 203, 212, and 236 in Fig. 3). During this phase, the light-curve evolution also hints to the presence of multiple maxima (Fig. 1). The emergence of multiple spectral absorption components in the spectra of ‘flaring novae’, correlating with the multiple peaks in the light curve, has been investigated previously (e.g. Chochol et al. 2003; Pejcha 2009; Tanaka et al. 2011a, b; Walter 2016; Harvey et al. 2018; Shore et al. 2018; Aydi et al. 2019, 2020a, b) and has been suggested to be due to either multiple phases of mass-loss (multiple ejections) or due to changes in the radius of the optical photosphere, caused by variability in the emitting central source. After d 300, the light curve shows a dramatic decline, possibly due to dust condensation. During the decline, the spectral line profiles show a broad emission (FWZI  $\gtrsim 3000$  km s $^{-1}$ ), with slower absorption/emission superimposed on top of the broad base emission (see Fig. 3 last panel; d 331). This is consistent with the spectral evolution of classical novae past peak brightness as described by Aydi et al. (2020b).




**Figure 2.** The spectral evolution of nova Gaia22alz representing the different spectral stages: early-rise (blue), mid-rise (red), late-rise (green), and post-optical-peak (black). Numbers between brackets are days after  $t_0$ . The resolving power  $R$  of each spectrum is added on the left-hand side of each panel.

### 3.3 Reddening and distance

Based on Gaia DR3, which observed Gaia22alz in quiescence (before its 2022 outburst), the system has a bias-corrected (Gaia Collaboration 2023) parallax of  $\varpi = 0.084 \pm 0.122$  mas. Based on this reported parallax and its uncertainty, it is not feasible to derive an accurate distance estimate to Gaia22alz, except to say that formally the distance is  $\gtrsim 3.0$  kpc at the 95 per cent level, so very nearby distances are disfavoured. This is a weak constraint but

consistent with the distance derived from the larger reddening values (see below).

The Galactic reddening maps of Schlafly & Finkbeiner (2011) estimate a total reddening value of  $A_V = 3.47$  mag in the direction of Gaia22alz. Chen et al. (2019) use measurements from the Gaia DR2, 2MASS, and WISE surveys to estimate reddening as a function of distance in the Galactic plane. Along the line of sight to Gaia22alz, they find that at around 2 kpc; there is a significant jump in the reddening towards Gaia22alz (see Fig. 5, which plots the reddening



**Figure 3.** The line profile evolution of H $\beta$  during the first 300 d of the eruption. The red dashed line represent rest velocity ( $v_{\text{rad}} = 0 \text{ km s}^{-1}$ ). Note that the spectra on d 45 and 80 have lower resolution ( $R \approx 1000$ ), compared to the other spectral epochs ( $R \approx 4400$  or  $14\,000$ ).


along the line of sight to Gaia22alz as a function of distance). However, the Chen et al. maps only extend to  $\approx 4.5 \text{ kpc}$  in this direction. Marshall et al. (2006) use the 2MASS survey and a slightly lower resolution grid to estimate a 3D reddening map out to larger distances. This map extends to around  $8 \text{ kpc}$  in the direction of Gaia22alz.

For a reddening estimate along the line of sight to Gaia22alz, which can in turn be used to constrain the distance, we use absorption lines from diffuse interstellar bands (DIBs), which are tracers of interstellar dust. We measure the equivalent width (EW) of the lines at  $5780.5$ ,  $5797.1$ ,  $6196.0$ ,  $6204.5$ , and  $6613.6 \text{ \AA}$  and use the relations of Friedman et al. (2011) to derive an average  $E(B - V) = 1.13 \text{ mag}$ . This implies  $A_V = 3.5 \text{ mag}$ , assuming a reddening law  $R_V = 3.1$ . This value is consistent with the integrated reddening along this line of sight, as estimated by the Schlafly & Finkbeiner (2011) Galactic reddening map.


We also use the EW of the NaI D and K1 interstellar lines to derive an additional reddening estimate. The EW of the NaI D1 and D2 lines are  $0.58$  and  $0.52 \text{ \AA}$ , while that of the K1 line at  $7699 \text{ \AA}$  is  $0.105 \text{ \AA}$ . These interstellar lines are shallow and narrow, arguing that they are not saturated. Using these values along with the empirical relations from Munari & Zwitter (1997), we derive  $E(B - V) \approx 0.5 \text{ mag}$ , translating to  $A_V = 1.55 \text{ mag}$  assuming  $R_V = 3.1$ . This value is half that of the Schlafly & Finkbeiner (2011) Galactic reddening maps and the value we derive from the DIBs.

We use our reddening estimates and the 3D Galactic reddening maps of Chen et al. (2019) to estimate the distance to Gaia22alz. In Fig. 5, we compare reddening as a function of distance (along several lines of sight that pass close to Gaia22alz) with our reddening values derived from the DIBs and the NaI D/K1 lines. Since the maps of Chen et al. (2019) use measurements from the Gaia DR2, 2MASS and WISE surveys, we use the reddening laws of Wang & Chen






**Figure 4.** The spectral evolution during the first two spectral epochs, but focusing on the evolution of HeII 4686 Å and H $\beta$  emission lines. The spectra shows the fading/disappearance of the HeII line and Bowen line between early- and mid-rise.




**Figure 5.** The extinction along the line of sight to Gaia22alz plotted against the distance. We choose the closest directions to the position of Gaia22alz from the Chen et al. (2019) and Marshall et al. (2006) Galactic reddening maps. The blue and red curves are the extinction values from the Chen et al. (2019) maps in the direction  $([l, b] = [277.55^\circ, -3.75^\circ])$  and  $([l, b] = [277.55^\circ, -3.65^\circ])$ , respectively. The magenta curve represents the extinction values from the Marshall et al. (2006) maps in the direction  $([l, b] = [277.50^\circ, -3.75^\circ])$ . Distances allowed by the *Gaia* parallax at 95 per cent confidence are shaded in green ( $> 3$  kpc).

(2019) to convert  $E(B - V) = 1.13$  mag into  $E(G_B - G_R) = 1.44$ ,  $E(G - K_s) = 2.49$ , and  $E(H - K_s) = 0.18$  mag. All these values yield a distance larger than 4.5 kpc. However, if we use  $E(B - V) = 0.5$  mag (derived from the NaI D and KI interstellar lines), we get a distance of  $2.5 \pm 0.5$  kpc using the same maps. This lower reddening value, and correspondingly lower distance, is not consistent with the *Gaia* parallax, which suggests a distance likelihood in excess of 3 kpc.



**Figure 6.** A  $1 \times 1$  arcmin DECaPS *grz* image of the field of Gaia22alz taken on 2017 October 4. The candidate progenitor system of Gaia22alz is circled in green.




**Figure 7.** The observed spectral energy distribution of the progenitor system of Gaia22alz constructed using DECaPS and VPHAS+ data. We fit two models to the SED: a blackbody model with Rayleigh-Jeans tail (black line) and an optically thick steady accretion disc model (magenta line) to the VPHAS+ (solid lines) and De-CaPS (dashed lines) data.

Therefore, throughout the paper, we assume a distance  $d > 4.5$  kpc and a reddening value  $A_V = 3.5$  mag towards Gaia22alz, as implied by the DIBs and consistent with the *Gaia* parallax and the Schlafly & Finkbeiner (2011) maps. However, we also discuss the implications of shorter distances and lower reddening towards Gaia22alz in the following sections.

### 3.4 The progenitor system

Archival data from the Optical Gravitational Lensing Experiment (Udalski, Szymański & Szymański 2015), taken between 2017 and 2020, measured average magnitudes  $I = 17.57 \pm 0.12$  and  $V = 18.66 \pm 0.09$  of the progenitor system of Gaia22alz during quiescence. In archival data from DECaPS, the source has median pre-eruption magnitudes of  $g \simeq 19.0$ ,  $r \simeq 18.2$ ,  $i \simeq 18.1$ ,



**Figure 8.** Colour-magnitude diagram adopted from Abril et al. (2020) showing the colours and absolute magnitudes, measured by Gaia, for several CV systems from different sub-types (classical nova and dwarf nova systems in quiescence, nova-likes, polars, intermediate polars, all taken from the Catalogue and Atlas of Cataclysmic Variables Downes et al. 2001). The dashed ellipses represent 1 and  $3\sigma$  of each sub-type bivariate Gaussian distribution (see Abril et al. 2020 for more details). We overplot the progenitor system of Gaia22alz at different distances; the red star corresponds to  $d = 4.5$  kpc and  $A_V = 3.5$  mag (derived based on the DIBs), while the blue star is for  $d = 2.5$  kpc and  $A_V = 1.55$  mag (derived based on the Na I D and K I lines). We also add a case of an intermediate distance/reddening (green square;  $d = 3.5$  kpc and  $A_V = 2.5$  mag).

$z \simeq 18.0$ , and  $y \simeq 17.9$ . Moreover, the AB magnitudes of the progenitor system,  $u = 19.56 \pm 0.02$ ,  $g = 19.04 \pm 0.01$ ,  $r = 18.40 \pm 0.02$ ,  $m(H\alpha) = 18.23 \pm 0.02$ , and  $i = 18.01 \pm 0.02$ , measured by the VST Photometric  $H\alpha$  Survey of the Southern Galactic Plane and Bulge (VPHAS+; Drew et al. 2014) on 2012 December 20 (HJD = 2456282.76), indicate a hot source and no substantial  $H\alpha$  emission.

The above measurements imply an absolute quiescent  $V$ -magnitude  $M_V \leq 1.8$ , assuming  $A_V = 3.5$  mag and a distance  $\geq 4.5$  kpc. This means that the system is luminous compared to other cataclysmic variables (CVs) in quiescence (e.g. Warner 1987). At such a brightness, the secondary star could be more evolved than a dwarf star, i.e. a sub-giant or even a giant. However, the colours of the system,  $(g - i)_0 \simeq -0.9$  and  $(V - I)_0 \simeq -1.73$  (assuming  $A_V = 3.5$  mag), are blue for a typical CV (whether it has a dwarf or giant companion; Bruch & Engel 1994; Szkody 1994; Zwitter & Munari 1995; Szkody et al. 2011; Kato, Maehara & Uemura 2012), ruling out the possibility of a giant evolved secondary. In Fig. 6, we show a combined  $grz$  colour image of the region of Gaia22alz from the DECaPS survey (Schlafly et al. 2018), showing the progenitor system.

We created a spectral energy distribution (SED) using the multiband photometry taken during quiescence by VPHAS+ and DECaPS, and we successfully fit the resulting SED with a reddened Rayleigh–Jeans tail of a hot blackbody (Fig. 7). The fit to both data sets provide almost identical outcomes. The best fit has been obtained with a radius  $R = (1.595 \pm 0.059) \times 10^{10} (T/10^5 K)^{-1} (d/5 \text{ kpc}) \text{ cm}$  and  $E(B - V) = 1.11 \pm 0.02$  mag. Note that this reddening value is practically identical to the one we derived from the DIBs and the total Galactic reddening resulting from Schlafly & Finkbeiner (2011) maps in the direction of Gaia22alz. There is no evidence for a cool companion, as up to  $\sim 1$  micron

( $Y$  band) the SED is well-reproduced by the hot blackbody, and in particular, any companion could not contribute more than a few per cent to the SED, which rules out a yellow/red giant donor.

Integrating under the blackbody, the source is luminous with a luminosity  $L = (4740 \pm 350) (T/10^5 K)^2 (d/5 \text{ kpc})^2 L_\odot$ , indicating the presence of a bright emitting source, which could be due to ongoing nuclear burning on the white dwarf surface (Schaefer & Collazzi 2010; Wolf et al. 2013; Zemko, Mukai & Orio 2015; Zemko et al. 2016; Aydi et al. 2018). However, since a thermonuclear runaway took place on the white dwarf in 2022, leading to the nova eruption of Gaia22alz, it is unlikely that the system has been undergoing nuclear shell burning. CVs are known to show strong emission lines during quiescence (Jiang et al. 2013; Breedts et al. 2014; Hou et al. 2020, 2023). These emission lines could potentially affect the colours of the system during quiescence, leading to bluer colours. While based on the VPHAS+ photometry, we do not observe significant  $H\alpha$  emission above the SED BB, emission lines might still have an effect on the overall emission and colours. Therefore, using a blackbody fit to describe the emission in CVs at quiescence is not necessarily an ideal assumption. Noting this caveat, we also attempt to fit the SED with an optically thick steady accretion disc model ( $F_\nu \propto \nu^{1/3}$ ; Fig. 7; magenta line). The best fit to the observed SED results in  $E(B - V) = 0.639 \pm 0.012$  mag. This value is more consistent with the lower reddening values derived from the Na I D and K I interstellar lines.

In order to compare the brightness and colours of Gaia22alz with other CVs and to better constrain the nature of the secondary star, we place the progenitor system of Gaia22alz on a colour-magnitude-diagram (CMD) adopted from Abril et al. (2020) in comparison with different CV sub-types (e.g. nova systems, dwarf novae, nova-likes, and magnetic CVs; Fig. 8). The colours and magnitudes of these CV



systems are all Gaia measurements. At a distance of 4.5 kpc and  $A_V = 3.5$  mag (red star in Fig. 8), the system is much bluer than other CV systems and is relatively brighter. However, for a distance of 2.5 kpc and  $A_V = 1.55$  mag (blue star in Fig. 8), the system has comparable colours and brightness to other CV systems, particularly novae and nova-like systems. Both cases, small or large distance/reddening, still rule out the presence of a giant, evolved secondary. It is possible that the distance and reddening derived from the DIBs are overestimated. Therefore, intermediate distances and reddening values (e.g.  $d = 3.5$  kpc and  $A_V = 2.5$ ; green square in Fig. 8), could bridge the gap between the colours/brightness of Gaia22alz and other CVs.

In summary, the Gaia parallax of the progenitor system of Gaia22alz favours larger distances ( $d > 3$  kpc) and therefore larger reddening towards the system. However, at such large distances and high reddening values, the progenitor system is relatively luminous and characterized by significantly bluer colours compared to other CVs, as shown in Fig. 8. This might lead us to suggest that the lower reddening/shorter distances are more reasonable to explain the colours and brightness of the system during quiescence, despite them being contradictory with the Gaia parallax. However, at such a distance and reddening, the absolute peak magnitude of Gaia22alz during eruption would be just  $M_V = -2.7$  (assuming  $d = 2.5$  kpc and  $A_V = 1.55$ ), which is much smaller than typical peak brightness of novae ( $M_V \approx -5$  to  $-10$ ; Shafter 2017; Shara et al. 2017). Could this mean that Gaia22alz is not a classical nova? We further elaborate on these speculations in the following sections. It is also possible that the actual distance and reddening towards Gaia22alz is intermediate between the values we derive from the DIBs and the ones we derive from the NaI D and KI lines.

## 4 DISCUSSION

### 4.1 Detecting the visible signature of the UV/X-ray flash

Novae observed during the rise to peak brightness typically show spectra dominated by P Cygni lines of H I and either Fe II or He I/N, originating in the dense nova ejecta (e.g. McLaughlin 1944; McLaughlin 1947; Gaposchkin 1957). The line profiles are characterized by velocities ranging from a few hundreds  $\text{km s}^{-1}$  (Aydi et al. 2020b) up to a few thousands  $\text{km s}^{-1}$  in some extreme cases (e.g. the very fast nova V1674 Her; Aydi et al. 2021). Aydi et al. (2020b) observed a large sample of novae before they reached optical peak and found that all of them show the same evolution. Particularly, they found that the optical spectra before peak are dominated by P Cygni profiles originating in the dense nova outflow. Gaia22alz is an exception. The first two spectra obtained on d 45 and 49 were not typical of classical novae observed before peak. More precisely, the absence of P Cygni absorption, the relatively narrow emission lines ( $\text{FWHM} \approx 400 \text{ km s}^{-1}$ ), and the strong He II and C IV emission lines are unusual for a classical nova pre-maximum spectrum. These features, observed at a stage when the eruption was less than four magnitudes above quiescence, led Brink et al. (2022) to suggest that Gaia22alz could be an unusual disc instability dwarf nova event or something more exotic. Here, we discuss the origin of these features and speculate about the reasons for the gradual rise in the optical light curve of nova Gaia22alz.

After the onset of the thermonuclear runaway, the CNO chain reactions become the main source of energy generation in a nova, producing radioactive elements that decay ( $\beta^+$ -decay), releasing a large amount of energy. It has been theorized (Starrfield et al. 1990; Krautter 2002; Hillman et al. 2014) and now observationally confirmed (König et al. 2022) that the dramatic increase in fusion rate

is followed by a short X-ray (and even UV) flash before the dramatic increase in the visible brightness. The X-ray/UV flash terminates when the nova envelope has had time to expand, thereby cooling down rapidly and becoming optically thick to the emission from the hot white dwarf. The peak of the nova's SED shifts into the visible band, turning the nova into an optical transient. The optical spectrum of the nova during this phase (rise to visible peak) is typically dominated by P Cygni profiles (Aydi et al. 2020b). However, it is not clear how the visible emission from the nova manifests during the X-ray/UV flash phase.

The first two spectra we obtained for Gaia22alz possibly capture the late stages of an extended, slow X-ray/UV flash. Unfortunately, we do not have any X-ray or UV observations taken during the rise to peak. However, the strong high-excitation He II 4686 Å and C IV emission lines in the first two spectra imply emission from a hot ionized gas (a few times  $10^5$  K; e.g. Morisset & Pequignot 1996; Bonning et al. 2013). Below, we suggest two possible origins for the early, gradual optical rise and the relatively narrow, high ionization spectral features (with no P Cygni absorptions) we observe in the early optical spectra (first 50 d):

(1) The optical emission is originating in the outer layers of the (yet optically thin) nova envelope in a scenario where the bulk of the envelope has not yet reached the escape velocity and has not expanded enough in size to cause a bright optical transient (i.e. a transient characterized by an amplitude of more than 8 mag). The outer layers of this slightly puffed-up nova envelope reprocess the high-energy X-ray/UV emission from the white dwarf surface, resulting in the spectral features we observe in the first 50 d and the early brightness increase. This means that the four magnitudes of gradually increasing optical brightness during the first 50 d might be taking place during the X-ray/UV flash. While the X-ray/UV flash is often predicted to be short (of the order of a few hours; Starrfield et al. 1990; Krautter 2002; König et al. 2022), Hillman et al. (2014) modelled the precursor UV flash for a range of system parameters, and found that for some models (e.g. a low mass white dwarf with  $M_{\text{WD}} = 0.65 M_{\odot}$  and a core temperature  $T_c = 10^7$  K, accreting at a rate  $\dot{M} = 10^{-10} M_{\odot} \text{ yr}^{-1}$ ), a UV flash preceding the optical peak emission can last for up to 60 d. While these models might not necessarily represent the accurate duration of the UV flash, they show that these UV flashes/precursors could last for days (see also Cao et al. 2012; Kato, Saio & Hachisu 2022). This time-scale is consistent with the duration of the early rise in Gaia22alz ( $\approx 50$  d). Moreover, based on the slow evolution of the light curve, we indeed expect that the WD mass in Gaia22alz is lower than 1 solar mass. Given that the white dwarf mass of Gaia22alz might be on the lower end of white dwarf masses in novae ( $M_{\text{WD}} \sim 0.6$  to  $-0.8 M_{\odot}$ ), it is likely that the high-energy emission from the WD is mostly in the far-UV and the X-ray flux is relatively weaker (see e.g. Kato et al. 2022).

The wings of the Balmer lines, which extend to velocities  $\gtrsim 2000 \text{ km s}^{-1}$  at this early epoch (see Fig. 3), possibly originate in a low-density, radiation-driven wind from the white dwarf surface, before the bulk of the envelope is ejected. Hachisu & Kato (2014) modelled the early emission of novae and suggested the presence of an optically thin expanding ejecta/wind during the early days of the eruption, which manifest as emission lines in the spectra with no P Cygni profiles.

After the X-ray/UV-flash and the early visible rise, we expect that the bulk of the nova envelope is ejected, possibly aided by the binary motion (Chomiuk et al. 2014). This would lead to the final climb towards peak brightness (days  $\sim 50$ –180). During this stage, the envelope cools to the critical temperature at which a recombination wave forms, so the envelope becomes opaque to the UV emission

**Table 1.** A comparison between the parameters of Gaia22alz (for different distance and extinction values) with the properties of classical novae and dwarf novae, along with the 2005 and 2019 outbursts of V1047 Cen (Aydi et al. 2022a), and the 2011 eruption of T Pyx (Arai et al. 2015; Godon et al. 2018).

| System                                | $\Delta m^a$<br>(mag) | $M_{V, \text{peak}}$ | FeII P Cygni $^b$<br>profiles | HeII/CVI<br>during the rise | Dust formation |
|---------------------------------------|-----------------------|----------------------|-------------------------------|-----------------------------|----------------|
| Gaia22alz ( $d=4.5$ kpc; $A_V=3.5$ )  | 7.8                   | −5.9                 | Yes                           | Yes                         | Yes            |
| Gaia22alz ( $d=2.5$ kpc; $A_V=1.55$ ) | 7.8                   | −2.7                 | Yes                           | Yes                         | Yes            |
| Classical novae                       | 8–15                  | −5 to −11            | Yes                           | No                          | Yes            |
| Dwarf novae                           | 2–5                   | > 0.0                | No                            | HeII Yes                    | No             |
| V1047 Cen (2019)                      | 6                     | −2.0                 | No                            | Yes                         | No             |
| V1047 Cen (2005; classical nova)      | 13                    | −8.0                 | Yes                           | No                          | No data        |
| T Pyxidis (2011; classical nova)      | 9                     | −8.1                 | Yes                           | Yes                         | No             |

<sup>a</sup>  $\Delta m$  is outburst amplitude.<sup>b</sup> Appearance of prominent FeII P Cygni profiles near optical peak.

(Shore 2014). This manifests as a transition in the optical spectra from a spectrum dominated by narrow emission lines to one dominated by higher velocity P Cygni profiles. This also manifests as the sudden disappearance of the high excitation lines of HeII and CIV (see first two panels in Fig. 2). This is similar to the overall spectral evolution observed during the 2011 eruption of T Pyx (Arai et al. 2015; see also Table 1). T Pyx also shares other similarities with Gaia22alz, particularly the optical light-curve evolution that showed a relatively slow rise (40 d) to visible brightness with a plateau phase at peak that lasted for more than a month (Schaefer et al. 2013).

Optical multiband photometry taken during the first 100 d would have been very informative to check if there is any change in the temperature of the emitting source and to derive its size. However, reported AAVSO observations only start after d 97, missing the crucial early- and mid-rise phases. Nevertheless, if the temperature of the emitting source is of the order of a few times  $10^5$  K (as implied by the high-excitation emission lines) and the brightness is  $g \approx 15$  (implying  $M_g \approx 2.5$  mag for a distance of 4.5 kpc), the optical emission should be originating close to the surface of the white dwarf (at  $r \sim 10^9$  cm; an emitting region comparable to the radius of the white dwarf).

(2) Another, less likely, possibility is that the optical emission is originating in the accretion disc, which is reprocessing high-energy (X-ray and/or UV) emission from the white dwarf surface. Similar to the scenario described above, the first 50 d represent the optical signature of an extended X-ray/UV flash, but in this case, the visible emission is originating in the disc rather than the outer layers of a slightly puffed-up nova envelope. Emission lines of Balmer, HeI, and HeII are often observed in CV systems and dwarf nova outbursts (e.g. Morales-Rueda & Marsh 2002), and they are characterized by velocities of a few hundreds  $\text{km s}^{-1}$  – similar to the ones we observe in the first two spectra of Gaia22alz. Could reprocessed emission by the accretion disc lead to a visible brightness increase of  $\sim 4$  mag and absolute optical magnitudes spanning +2 to −2.5 (assuming  $d = 4.5$  kpc and  $A_V = 3.5$  mag), during the first 50 d of the outburst? In the case of supersoft X-ray sources (e.g. van den Heuvel et al. 1992), which are similarly powered by nuclear burning on the white dwarf and which part of their visible emission is associated with emission from the accretion disc (in addition to the companion star), we observe absolute magnitudes between +1.5 and −2.5 (e.g. Smale et al. 1988), comparable to the brightness of Gaia22alz during the first 50 d of the eruption. However, this is merely an indirect argument that visible emission from the accretion disc could potentially explain the early rise of Gaia22alz. We note that the accretion disc could well be engulfed by the nova envelope from the early days of the eruption, favouring the first suggestion we present in this section.

Whether the high-energy emission is reprocessed by the outer layers of the still-optically thin and hot nova envelope or the accretion disc, Gaia22alz could be one of the rare times we observe the optical signature of the hot X-ray/UV flash, which is often missed in other novae. We can speculate that the extremely slow and gradual rise of nova Gaia22alz and the diverse all-sky surveys that are observing the sky more efficiently at deep magnitudes (visible magnitudes down to 17–18 mag) is what allowed us to catch this early phase of a nova eruption. Given the lack of multiwavelength observations during this early and critical phase, which prevent us from drawing definitive conclusions, we consider alternative scenarios below and explore their pros and cons for explaining the unusual observational features of Gaia22alz.

## 4.2 Alternative explanations

Below we suggest some alternative explanations to the early spectral features of Gaia22alz and its unusual, gradual rise. The characteristics of some of these different scenarios are compared in Table 1, considering two different distance and reddening assumptions for Gaia22alz.

(i) **Shock interaction from an early mass-loss phase?** The main alternative scenario we suggest here is that, during the first weeks of the eruption (before the bulk of the nova envelope is ejected), the system loses mass from the L2 outer Lagrangian point. The gravitational potential of the rotating binary is not axially symmetric, so the matter leaving from L2 exhibits small fluctuations in its radial velocity (see Pejcha, Metzger & Tomida 2016 for a detailed description of the model). This causes the matter being ejected from L2 as spiral arms to collide with itself at around 10 times the semimajor axis of the binary, creating stationary shocks that radiate some of the kinetic energy of the L2 outflow. In Section B of the Appendix, we present calculations for the temperature of the shocked gas and the luminosity of the line emission from these stationary shocks. Our calculations show that the shocks would heat the gas to a temperature of  $\sim$ few times  $10^5$  K with emission line luminosity of  $10^{35} \text{ erg s}^{-1}$ , consistent with the early spectra of Gaia22alz.

(ii) **A dwarf nova outburst preceding/triggering a classical nova?** The features detected in the first two spectral epochs of Gaia22alz resemble features observed in disc instability dwarf nova outbursts (see Warner 1995 for a review). During outburst, the spectrum of a dwarf nova is typically dominated by absorption features with fill-in by emission lines. Moreover, the spectra could exhibit strong emission lines, particularly at later stages of the outburst (e.g. Bruch 1989; Bruch & Schimpke 1992; Morales-Rueda

& Marsh 2002; Han et al. 2020). Strong emission lines of HeII at 4686 Å are also commonly observed in the spectra of dwarf novae (Han et al. 2020). The gradual increase of just a few magnitudes in the optical light curve during the first 50 d of Gaia22alz eruption is consistent with the amplitude of a dwarf nova outbursts ( $\approx 2\text{--}5$  mag). Therefore, could it be that the classical nova eruption of Gaia22alz was coincidentally preceded or maybe even triggered by a dwarf nova outburst? Mróz et al. (2016) observed multiple dwarf nova outbursts preceding the classical nova eruption of nova V1213 Cen in 2009. Similarly, V392 Per and V1017 Sgr showed DN outbursts 2 and 8 yr, respectively, prior to their classical nova eruptions (Salazar et al. 2017; Murphy-Glasyher et al. 2022). Meaning that DN outburst could precede a classical nova eruption by a few years, and potentially in the case of Gaia22alz, happening around the same time as the classical nova eruption started. Could this be the case? While this is possibility, it is unlikely that a disc instability event would be the trigger for the classical nova eruption. In particular, we note that the onset of the thermonuclear runaway is expected to take place years before the classical nova becomes a visible transient (Starrfield, Sparks & Truran 1974; Prialnik 1986; Shen & Quataert 2022). Moreover, the material dumped on to the white dwarf from a dwarf nova outburst is suggested to be of the order of  $10^{24}$  g ( $\sim 10^{-9} M_{\odot}$ ) in systems with dwarf companions (e.g. Cannizzo 1993), meaning that the total accreted mass due to the dwarf nova outburst is likely negligible compared to the total mass of the classical nova accreted/ejected envelope ( $10^{-7}\text{--}10^{-3} M_{\odot}$ ; Yaron et al. 2005). In conclusion, it is unlikely that the eruption of Gaia22alz is triggered by a dwarf nova outburst. However there remains a slight chance of coincidence in Gaia22alz: in this scenario, the early rise of the light curve and the early spectral features (narrow emission lines of Balmer, HeI, and HeII) are due to a disc instability event that coincidentally directly precedes the classical nova eruption. Nevertheless, this possibility also does not agree well with the pre-eruption magnitudes and SED fitting (Fig. 7). The SED fitting to the VPHAS+ and DECaPS data implies a steady optically thick accretion disc for at least the epochs covered by these observations. However, a steady optically thick accretion disc is inconsistent with a dwarf nova because their quiescent accretion disc are usually optically thin (Han et al. 2020). Similarly, the colours and brightness of the system during quiescence are different from most DN systems (Fig. 8).

(iii) **An outburst similar to the 2019 outburst of V1047 Cen?** Gaia22alz share similarities with the peculiar 2019 outburst of V1047 Cen, which was first identified as an outbursting classical nova in 2005 (Aydi et al. 2022a). The 2019 outburst was characterized by a long rise (lasting more than 2 months), a slowly evolving light curve (with an outburst duration  $>400$  d), and showed significant differences with the 2005 classical nova eruption experienced by the same system (fig. 1 of Aydi et al. 2022a). In Fig. A1, we show a comparison of the light curves of Gaia22alz and V1047 Cen, where both show a slow rise and continuous variability near peak brightness for several months. The spectra of the 2019 outburst of V1047 Cen also showed narrow emission lines (FWHM of a few hundred  $\text{km s}^{-1}$ ) of Balmer and HeI, HeII, and CIV during the rise to peak. As in Gaia22alz, Aydi et al. (2022a) noted that these high-excitation HeII and CIV emission lines disappeared as the outburst evolved. The colours of the system prior to the 2019 outburst were also significantly blue ( $[V - I_0] = -1.19$ ), which led Aydi et al. (2022a) to suggest that ongoing nuclear burning was taking place on the surface of the white dwarf. Aydi et al. (2022a) suggested that the outburst of V1047 Cen may have started with a brightening of the disc due to enhanced mass transfer or disc instability, possibly leading to enhanced nuclear shell burning on the white dwarf, which

was already experiencing some level of quasi-steady shell burning in the wake of the 2005 nova. The nuclear shell burning eventually led to the generation of a wind/outflow, resulting in significant broadening in the emission lines (see Aydi et al. 2022a for more details).

Could Gaia22alz's outburst be similar to that of the 2019 outburst of V1047 Cen rather than being a classical nova? While there are some similarities between V1047 Cen and Gaia22alz (e.g. the slow rise to peak brightness, the high-excitation lines during the rise to visible peak, the blue colours of the progenitors, and the slowly evolving outbursts), the two events also show discrepancies. As the event evolved, the 2019 outburst of V1047 Cen did not show typical observational features of classical novae, while Gaia22alz did develop such features (e.g. peak absolute magnitude and emergence of FeII P Cygni profiles, potential dust formation). As detailed in Table 1, at a distance of 4.5 kpc and for the relatively large extinction value, Gaia22alz would have a peak absolute magnitude of  $M_V = -5.9$ , consistent with a classical nova eruption. However, V1047 Cen only reached  $M_V \approx -2$  in 2019 (much smaller than typical brightness of classical novae). The gap is bridged if we consider shorter distances and lower extinction towards Gaia22alz. Nevertheless, the optical spectral evolution of the two systems is still different, with Gaia22alz exhibiting spectra dominated by strong Balmer and FeII P Cygni profiles as the outburst evolved, which was not the case of V1047 Cen, i.e. no prominent FeII P Cygni profiles. Note that the 2005 classical nova eruption of V1047 Cen, showed typical spectral features of classical novae with spectra dominated by P Cygni profiles of Balmer and FeII near optical peak; Walter et al. 2012; Aydi et al. 2022a.

(iv) **Pre-existing circumbinary material?** Could the early spectral features observed in Gaia22alz originate in circumbinary material reprocessing the high-energy emission from the white dwarf surface? In this scenario, we still require high-energy luminous emission from the X-ray/UV flash, but it is then absorbed by the neutral gas in the circumbinary medium and re-emitted in the visible band. Many nova systems, particularly ones with evolved giant secondaries, are known to have dense circumbinary material enriched by the mass-losing, giant companion (e.g. V3890 Sgr- Harrison, Johnson & Spyromilio 1993; RS Oph- O'Brien et al. 2006; V407 Cyg- Munari et al. 2011b; Booth, Mohamed & Podsiadlowski 2016; Orío et al. 2020; Mikołajewska et al. 2021). In contrast, novae with main-sequence, dwarf secondaries are characterized by low-density circumbinary media (e.g. Bode, Seaquist & Evans 1987). The dense circumbinary material in the case of novae with giant secondaries manifests as low-velocity ( $\approx 10\text{--}50 \text{ km s}^{-1}$ ) P Cygni profiles in their optical spectra (e.g. Dobrzycka & Kenyon 1994; Brandi et al. 2009; Mondal et al. 2018). These velocities reflect the relatively slow mass-loss from the red giant companion (Bowen 1988).

The optical spectra of Gaia22alz do not show such features. The relatively narrow emission lines observed in its early spectra are characterized by FWHM  $\approx 400 \text{ km s}^{-1}$ , which is significantly faster than the velocities typically observed for winds/mass-loss from evolved stars in symbiotic systems (Brandi et al. 2009; Munari et al. 2011a). Moreover, the blue colour of the system during quiescence argues against an evolved secondary star (see Section 3.4 and Fig. 8). Therefore, it is unlikely that the early spectra of Gaia22alz originate in circumbinary material reprocessing the emission from the nuclear burning on the white dwarf.

(v) **An unusual symbiotic nova?** Symbiotic novae are thermonuclear runaway events that take place on accreting white dwarfs in wide binary systems (e.g. Kenyon & Truran 1983; Mikołajewska 2010). These systems consist of a white dwarf accreting material from an evolved, red giant companion (Belczyński et al. 2000;



Mikolajewska 2007). Traditional symbiotic novae are characterized by slow outbursts that last years and even decades (in contrast with recurrent novae in symbiotic systems, which tend to evolve rapidly; see Mikolajewska 2010). The orbital periods of such systems are much longer than those of classical nova systems ( $P_{\text{orb}} > 100$  d for symbiotic nova systems, compared to  $P_{\text{orb}} < 1$  d for classical nova systems), and the white dwarf hosts are characterized by low masses ( $0.4\text{--}0.6 M_{\odot}$ ; Mikolajewska 2003).

While the outburst of Gaia22alz is slow compared to classical novae ( $t_2 = 212$  d, qualifying it as a ‘very slow’ nova; Gaposchkin 1957), the time-scales on which symbiotic novae evolve tend to be even slower (a few years up to decades; e.g. Chochol et al. 2003; Mikolajewska 2010). Could Gaia22alz be an unusually rapidly evolving symbiotic nova? There is generally a scarcity of spectroscopic follow-up of slow symbiotic novae during their rise to optical peak to compare with Gaia22alz early spectra, but Munari et al. (2009) observed the slow symbiotic nova V4368 Sgr across 16 yr of its outburst that started in 1993. The pre-maximum spectra show relatively narrow emission lines of Balmer with no P Cygni profiles, similar to the early spectra of Gaia22alz. However, they do not detect high-ionization HeII or CIV lines. The spectral features of V4368 Sgr remained similar after its optical peak, with no P Cygni profiles emerging. This led Munari et al. (2009) to suggest that up until the time of their study, mass-loss from the nova was still very low or even absent. While the spectral evolution of V4368 Sgr might not necessarily represent all slow symbiotic nova, its overall evolution is different from Gaia22alz, which clearly showed evidence for ejected material and mass-loss as the eruption evolved. Moreover, based on the archival data of the progenitor system (see Section 3.4 and Fig. 8), it is unlikely that the companion star is a red/yellow giant, arguing against a symbiotic nova possibility.

(vi) **A stellar merger?** One alternative suggestion to explain the unusual spectral and photometric evolution of Gaia22alz is that the eruption is more exotic than a nova outburst. While the spectra observed after d 50 become consistent with a classical nova eruption, the amplitude of the eruption ( $\approx 8$  mag) is on the lower end of eruption amplitudes in novae (Warner 2008; Kawash et al. 2021). The optical light curve of Gaia22alz resembles the optical light curves of stellar mergers (Luminous Red Novae; see fig. 6 in Pastorello et al. 2019), showing a relatively slow evolution and secondary peaks. Could it be that Gaia22alz is a stellar merger? The current data set does not support this hypothesis; in particular, the spectra never showed a red continuum nor particular absorption bands that are usually seen in luminous red novae (e.g. TiO and VO; Williams et al. 2015; Pastorello et al. 2019). Nevertheless, further follow-up are encouraged to confirm the nature of this event and obtain the parameters of the binary after the system has returned to quiescence.

### 4.3 The progenitor brightness and colours

Photometry of Gaia22alz’s progenitor system during quiescence indicates emission from a luminous and hot, blue source at the large distance and extinction estimates ( $d = 4.5$  kpc and  $A_V = 3.5$ ; see Section 3.4 and Fig. 8). Here, we speculate about potential causes for this unusually high luminosity and remarkably blue colours.

To first order, ongoing nuclear burning on the surface of the white dwarf seems to be the most reasonable explanation for the brightness and colours of the progenitor. Ongoing nuclear burning could be due to a relatively high mass transfer rate ( $10^{-8}\text{--}10^{-7} M_{\odot} \text{ yr}^{-1}$ ; e.g. Wolf et al. 2013). However, sustained nuclear shell burning on the white dwarf could not directly precede a 2022 thermonuclear nova eruption (Starrfield 1989; Wolf et al. 2013). We can, in turn, speculate that

the 2022 eruption of Gaia22alz is not a thermonuclear nova eruption, but rather a dramatic brightness increase due to wind/shell ejection driven by the nuclear shell burning, particularly due to the extremely slow and gradual rise – in a scenario similar to that suggested for the 2019 outburst of V1047 Cen (Aydi et al. 2022a), as discussed above. However, we have showed in Section 4.2 that there are differences in the overall evolution of Gaia22alz and V1047 Cen.

Could the accretion disc be undergoing an active state prior to the nova eruption? Several recurrent novae are shown to go through active high-accretion states prior to a nova eruption. This manifests as an increase in the system brightness by a fraction of a magnitude accompanied with brightness flickering, a change in the colours towards the blue (in the optical), and appearance of high excitation emission lines, such as HeII, in the optical spectra (e.g. Munari, Dallaporta & Cherini 2016). This active state is suggested to be due to an increase in the accretion rate on to the white dwarf, with the flickering originating in the accretion disc (Luna et al. 2018). While an increased accretion rate leading to the 2022 eruption of Gaia22alz could increase the brightness of the system, the lack of multiwavelength observations during quiescence prevents us from reaching a conclusive explanation for the brightness/colours of the system.

Similarities with T Pyx in quiescence: we have discussed above the similarities in the spectral and light-curve evolution of Gaia22alz and T Pyx. The two systems also share similarities during quiescence. The absolute magnitude of T Pyx during quiescence is  $M_V = 1.8 \pm 0.8$ , assuming a distance between 2.5 and 4.5 kpc and  $E(B - V) = 0.35$  (see e.g. Patterson et al. 1998; Godon et al. 2018). This is also significantly bright compared to other CVs with dwarf companions and is equal to the brightness of Gaia22alz during quiescence ( $M_V = 1.8$ ; for  $d = 4.5$  kpc and  $A_V = 3.5$  mag). The colours of T Pyx during quiescence are also bluer than typical CVs, with  $(B - V)_0 = -0.35$ . This means that both systems are significantly bright and blue in comparison to other nova systems in quiescence, which might imply certain physical mechanisms or system properties that are responsible for the quiescent characteristics of systems like Gaia22alz and T Pyx. However, investigating these mechanisms are beyond the scope of this paper.

A lower reddening and distance estimate? The reddening we derive from the NaI D and KI interstellar lines ( $E(B - V) = 0.5$ ) suggests a smaller distance towards Gaia22alz. At these distances and reddening, the colours of the progenitor system are consistent with other CVs in quiescence (see Fig. 8). However, as mentioned above, at such distances and reddening, the peak absolute magnitude of the nova would be  $M_V \approx -2.7$ . This is much fainter than the peak intrinsic brightness of typical classical novae (Shafter 2017). Could this be an indication that Gaia22alz is an unusually faint classical nova (particularly due to the unusual spectral evolution and very slow and gradual rise of the eruption)? While unlikely, this would make Gaia22alz one of the least luminous classical novae on record.

As discussed above, Intermediate values of reddening and distance (e.g.  $d = 3.5$  kpc and  $A_V = 2.5$  mag) could potentially bridge the gaps and result with more reasonable quiescent colours [ $(V - I) = -0.2$  &  $(G_{BP} - G_{RP}) = 0.037$ ; see Fig. 8] and acceptable peak absolute magnitude ( $M_{V, \text{peak}} \approx -4.4$ ).

## 5 CONCLUSIONS

We present a detailed analysis of the spectral evolution of the slow nova Gaia22alz during its rise to peak brightness. The spectra obtained during the first 50 d were inconsistent with typical classical nova spectra, showing relatively narrow (FWHM  $\approx 400 \text{ km s}^{-1}$ )



emission lines and strong high-excitation lines of HeII and CIV, and no P Cygni absorptions. However, as the eruption progressed, the spectra underwent dramatic changes, becoming more akin to classical novae. Based on the features observed in the early spectra (first 50 d), we suggest that the early optical emission, particularly the relatively narrow emission lines with no P Cygni absorptions, originates in the outer layers of a slightly puffed-up (but not yet ejected) optically thin nova envelope or in the accretion disc, which are reprocessing high-energy (UV/X-ray) emission from the white dwarf surface during the so-called X-ray/UV flash phase. The spectral line profiles also suggest the presence of a fast wind, possibly driven by radiation of the nuclear burning white dwarf, before the bulk of the nova envelope is ejected. If confirmed, this would be one of the rare incidents we have observed the optical signature of a nova during the X-ray/UV flash. Thanks to ongoing all-sky surveys, we anticipate more slow classical novae to be detected at early stages of eruption, which will help us to better understand and constrain the physics of the poorly observed X-ray/UV flash phases in classical novae.

The lack of multiwavelength observations prevent us from confirming these suggestions, and we speculate about alternative explanations/scenarios to explain the nature of this event and its unusual observational features. Follow-up observations in the future might allow us to constrain the parameters of the progenitor system and its physical characteristics, which could help us gain insight into the nature of the eruption of Gaia22alz.

## ACKNOWLEDGEMENTS

We thank the American Association of Variable Stars Observers (AAVSO) observers from around the world who contributed their magnitude measurements to the AAVSO International Database used in this work.

EA acknowledges support by the National Aeronautics and Space Administration (NASA) through the NASA Hubble Fellowship grant HST-HF2-51501.001-A awarded by the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., for NASA, under contract NAS5-26555. EA, LC, and KVS acknowledge National Science Foundation (NSF) awards AST-1751874 and AST-2107070, NASA award 11-Fermi 80NSSC18K1746, and a Cottrell fellowship of the Research Corporation. JS was supported by the Packard Foundation. DAHB gratefully acknowledges the receipt of research grants from the National Research Foundation of South Africa. JM was supported by the National Science Centre, Poland, grant OPUS 2017/27/B/ST9/01940. A part of this work is based on observations made with the Southern African Large Telescope (SALT), with the Large Science Programme on transients 2021-2-LSP-001 (PI: DAHB). Polish participation in SALT is funded by grant No. MEiN 2021/WK/01. This paper was partially based on observations obtained at the SOAR telescope, which is a joint project of the Ministério da Ciência, Tecnologia e Inovações (MCTI/LNA) do Brasil, the US National Science Foundation's NOIRLab, the University of North Carolina at Chapel Hill (UNC), and Michigan State University (MSU). Analysis made significant use of PYTHON 3.7.4, and the associated packages NUMPY, MATPLOTLIB, SEABORN, SCIPY. Data reduction made significant use of MIDAS FEROS (Stahl et al. 1999), ECHELLE (Ballester 1992), PYSALT (Crawford et al. 2010), and IRAF (Tody 1986, 1993).

## DATA AVAILABILITY

The data are available as online material and can be found here: <https://www.dropbox.com/s/n2qxklfizgyuok/Gaia22alz.zip?dl=0>

## REFERENCES

- Abril J., Schmidtbreick L., Ederoclite A., López-Sanjuan C., 2020, *MNRAS*, 492, L40
- Arai A., Isogai M., Yamanaka M., Akitaya H., Uemura M., 2015, *Acta Polytechnica CTU Proc.*, 2, 257
- Aydi E. et al., 2018, *MNRAS*, 480, 572
- Aydi E. et al., 2019, preprint ([arXiv:1903.09232](https://arxiv.org/abs/1903.09232))
- Aydi E., Chomiuk L., Sokolovsky K. V., Steinberg E., 2020a, *Nature Astron.*, 2, 697
- Aydi E. et al., 2020b, *ApJ*, 905, 62
- Aydi E. et al., 2021, *Astron. Telegram*, 14710, 1
- Aydi E. et al., 2022a, *ApJ*, 939, 6
- Aydi E. et al., 2022b, *Astron. Telegram*, 15355, 1
- Aydi E. et al., 2022c, *Astron. Telegram*, 15395, 1
- Ballester P., 1992, in Grosbøl P. J., de Ruijscher R. C. E. eds, *European Southern Observatory Conference and Workshop Proceedings Vol. 41*, European Southern Observatory Conference and Workshop Proceedings. p. 177
- Barnes S. I. et al., 2008, The optical design of the Southern African Large Telescope high resolution spectrograph: SALT HRS, in *Ground-based and Airborne Instrumentation for Astronomy II. Proc. SPIE 7014*, p. 70140K.
- Belczyński K., Mikołajewska J., Munari U., Ivison R. J., Friedjung M., 2000, *A&AS*, 146, 407
- Bode M. F., Evans A., 2008, in Bode M. F., Evans A. eds, *Classical Novae*. Cambridge Astrophysics Series Vol. 43, Cambridge University Press, Cambridge, UK
- Bode M. F., Seaquist E. R., Evans A., 1987, *MNRAS*, 228, 217
- Bonning E. W., Shields G. A., Stevens A. C., Salviander S., 2013, *ApJ*, 770, 30
- Booth R. A., Mohamed S., Podsiadlowski P., 2016, *MNRAS*, 457, 822
- Bowen G. H., 1988, *ApJ*, 329, 299
- Bramall D. G. et al., 2010, in *The SALT HRS spectrograph: final design, instrument capabilities, and operational modes*, *Ground-based and Airborne Instrumentation for Astronomy III*. p. 77354F
- Bramall D. G. et al., 2012, in *The SALT HRS spectrograph: instrument integration and laboratory test results*, *Ground-based and Airborne Instrumentation for Astronomy IV. Proc. SPIE 8446*, p. 84460A
- Brandt E., Quiroga C., Mikołajewska J., Ferrer O. E., García L. G., 2009, *A&A*, 497, 815
- Breedt E. et al., 2014, *MNRAS*, 443, 3174
- Brink J. et al., 2022, *Astron. Telegram*, 15270, 1
- Bruch A., 1989, *A&AS*, 78, 145
- Bruch A., Engel A., 1994, *A&AS*, 104, 79
- Bruch A., Schimpke T., 1992, *A&AS*, 93, 419
- Buckley D. A. H., Swart G. P., Meiring J. G., 2006, in *Completion and commissioning of the Southern African Large Telescope*, *Society of Photo-Optical Instrumentation Engineers (SPIE) Conference Series. Proc. SPIE 6267*, p. 62670Z
- Burgh E. B., Nordsieck K. H., Kobulnicky H. A., Williams T. B., O'Donoghue D., Smith M. P., Percival J. W., 2003, in Iye M., Moorwood A. F. M. eds, *Proc. SPIE Conf. Ser. Vol. 4841, Instrument Design and Performance for Optical/Infrared Ground-based Telescopes*. SPIE, Bellingham, p. 1463
- Cannizzo J. K., 1993, *ApJ*, 419, 318
- Cao Y. et al., 2012, *ApJ*, 752, 133
- Chen B. Q. et al., 2019, *MNRAS*, 483, 4277
- Chochol D., Pribulla T., Parimucha Š., Vaňko M., 2003, *Balt. Astron.*, 12, 610
- Chomiuk L. et al., 2014, *Nature*, 514, 339
- Chomiuk L., Metzger B. D., Shen K. J., 2021, *ARA&A*, 59, 391
- Clemens J. C., Crain J. A., Anderson R., 2004, in Moorwood A. F. M., Iye M. eds, *Proc. SPIE Conf. Ser. Vol. 5492, Ground-based Instrumentation for Astronomy*. SPIE, Bellingham, p. 331
- Crause L. A. et al., 2014, in *Performance of the Southern African Large Telescope (SALT) High Resolution Spectrograph (HRS)*, *Ground-based and Airborne Instrumentation for Astronomy V. Proc. SPIE 9147*, p. 91476T

- Crawford S. M. et al., 2010, in *PySALT: the SALT science pipeline, Observatory Operations: Strategies, Processes, and Systems III*. Proc. SPIE 7737, p. 773725
- Della Valle M., Izzo L., 2020, *A&AR*, 28, 3
- Dobrzycka D., Kenyon S. J., 1994, *AJ*, 108, 2259
- Downes R. A., Webbink R. F., Shara M. M., Ritter H., Kolb U., Duerbeck H. W., 2001, *PASP*, 113, 764
- Draine B. T., 2009, in Henning T., Grün E., Steinacker J.eds, 414, ASP Conf. Ser. Vol. 414, *Cosmic Dust-Near and Far*. Astron. Soc. Pac., San Francisco, p. 453
- Drew J. E. et al., 2014, *MNRAS*, 440, 2036
- Ederoclite A., 2014, in Woudt P. A., Ribeiro V. A. R. M.eds, ASP Conf. Ser. Vol. 490, *Stellar Novae: Past and Future Decades*. Astron. Soc. Pac., San Francisco, p. 163
- Friedjung M., 1992, *A&A*, 262, 487
- Friedman S. D. et al., 2011, *ApJ*, 727, 33
- Gaia Collaboration, 2021, *A&A*, 649, A1
- Gaia Collaboration et al., 2023, *A&A*, 674, A1
- Gaposchkin C. H. P., 1957, *The galactic novae*, Amsterdam, North-Holland, Pub. Co., Interscience Publishers, New York
- Godon P., Sion E. M., Williams R. E., Starrfield S., 2018, *ApJ*, 862, 89
- Goranskij V. P. et al., 2007, *Astrophys. Bull.*, 62, 125
- Hachisu I., Kato M., 2004, *ApJ*, 612, L57
- Hachisu I., Kato M., 2014, *ApJ*, 785, 97
- Hachisu I., Kato M., 2022, *ApJ*, 939, 1
- Han Z., Boonruksar S., Qian S., Xiaohui F., Wang Q., Zhu L., Dong A., Zhi Q., 2020, *PASJ*, 72, 76
- Harrison T. E., Johnson J. J., Spyromilio J., 1993, *AJ*, 105, 320
- Harvey E. J., Redman M. P., Darnley M. J., Williams S. C., Berdyugin A., Pirola V. E., Fitzgerald K. P., O'Connor E. G. P., 2018, *A&A*, 611, A3
- Hillman Y., Prialnik D., Kovetz A., Shara M. M., Neill J. D., 2014, *MNRAS*, 437, 1962
- Hodgkin S. T. et al., 2022, *Transient Name Serv. Discovery Rep.*, 2022-313, 1
- Hou W., Luo A. L., Li Y.-B., Qin L., 2020, *AJ*, 159, 43
- Hou W., Luo A. L., Dong Y.-Q., Chen X.-L., Bai Z.-R., 2023, *AJ*, 165, 148
- Hounsell R. et al., 2010, *ApJ*, 724, 480
- Hounsell R. et al., 2016, *ApJ*, 820, 104
- Izzo L. et al., 2012, *Mem. Soc. Astron. Ital.*, 83, 830
- Jiang B., Luo A., Zhao Y., Wei P., 2013, *MNRAS*, 430, 986
- Kafka S., 2020, *Obs. AAVSO Int. Database*, <https://www.aavso.org>
- Kato M., Hachisu I., 2011, *ApJ*, 743, 157
- Kato T., Takamizawa K., 2001, *Inf. Bull. Var. Stars*, 5100
- Kato T., Maehara H., Uemura M., 2012, *PASJ*, 64, 63
- Kato M., Hachisu I., Henze M., 2013, *ApJ*, 779, 19
- Kato M., Saio H., Hachisu I., 2022, *PASJ*, 74, 1005
- Kawash A. et al., 2021, *ApJ*, 910, 120
- Kenyon S. J., Truran J. W., 1983, *ApJ*, 273, 280
- Kniazev A. Y., Gvaramadze V. V., Berdnikov L. N., 2016, *MNRAS*, 459, 3068
- Kobulnicky H. A., Nordsieck K. H., Burgh E. B., Smith M. P., Percival J. W., Williams T. B., O'Donoghue D., 2003, in Iye M., Moorwood A. F. M.eds, 4841, *Proc. SPIE Conf. Ser. Vol. 4841, Instrument Design and Performance for Optical/Infrared Ground-based Telescopes*. SPIE, Bellingham, p. 1634
- Kochanek C. S. et al., 2017, *PASP*, 129, 104502
- König O. et al., 2022, *Nature*, 605, 248
- Krautter J., 2002, in Hernanz M., José J.eds, AIP Conf. Ser. Vol. 637, *Classical Nova Explosions*. Am. Inst. Phys., New York, p. 345
- Luna G. J. M. et al., 2018, *A&A*, 619, A61
- Marshall D. J., Robin A. C., Reylé C., Schultheis M., Picaud S., 2006, *A&A*, 453, 635
- Matsumoto T., Metzger B. D., 2022, *ApJ*, 938, 5
- McLaughlin D. B., 1944, *Pop. Astron.*, 52, 109
- McLaughlin D. B., 1947, *PASP*, 59, 244
- Mikolajewska J., 2003, in Corradi R. L. M., Mikolajewska J., Mahoney T. J.eds, ASP Conf. Ser. Vol. 303, *Symbiotic Stars Probing Stellar Evolution*. Astron. Soc. Pac., San Francisco, p. 9
- Mikolajewska J., 2007, *Balt. Astron.*, 16, 1
- Mikolajewska J., 2010, preprint ([arXiv:1011.5657](https://arxiv.org/abs/1011.5657))
- Mikolajewska J., Ikiewicz K., Gałan C., Monard B., Otulakowska-Hypka M., Shara M. M., Udalski A., 2021, *MNRAS*, 504, 2122
- Mondal A., Anupama G. C., Kamath U. S., Das R., Selvakumar G., Mondal S., 2018, *MNRAS*, 474, 4211
- Morales-Rueda L., Marsh T. R., 2002, *MNRAS*, 332, 814
- Morisset C., Pequignot D., 1996, *A&A*, 312, 135
- Mróz P. et al., 2016, *Nature*, 537, 649
- Munari U., Zwitter T., 1997, *A&A*, 318, 269
- Munari U. et al., 1996, *A&A*, 315, 166
- Munari U., Ochner P., Siviero A., Jones A., Moretti S., Tomaselli S., Dallaporta S., 2009, *Balt. Astron.*, 18, 75
- Munari U., Ribeiro V. A. R. M., Bode M. F., Saguner T., 2011a, *MNRAS*, 410, 525
- Munari U. et al., 2011b, *MNRAS*, 410, L52
- Munari U., Dallaporta S., Cherini G., 2016, *New Atron.*, 47, 7
- Murphy-Glasyher F. J. et al., 2022, *MNRAS*, 514, 6183
- O'Brien T. J. et al., 2006, *Nature*, 442, 279
- O'Donoghue D. et al., 2006, *MNRAS*, 372, 151
- Orio M. et al., 2020, *ApJ*, 895, 80
- Pastorello A. et al., 2019, *A&A*, 630, A75
- Patterson J. et al., 1998, *PASP*, 110, 380
- Pejcha O., 2009, *ApJ*, 701, L119
- Pejcha O., Metzger B. D., Tomida K., 2016, *MNRAS*, 461, 2527
- Pejcha O., Metzger B. D., Tyles J. G., Tomida K., 2017, *ApJ*, 850, 59
- Prialnik D., 1986, *ApJ*, 310, 222
- Salazar I. V., LeBleu A., Schaefer B. E., Landolt A. U., Dvorak S., 2017, *MNRAS*, 469, 4116
- Schaefer B. E., 2022, *MNRAS*, 517, 6150
- Schaefer B. E., Collazzi A. C., 2010, *AJ*, 139, 1831
- Schaefer B. E. et al., 2013, *ApJ*, 773, 55
- Schlafly E. F., Finkbeiner D. P., 2011, *ApJ*, 737, 103
- Schlafly E. F. et al., 2018, *ApJS*, 234, 39
- Shafter A. W., 2017, *ApJ*, 834, 196
- Shappee B. J. et al., 2014, *ApJ*, 788, 48
- Shara M. M. et al., 2017, *ApJ*, 839, 109
- Shen K. J., Quataert E., 2022, *ApJ*, 938, 31
- Shore S. N., 2014, in Woudt P. A., Ribeiro V. A. R. M.eds, ASP Conf. Ser. Vol. 490, *Stellar Novae: Past and Future Decades*. Astron. Soc. Pac., San Francisco, p. 145
- Shore S. N., Kuin N. P., Mason E., De Gennaro Aquino I., 2018, *A&A*, 619, A104
- Shu F. H., Lubow S. H., Anderson L., 1979, *ApJ*, 229, 223
- Smale A. P. et al., 1988, *MNRAS*, 233, 51
- Stahl O., Kaufer A., Tubbesing S., 1999, in Guenther E., Stecklum B., Klose S.eds, ASP Conf. Ser. Vol. 188, *Optical and Infrared Spectroscopy of Circumstellar Matter*. Astron. Soc. Pac., San Francisco, p. 331
- Starrfield S., 1989, in Bode M. F., Evans A.eds, *Classical novae*, J. Wiley, Chichester, UK, p. 39
- Starrfield S., Sparks W. M., Truran J. W., 1974, *ApJS*, 28, 247
- Starrfield S., Truran J. W., Sparks W. M., Krautter J., MacDonald J., 1990, in Cassatella A., Viotti R.eds, *Proceedings of Colloquium No.122 of the International Astronomical Union, Physics of Classical Novae*. Vol. 369, Springer-Verlag, Berlin, Germany, New York, NY, p. 306
- Strope R. J., Schaefer B. E., Henden A. A., 2010, *AJ*, 140, 34
- Szkody P., 1994, *AJ*, 108, 639
- Szkody P. et al., 2011, *AJ*, 142, 181
- Tanaka J., Nogami D., Fujii M., Ayani K., Kato T., 2011a, *PASJ*, 63, 159
- Tanaka J., Nogami D., Fujii M., Ayani K., Kato T., Maehara H., Kiyota S., Nakajima K., 2011b, *PASJ*, 63, 911
- Tody D., 1986, in Crawford D. L.eds, *Proc. SPIE Conf. Ser. Vol. 627, Instrumentation in astronomy VI*. SPIE, Bellingham, p. 733

- Tody D., 1993, in Hanisch R. J., Brissenden R. J. V., Barnes J. eds, ASP Conf. Ser. Vol. 52, Astronomical Data Analysis Software and Systems II. Astron. Soc. Pac., San Francisco, p. 173
- Udalski A., Szymański M. K., Szymański G., 2015, *Acta Astron.*, 65, 1
- Vogt N., 1990, *ApJ*, 356, 609
- Walter F. M., 2016, in Conference on Shocks and Particle Acceleration in Novae and Supernovae. <https://www.simonsfoundation.org/event/conference-on-shocks-and-particle-acceleration-in-novae-and-supernovae/> (Accessed June 8th, 2023)
- Walter F. M., Battisti A., Towers S. E., Bond H. E., Stringfellow G. S., 2012, *PASP*, 124, 1057
- Wang S., Chen X., 2019, *ApJ*, 877, 116
- Warner B., 1987, *MNRAS*, 227, 23
- Warner B., 1995, *Cataclysmic variable stars*. Cambridge Astrophysics Series Vol. 28, Cambridge University Press, Cambridge, UK
- Warner B., 2008, in Bode M. F., Evans A. eds, *Classical Novae*, 2nd Edition. Cambridge Astrophysics Series, No. 43, Cambridge University Press, Cambridge, p. 16
- Williams R. E., 1992, *AJ*, 104, 725
- Williams R. E., 2012, *AJ*, 144, 98

- Williams S. C., Darnley M. J., Bode M. F., Steele I. A., 2015, *ApJ*, 805, L18
- Wolf W. M., Bildsten L., Brooks J., Paxton B., 2013, *ApJ*, 777, 136
- Yaron O., Prialnik D., Shara M. M., Kovetz A., 2005, *ApJ*, 623, 398
- Zemko P., Mukai K., Orio M., 2015, *ApJ*, 807, 61
- Zemko P., Orio M., Mukai K., Bianchini A., Ciroi S., Cracco V., 2016, *MNRAS*, 460, 2744
- Zwitter T., Munari U., 1995, *A&AS*, 114, 575
- van den Heuvel E. P. J., Bhattacharya D., Nomoto K., Rappaport S. A., 1992, *A&A*, 262, 97

## SUPPORTING INFORMATION


Supplementary data are available at *MNRAS* online.

### online\_material.zip


Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

## APPENDIX A: SUPPLEMENTARY PLOTS AND TABLES

In this section, we present supplementary plots and tables.




**Figure A1.** *Left:* the complete ASAS-SN optical *g*-band light curve of nova Gaia22alz up until the time of the writing of this paper. *Right:* the *I*-band light curve of the 2019 outburst of V1047 Cen, adopted from Aydi et al. (2022a).




**Figure A2.** The spectral evolution of nova Gaia22alz representing the different spectral stages: early-rise (blue), mid-rise (red), late-rise (green), and post-optical-peak (black). Numbers between brackets are days after  $t_0$ . The resolving power  $R$  of each spectrum is added on the left-hand side of each panel.





**Figure A3.** The spectral evolution of nova Gaia22alz representing the different spectral stages: early-rise (blue), mid-rise (red), late-rise (green), and post-optical-peak (black). Numbers between brackets are days after  $t_0$ . The resolving power  $R$  of each spectrum is added on the left-hand side of each panel.



**Figure A4.** The spectral evolution of nova Gaia22alz representing the different spectral stages: early-rise (blue), mid-rise (red), late-rise (green), and post-optical-peak (black). Numbers between brackets are days after  $t_0$ . The resolving power  $R$  of each spectrum is added on the left-hand side of each panel.

**Table A1.** Optical spectroscopic observations log.

| Telescope | Instrument | date       | $t - t_0$<br>(d) | Resolving power | $\lambda$ Range<br>(Å) |
|-----------|------------|------------|------------------|-----------------|------------------------|
| SALT      | RSS        | 2022-03-10 | 45               | 1500            | 4200–7300              |
| SALT      | HRS        | 2022-03-15 | 49               | 14000           | 4000–9000              |
| SALT      | HRS        | 2022-04-03 | 68               | 14000           | 4000–9000              |
| SOAR      | Goodman    | 2022-04-09 | 76               | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-04-10 | 76               | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-04-14 | 80               | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-04-21 | 86               | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-04-21 | 86               | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-05-01 | 96               | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-05-01 | 96               | 4400            | 4500–5170              |
| SALT      | HRS        | 2022-05-08 | 103              | 14000           | 4000–9000              |
| SALT      | HRS        | 2022-05-11 | 106              | 14000           | 4000–9000              |
| SOAR      | Goodman    | 2022-05-22 | 118              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-05-22 | 118              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-05-24 | 120              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-05-24 | 120              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-06-09 | 135              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-06-09 | 135              | 4400            | 4500–5170              |
| Magellan  | IMACS      | 2022-07-21 | 178              | 1500            | 3800–6600              |
| SOAR      | Goodman    | 2022-08-15 | 203              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-08-15 | 203              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-08-24 | 212              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-08-24 | 212              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-09-17 | 236              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-09-17 | 236              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-10-08 | 257              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-10-08 | 257              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-10-09 | 258              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-10-18 | 267              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-10-18 | 267              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-10-29 | 278              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-10-29 | 278              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-12-01 | 310              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-12-01 | 310              | 4400            | 4500–5170              |
| SOAR      | Goodman    | 2022-12-22 | 331              | 1100            | 3800–7800              |
| SOAR      | Goodman    | 2022-12-22 | 331              | 4400            | 4500–5170              |

**APPENDIX B: ANALYTICAL MODEL OF STATIONARY SHOCK WITHIN THE L2 OUTFLOW**

In this section, we present a set of calculations supporting the main, alternative model with suggest in Section 4. A binary of period  $P$  and mass  $M$  has a semimajor axis given by

$$a \approx 3.5 \times 10^{10} \text{cm} \left( \frac{M}{M_\odot} \right)^{1/3} \left( \frac{P}{\text{h}} \right)^{2/3}, \quad (\text{B1})$$

with binary escape speed

$$v_{\text{esc}} \approx \left( \frac{2GM}{a} \right)^{1/2} \approx 870 \text{ km s}^{-1} \left( \frac{M}{M_\odot} \right)^{1/3} \left( \frac{P}{\text{h}} \right)^{-1/3}. \quad (\text{B2})$$

Consider a mass-loss from the  $L_2$  point at a rate  $\dot{M}_L$ , which we assume is monotonically rising after the first few days of the eruption. Shu, Lubow & Anderson (1979) and Pejcha et al. (2016) show that cold mass-loss from the  $L_2$  point for a binary of  $0.064 \lesssim q \equiv M_*/M_{\text{WD}} \lesssim 0.78$  reaches a terminal velocity up to

$$v_{L2, \text{min}} \approx v_{\text{esc}}/4 \approx 220 \text{ km s}^{-1} \left( \frac{M}{M_\odot} \right)^{1/3} \left( \frac{P}{\text{h}} \right)^{-1/3}. \quad (\text{B3})$$

Higher velocities are predicted for matter ejected with finite thermal temperature (Pejcha et al. 2017), so we take  $v_{L2} \approx 2v_{L2, \text{min}}$ , to match the  $400 \text{ km s}^{-1}$  line widths observed in Gaia22alz. The vertical surface density and optical depth of the equatorial outflow at radius  $r \gg a$  from the binary is given by

$$\Sigma \approx \frac{\dot{M}_{L2}}{2\pi r v_{L2}}; \quad \tau_T \approx \frac{\dot{M}_{L2} \kappa_T}{2\pi r v_{L2}} \approx 1.0 \left( \frac{\dot{M}_{L2}}{10^{-7} M_\odot \text{wk}^{-1}} \right) \left( \frac{r}{10a} \right)^{-1} \left( \frac{M}{M_\odot} \right)^{-2/3} \left( \frac{P}{\text{h}} \right)^{-1/3} \quad (\text{B4})$$

where  $\kappa_T \approx 0.34 \text{ cm}^2 \text{ g}^{-1}$  is the Thomson optical depth. The  $L_2$  streams collide at a radius  $r_{\text{coll}} \approx 10a_{\text{bin}}$ , with a relative velocity  $\Delta v \approx 0.2v_{L2}$ , shock-heating the material to a temperature

$$T_c \simeq \frac{3}{16} \frac{m_p}{k} (\Delta v)^2 \approx 1.7 \times 10^5 \text{ K} \left( \frac{M}{M_\odot} \right)^{2/3} \left( \frac{P}{\text{h}} \right)^{-2/3}, \quad (\text{B5})$$

high enough to (re-)ionize the gas. The mid-plane density on the radial scale of the collision can likewise be estimated as Pejcha et al. (2016)

$$\rho_c \sim \frac{\Sigma}{2a_{\text{bin}}} \sim 5 \times 10^{-11} \text{ g cm}^{-3} \left( \frac{\dot{M}_{L2}}{10^{-7} M_\odot \text{ wk}^{-1}} \right) \left( \frac{r_{\text{coll}}}{10a} \right)^{-1} \left( \frac{M}{M_\odot} \right)^{-1} \left( \frac{P}{\text{h}} \right)^{-1}. \quad (\text{B6})$$

For typical values of  $(\rho_c, T_c)$ , we estimate the opacity (e.g. from OPAL) to be close to the Thomson value assumed above (e.g. fig. 3 of Matsumoto & Metzger 2022; see also below). The kinetic luminosity of the shocks is given by

$$L_{L2} \approx \dot{M}_{L2} v_{L2} \Delta v \approx 1.3 \times 10^{35} \text{ erg s}^{-1} \left( \frac{\dot{M}_{L2}}{10^{-7} M_\odot \text{ wk}^{-1}} \right) \left( \frac{M}{M_\odot} \right)^{2/3} \left( \frac{P}{\text{h}} \right)^{-2/3}, \quad (\text{B7})$$

while the radiative cooling time of the gas,

$$t_{\text{cool}} \sim \frac{3kT_c}{(4\rho_c/m_p)\Lambda(T_c)} \sim 10^{-3} \text{ s} \left( \frac{\dot{M}_{L2}}{10^{-7} M_\odot \text{ wk}^{-1}} \right)^{-1}, \quad (\text{B8})$$

is extremely short, where  $\Lambda(T) \simeq 2.8 \times 10^{-18} (T/\text{K})^{-0.7} \text{ erg s}^{-1} \text{ cm}^3$  is the atomic line cooling rate (e.g. Draine 2009).

Using equation (B4 and B7), we can eliminate  $\dot{M}_{L2}$  to write

$$L_{L2} \approx 10^{35} \text{ erg s}^{-1} \left( \frac{\tau_T}{1} \right) \left( \frac{M}{M_\odot} \right)^{4/3} \left( \frac{P}{\text{hr}} \right)^{-1/3}, \quad (\text{B9})$$

Up to what value of  $\tau_T$  can we still see line emission from the shock-heated gas? Considering that the free-free opacity for an optical photon (e.g. H II) of energy  $\epsilon_\gamma \sim 1 \text{ eV}$ ,

$$\kappa_{\text{ff}} \approx 0.35 \text{ cm}^2 \text{ g}^{-1} \left( \frac{T}{10^5 \text{ K}} \right)^{-3/2} \left( \frac{\rho_c}{10^{-10} \text{ g cm}^{-3}} \right) \left( \frac{\epsilon_\gamma}{1 \text{ eV}} \right)^{-2} \bar{g}_{\text{ff}}, \quad (\text{B10})$$

is close to the Thomson opacity. Therefore, the effective absorption optical depth  $\tau_{\text{th}} \simeq \sqrt{\tau_{\text{ff}}(\tau_T + \tau_{\text{ff}})}$  is close to  $\tau_T$ . In other words, we expect to observe line emission up to  $\tau_T \sim 1$ , or  $L_{L2} \sim 10^{35} \text{ erg s}^{-1}$ .

<sup>1</sup>Center for Data Intensive and Time Domain Astronomy, Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824, USA

<sup>2</sup>Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, Bartycka 18, PL 00-716 Warsaw, Poland

<sup>3</sup>South African Astronomical Observatory, P.O. Box 9, 7935 Observatory, South Africa

<sup>4</sup>Department of Astronomy, University of Cape Town, Private Bag X3, Rondebosch 7701, South Africa

<sup>5</sup>Department of Physics and Columbia Astrophysics Laboratory, Columbia University, New York, NY 10027, USA

<sup>6</sup>Center for Computational Astrophysics, Flatiron Institute, 162 5th Ave, New York, NY 10010, USA

<sup>7</sup>UK Astronomy Technology Centre, Royal Observatory, Blackford Hill, Edinburgh, EH9 3HJ, UK

<sup>8</sup>Carnegie Observatories, 813 Santa Barbara Street, Pasadena, CA 91101, USA

<sup>9</sup>DARK, Niels Bohr Institute, University of Copenhagen, Jagtvej 128, D-2200 Copenhagen Ø, Denmark

<sup>10</sup>National Radio Astronomy Observatory, P.O. Box O, Socorro, NM 87801, USA

<sup>11</sup>INAF-Osservatorio Astronomico di Trieste, Via G.B. Tiepolo 11, I-34143 Trieste, Italy

<sup>12</sup>Institute of Fundamental Physics of the Universe, Via Beirut 2, Miramare, 34151 Trieste, Italy

<sup>13</sup>Astronomical Observatory, University of Warsaw, Al. Ujazdowskie 4, PL-00-478 Warszawa, Poland

<sup>14</sup>CRESST and X-ray Astrophysics Laboratory, NASA/GSFC, Greenbelt, MD 20771, USA

<sup>15</sup>Department of Physics, University of Maryland, Baltimore County, 1000 Hilltop Circle, Baltimore, MD 21250, USA

<sup>16</sup>INAF-Osservatorio di Padova, vicolo dell'Osservatorio 5, I-35122 Padova, Italy

<sup>17</sup>Department of Astronomy, University of Wisconsin, 475 N. Charter St., Madison, WI 53704, USA

<sup>18</sup>Institute for Astronomy, University of Hawai'i, 2680 Woodlawn Drive, Honolulu, HI 96822, USA

<sup>19</sup>Department of Astronomy and Theoretical Astrophysics Center, University of California, Berkeley, CA 94720, USA

<sup>20</sup>Columbia Astrophysics Laboratory and Department of Physics, Columbia University, New York, NY 10027, USA

<sup>21</sup>Department of Astronomy, University of Illinois at Urbana-Champaign, 1002 W. Green Street, Urbana, IL 61801, USA

<sup>22</sup>Sternberg Astronomical Institute, Moscow State University, Universitetskii pr. 13, 119992 Moscow, Russia

<sup>23</sup>Department of Astronomy & Astrophysics, University of California, Santa Cruz, 1156 High Street, Santa Cruz, CA 95064, USA

<sup>24</sup>Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218, USA

This paper has been typeset from a  $\text{\LaTeX}$  file prepared by the author.