

This article may be downloaded for personal use only. Any other use requires prior permission of the author and AIP Publishing. This article appeared in Bodaghee, A., J. A. Tomsick, J. Rodriguez, S. Chaty, K. Pottschmidt, R. Walter, and P. Romano. "Supergiant X-Ray Binaries Observed by Suzaku." AIP Conference Proceedings 1427, no. 1 (March 29, 2012): 52–55. <https://doi.org/10.1063/1.3696150>. and may be found at <https://doi.org/10.1063/1.3696150>.

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Supergiant X-ray binaries observed by *Suzaku*

A. Bodaghee*, J. A. Tomsick*, J. Rodriguez†, S. Chaty†, K. Pottschmidt**,
R. Walter‡ and P. Romano§

*Space Sciences Laboratory, University of California, Berkeley, USA

†Lab. AIM, CEA/IRFU - UPD – CNRS/INSU, CEA DSM/IRFU/SaP, Centre de Saclay, France

**CRESST/UMBC, NASA Goddard Space Flight Center, USA

‡ISDC, Observatoire de l'Université de Genève, Switzerland

§INAF, Istituto di Astrofisica Spaziale e Fisica Cosmica, Palermo, Italy

Abstract. *Suzaku* observations are presented for the high-mass X-ray binaries IGR J16207–5129 and IGR J17391–3021. For IGR J16207–5129, we provide the first X-ray broadband (0.5–60 keV) spectrum from which we confirm a large intrinsic column density ($N_{\text{H}} = 1.6 \times 10^{23} \text{ cm}^{-2}$), and we constrain the cutoff energy for the first time ($E_{\text{cut}} = 19 \text{ keV}$). A prolonged ($> 30 \text{ ks}$) attenuation of the X-ray flux was observed which we tentatively attribute to an eclipse of the probable neutron star by its massive companion, in a binary system with an orbital period between 4 and 9 days, and inclination angles > 50 degrees. For IGR J17391–3021, we witnessed a transition from quiescence to a low-activity phase punctuated by weak flares whose peak luminosities in the 0.5–10 keV band are only a factor of 5 times that of the pre-flare emission. These micro flares are accompanied by an increase in N_{H} which suggests the accretion of obscuring clumps of wind. We now recognize that these low-activity epochs constitute the most common emission phase for this system, and perhaps in other supergiant fast X-ray transients (SFXTs) as well. We close with an overview of our upcoming program in which *Suzaku* will provide the first ever observation of an SFXT (IGR J16479–4514) during a binary orbit enabling us to probe the accretion wind at every phase.

Keywords: accretion, accretion disks ; gamma-rays: general ; stars: neutron ; supergiants ; X-rays: binaries ; X-rays: individual (IGR J16207–5129; IGR J17391–3021 = XTE J1739–302)

PACS: 98.70.Qy, 97.80.Jp, 97.60.Jd, 97.20.Pm, 97.10.Gz

INTRODUCTION & OBSERVATIONS

High-mass X-ray binaries (HMXBs) are compact objects (usually a neutron star but sometimes a black hole) which accrete from a massive donor star. An HMXB can be further classified based on whether this donor is a B-emission line star (BEXBs) or a supergiant OB star (SGXBs). Recently, another subclass of HMXB has been recognized wherein the donor star is clearly a supergiant OB star as in SGXBs, but the X-ray emission shows a large degree of variability as in BEXBs. These systems are called supergiant fast X-ray transients (SFXTs).

Suzaku, with its high sensitivity over a broad X-ray bandpass, can help resolve several outstanding issues concerning the accretion physics and nature of HMXBs. We point the reader to *Suzaku* observations which: confirmed the presence of clumpy winds in the SFXT called IGR J17544–2619 [1]; provided the most accurate broadband X-ray spectrum of the highly-obscured IGR J16318–4848 [2]; and which found a low-level of activity in the SFXT IGR J08408–4503 even during quiescence [3]. In 2008, *Suzaku* observed the SGXB IGR J16207–5129 and the SFXT IGR J17391–3021 (= XTE J1739–302) for 33 and 37 ks of effective exposure time, respectively. The analysis of these sources is detailed in Bodaghee et al. [4, 5] and is summarized below.

IGR J16207–5129

Figure 1 shows that the emission from IGR J16207–5129 varies within 1–2 orders of magnitude on ks timescales, which is typical of SGXBs. Intriguingly, X-rays from the source, and the variability associated with that emission, are significantly diminished during the last ~ 30 ks of the observation. If the accretor were to enter an “off” state, this would last a few 100 s, or at least an order of magnitude shorter than the period of inactivity that we observed. The observed attenuation of the X-ray flux can not be explained by an increase in the column density (i.e., occulting wind clumps) since the measured N_{H} does not change in a significant way between O2a and O2b (Table 1). An occultation of the neutron star would imply unrealistic column densities at other orbital phases. A remaining viable explanation is an eclipse of the compact object by its companion, but the data do not allow us to confirm or reject this hypothesis. If the eclipse is real, it lasts at least 30 ks (0.35 d) setting a lower limit on the orbital period at 0.7 d. With this constraint, and assuming typical stellar parameters, a model for an eclipsing binary [6] yields orbital periods between 4 and 9 d with inclination angles $> 50^\circ$. We note that whatever mechanism is involved in driving the sporadic emission in SFXTs (e.g., the 33-ks dormant period of IGR J17391–3021 in Fig. 2) could be responsible for the prolonged attenuation seen in IGR J16207–5129. The spectrum of IGR J16207–5129 (Fig. 1) features a large absorbing column ($N_{\text{H}} > 10^{23} \text{ cm}^{-2}$) and a possible iron line, both of which suggest the presence of matter around the X-ray source. A cutoff near 20 keV favors a neutron star as the compact object, but coherent pulsations or cyclotron absorption lines that would confirm the presence of a neutron star are not detected.




FIGURE 1. *Left:* Background-subtracted light curve of IGR J16207–5129 from XIS (red: 0.5–10 keV) and HXD (blue: 15–60 keV). The upper panel displays the first observation (O1) while the lower panel shows the second observation (O2) which includes simultaneous data from XIS. Each bin collects 160 s worth of data. The dashed line represents MJD 54527.382. This corresponds to the onset of an unusually long period (> 30 ks) of suppressed flux which could represent an eclipse of the compact object by its supergiant stellar companion. *Right:* Spectrum of IGR J16207–5129 corrected for the background and fit with an absorbed power law with an exponential cutoff whose parameters are listed in Table 1. Each bin collects at least 150 photon counts from XIS-1 (red), XIS-0 paired with XIS-3 (blue), and PIN (black).

IGR J17391–3021

During the initial 33 ks of the observation of IGR J17391–3021, the source is in an extremely low-activity phase that is at or near quiescence (Fig. 2). The source then enters a period of enhanced activity in which the luminosity is only a factor of 5 that of the “quiescent” emission. Bozzo et al. [8] noticed similar behavior in a recent *XMM-Newton* observation of this source. Observations prior to and after MJD 54518.92 are referred to henceforth as “dormant” and “active” epochs, respectively. The active state features 3 quasi-periodic flares separated by ~ 15 ks intervals. The peak fluxes are at the level of the faint detections (and a few upper limits) from *Swift* monitoring [7]. Epochs of enhanced activity just above quiescence, but well below the bright flaring episodes typical of this class, represent over 60% of all observations, so they are the most common emission state. Given more exposure time, many *Swift* upper limits would be detections suggesting that the duty cycle is even higher. Based on the source ephemeris [9], which places the *XMM-Newton* and *Suzaku* observations at different phases, such low-activity states are not confined to a specific part of the orbit. Table 1 shows that when IGR J17391–3021 is in the active state, the N_{H} is at least twice as high as it is in the dormant phase. This is unlike what was seen by Bozzo et al. [8] who found that the N_{H} remained steady (within the statistical errors) during weak flares while Γ varied by around 50%. Differences in the spectral properties of the flares caught by *Suzaku* and *XMM-Newton* could be




FIGURE 2. *Left:* Light curve of IGR J17391–3021 from XIS, *Swift*-XRT, and *XMM-Newton*. The top panel presents the background-subtracted light curve from our *Suzaku* observation (red crosses, 240-s binning). In the bottom panel, the source light curve from the *Swift* monitoring campaign [7] is shown (blue circles, ~ 1 ks of exposure time per data point) along with their $3\text{-}\sigma$ upper limits for non-detections (green downward arrows). The blue upwards arrows designate the maximum flux of large outbursts detected by *Swift* that are situated beyond the scale of the graph. The average flux from our 37-ks *Suzaku* observation is plotted as a single red cross at the intersection of the dotted lines. The magenta triangle at MJD 54740 corresponds to the average flux in a 31-ks observation with *XMM-Newton* [8]. Fluxes are given in the 0.5–10 keV energy band as observed in units of $10^{-11} \text{ erg cm}^{-2} \text{ s}^{-1}$. *Right:* Background-corrected spectrum of IGR J17391–3021 fit with an absorbed power law. The data represent photon counts from XIS-1 (red), and XIS-0 combined with XIS-3 (blue). Each bin collects a minimum of 150 counts.

TABLE 1. Parameters from absorbed power laws fit to the *Suzaku* spectra of IGR J16207–5129 (0.5–60 keV) and IGR J17391–3021 (0.5–10 keV) for various epochs. The luminosity in the 0.5–10 keV band is corrected for absorption. Errors represent 90% confidence.

source	epoch	$10^{22} \frac{N_H}{\text{cm}^{-2}}$	Γ	$10^{33} \frac{L}{\text{erg s}^{-1}}$	χ^2_{ν}/dof
IGR J16207–5129	total	16±1	0.9±0.2	130	1.05/199
	O2a	19±1	1.3±0.1	240	0.91/182
	O2b	19±5	1.5±0.4	56	0.51/64
IGR J17391–3021	total	3.6±0.4	1.4±0.1	4.8	0.82/66
	dormant	1.0±0.6	1.0±0.3	1.3	0.68/8
	active	4.1±0.5	1.5±0.1	7.4	0.73/54

due to unequal geometric configurations of the system between the observations. The N_H during activity is 2–4 times the interstellar value which suggests that the source is not strongly absorbed (intrinsically) except when the compact object accretes a clump passing along our line of sight [see also 1].

CONCLUSIONS & PERSPECTIVES

The putative eclipse in the persistently-emitting SGXB IGR J16207–5129 awaits confirmation [4]. Micro-flares from IGR J17391–3021 hold valuable clues to the accretion processes of these intriguing transients [5]. The demarcation between SGXBs and SFXTs is not as clear as originally believed, and a few systems such as IGR J16479–4514 might represent an intermediate state. In this respect, our upcoming 150-ks long *Suzaku* observation of IGR J16479–4514 at various phases of its orbit will help us understand the extent to which perceived emissivity differences between these populations stem from their unequal wind and orbital characteristics.

REFERENCES

1. R. A. Rampy, D. M. Smith, and I. Negueruela, *ApJ* **707**, 243–249 (2009), 0904.1189.
2. L. Barragán, J. Wilms, K. Pottschmidt, M. A. Nowak, I. Kreykenbohm, R. Walter, and J. A. Tomsick, *A&A* **508**, 1275–1278 (2009), 0912.0254.
3. L. Sidoli, P. Esposito, and L. Ducci, *MNRAS* **409**, 611–618 (2010), 1007.1091.
4. A. Bodaghee, J. A. Tomsick, J. Rodriguez, S. Chaty, K. Pottschmidt, and R. Walter, *ApJ* **719**, 451–458 (2010), 1006.1911.
5. A. Bodaghee, J. A. Tomsick, J. Rodriguez, S. Chaty, K. Pottschmidt, R. Walter, and P. Romano, *ApJ* **727**, 59 (2011), 1012.0855.
6. S. A. Rappaport, and P. C. Joss, “X-ray pulsars in massive binary systems,” in *Accretion-Driven Stellar X-ray Sources*, edited by W. H. G. Lewin & E. P. J. van den Heuvel, 1983, pp. 1–39.
7. P. Romano, V. La Parola, S. Vercellone, G. Cusumano, L. Sidoli, H. A. Krimm, C. Pagani, P. Esposito, E. A. Hoversten, J. A. Kennea, K. L. Page, D. N. Burrows, and N. Gehrels, *MNRAS* **410**, 1825–1836 (2011), 1009.1146.
8. E. Bozzo, L. Stella, C. Ferrigno, A. Giunta, M. Falanga, S. Campana, G. Israel, and J. C. Leyder, *A&A* **519**, A6 (2010), 1004.2059.
9. S. P. Drave, D. J. Clark, A. J. Bird, V. A. McBride, A. B. Hill, V. Sguera, S. Scaringi, and A. Bazzano, *MNRAS* **409**, 1220–1226 (2010), 1007.3379.