

©2020. American Geophysical Union. All Rights Reserved. Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.



Journal of Geophysical Research: Atmospheres

Supporting Information for

Characterization of aerosol type over East Asia by Multi-angle Imaging SpectroRadiometer (MISR): evaluation and implication

Minghui Tao¹, Jun Wang², Rong Li³, Liangfu Chen⁴, Xiaoguang Xu⁵, Lunche Wang¹, Jinhua Tao⁴, Zifeng Wang⁴, Juan Xiang¹

¹Hubei Key Laboratory of Critical Zone Evolution, School of Geography and Information Engineering, China University of Geosciences, Wuhan 430074, China

²Dept. of Chemical & Environmental Engineering, University of Iowa, Iowa City 52242-1503, USA

³School of Resources and Environmental Science, Hubei University, Wuhan 430062, China

⁴State Key Laboratory of Remote Sensing Science, Institute of Remote Sensing and Digital Earth, Chinese Academy of Sciences, Beijing 100101, China

⁵Joint Center for Earth Systems and Technology, University of Maryland Baltimore County, Baltimore, Maryland 21250, USA

Contents of this file

Figures S1 to S3

Figure S1. Annual frequency of MISR (left) small, (middle) large-mode AOD, and MISR AOD >0.15 (right), respectively.




Figure S2. Annual mean of MISR spherical nonabsorbing AOD with effective radius a) $\leq 0.12 \mu\text{m}$, b) at $0.26 \mu\text{m}$, and c) $2.8 \mu\text{m}$, and d-f) their respective frequency.




Figure S3. Annual mean of MISR AOD in mixture group a) 1-30, b) 31-50, and c) 51-74, and d-f) their frequency, respectively.

