

Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0)

<https://creativecommons.org/licenses/by-nc-nd/4.0/>

DOI:

<https://doi.org/10.1016/j.jmb.2016.10.018>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

**Please provide feedback**

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.



# HHS Public Access

Author manuscript

*J Mol Biol.* Author manuscript; available in PMC 2018 June 30.

Published in final edited form as:

*J Mol Biol.* 2017 June 30; 429(13): 1946–1957. doi:10.1016/j.jmb.2016.10.018.

## Choose your own adventure: The role of histone modifications in yeast cell fate

Deepika Jaiswal<sup>a,b</sup>, Rashi Turniansky<sup>a,c</sup>, and Erin M. Green<sup>a,d</sup>

<sup>a</sup>Department of Biological Sciences, University of Maryland Baltimore County, 1000 Hilltop Circle, Baltimore, MD, USA, 21250

### Abstract

When yeast cells are challenged by a fluctuating environment, signaling networks activate differentiation programs that promote their individual or collective survival. These programs include the initiation of meiotic sporulation, the formation of filamentous growth structures and the activation of programmed cell death pathways. The establishment and maintenance of these distinct cell fates are driven by massive gene expression programs that promote the necessary changes in morphology and physiology. While these genomic reprogramming events depend on a specialized network of transcription factors, a diverse set of chromatin regulators, including histone-modifying enzymes, chromatin remodelers and histone variants, also play essential roles. Here, we review the broad functions of histone modifications in initiating cell fate transitions, with particular focus on their contribution to the control of expression of key genes required for the differentiation programs and chromatin reorganization that accompanies these cell fates.

### Graphical Abstract


---

<sup>d</sup>Corresponding author: egreen@umbc.edu; 410-455-1189.

<sup>b</sup>deej@umbc.edu

<sup>c</sup>iu26299@umbc.edu

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



### Keywords

chromatin; post-translational modifications; meiosis; pseudohyphal growth; programmed cell death

In yeast cells, a series of intrinsic and extrinsic signals initiate cell fate changes in a fluctuating environment. The signals are transduced through integrated networks that ultimately converge on highly-regulated gene expression programs, which are orchestrated by transcription factors, chromatin-modifying enzymes, chromatin remodelers and regulatory RNAs, such as long non-coding RNAs (lncRNAs) [1–3]. The well-characterized genome and tractable genetics of *Saccharomyces cerevisiae* has yielded numerous insights into the genomic reprogramming that occurs in response to diverse environmental signals, and the key mechanisms discovered parallel similar reprogramming events in metazoan systems [4]. Here, we discuss our current understanding of histone modifications and modifying enzymes in *S. cerevisiae* and review the role of chromatin dynamics during key cell fate decisions, including the differentiation of diploids into haploid spores (meiosis and sporulation), the formation of filamentous growth structures and the initiation of programmed cell death pathways. Nutrient availability dictates the cell's decision to enter

either the sporulation or filamentous growth program [5, 6], and reprogramming depends on the tightly-executed activation of key genes, such as *IME1* and *FLO11*, respectively, to drive the required cell cycle and morphological changes. Likewise, activation of programmed cell death pathways occurs in response to diverse cellular stresses in yeast and is accompanied by both transcriptional changes and massive changes to chromatin structure [7]. While these cell fate changes are orchestrated by a dedicated set of sequence-specific transcription factors, histone-modifying enzymes are critical to the reorganization of the chromatin landscape required for full execution of these fates.

## Histone Modifications of *Saccharomyces cerevisiae*

The packaging of DNA into nucleosomes provides the structural foundation for genomic reprogramming events that underlie cell fate transitions. Nucleosomes are comprised of the four core histones H3, H4, H2A and H2B, and may also have one of two variant histones in budding yeast, Htz1 (H2A.Z), which marks heterochromatin-euchromatin boundaries and poised promoters [8, 9], or the centromere-specific H3 variant Cse4 (CENP-A) [10]. The role of individual nucleosomes in defining transcriptional states is determined by their level of histone modification, their position relative to defined sequence elements, the presence of histone variants and their interaction with the transcriptional machinery, all of which are regulated by the coordinated activity of chromatin-associated protein complexes and modifying enzymes [11, 12].

Histone-modifying enzymes, in particular, play a key role in integrating signals at chromatin through the dynamic process of post-translational modification (PTM), including methylation, acetylation, phosphorylation, and ubiquitination [1–3]. The enzymes that add histone PTMs are often referred to as writers and include histone methyltransferases (HMTs), histone acetyltransferases (HATs), kinases, and ubiquitin-protein transferases (UbTs). The marks are removed by erasers- histone demethylases (HDMs), histone deacetylases (HDACs), phosphatases, and deubiquitinases (DUBs)- allowing for dynamic regulation. Effector proteins, known as readers, are recruited to or stabilized at chromatin by these marks and act to transduce signals received at chromatin, governing such pathways as transcription and DNA repair. Reader proteins contain well-defined domains, such as bromodomains, chromodomains and PHD fingers, that recognize specific PTMs on histones [13–15]. Importantly, these domains are often found in writers and erasers, allowing for crosstalk between modifications and integrated signaling networks at chromatin.

As summarized in Table 1, a number of marks on the core histones have been extensively studied using proteomic and molecular techniques, and have well-characterized writers, erasers and readers. These marks, such as H3 and H4 acetylation, H3K4 and H3K36 methylation, are relatively abundant and have been implicated in diverse roles throughout the genome, especially in the realm of gene expression regulation. However, recent advances in proteomic techniques and other methods have revealed the presence of several previously unknown histone PTMs [16–20] that may act in a context-dependent manner. For example, newly-discovered acetylation at H4K44 has been identified in pre-meiotic and meiotic cells and is required for proper double strand break formation and recombination during meiosis [21]. Also, H4K5, K8 and K12 mono-methyl marks, catalyzed by Set5, which are present in

approximately 1% of nucleosomes genome-wide [22], are likely to function in a context-dependent manner. Set5's activity has been linked to stress responses and gene repression in coordination with Set1 [23], however the mechanism by which these methyl marks interpret and transduce signals at chromatin remains unclear.

The genome reprogramming events underlying cell fate decisions rely on both widespread changes to the epigenome, as well as targeted events that may be specific to environmental stimuli or the regulation of individual genomic regions. Marks that appear to be lowly abundant or present only during specialized cellular programs or stress responses may have unique, context-dependent roles in the genome. Additionally, increasing evidence indicates an extensive role for so-called histone-modifying enzymes in the post-translational modification of non-histone proteins, which has been linked to critical signaling pathways [24–28]. The kinetochore protein Dam1 is methylated by Set1, supporting a role for Set1 in chromosome segregation independent of H3K4 methylation [29]. As another example, phosphoenolpyruvate carboxykinase Pck1 is acetylated by the HAT Esa1 and deacetylated by the HDAC Sir2 to dynamically regulate its activity in gluconeogenesis [30]. Furthermore, there are numerous potential histone-modifying enzymes for which no substrates have been identified, such as the candidate methyltransferases Set4 and Set6, and demethylase Ecm5, and also novel modifications, such as glutamate methylation and lysine succinylation [19, 31], for which the functional consequences are largely unclear. Altogether, these observations suggest that we have only begun to scratch the surface regarding the potential roles of histone modifications and histone-modifying enzymes in signaling networks that govern cell fate decisions and genomic reprogramming. Further investigation of these pathways in yeast is expected to reveal principles common to similar events in metazoan systems.

## Nutrient-responsive signaling and chromatin states

Changes to nutrient availability trigger multiple responses that promote survival of yeast cells under fluctuating environmental conditions (Figure 1). For example, nitrogen depletion, the absence of glucose and the presence of a nonfermentable carbon source, such as acetate, stimulates entry into sporulation [5, 32], a specialized cell fate in which *MAT $\alpha$ / $a$*  diploid cells undergo a meiotic division to produce haploid progeny (spores), which are packaged into a protective ascus. Sporulation is characterized by a genetic reprogramming event in which hundreds of genes are induced or repressed in a precisely timed synchrony that controls the progression of meiosis and the morphological changes required for spore formation [33, 34]. Alternatively, a change in environmental conditions in which glucose is abundant, but nitrogen is still limiting, promotes the dimorphic transition of diploid cells from the yeast state to a pseudohyphal state, in which filamentous structures (pseudohyphae) form, adhere to other cells and solid surfaces, such as agar [35]. Similar to sporulation, the filamentous state is also driven by a signaling cascade that activates a gene expression program required for the morphological and physiological changes associated with the new polarized growth structures. In both cases, the nutrient-responsive signaling pathways converge on genes that underlie these cell fate transitions, *IME1* and *FLO11*, respectively. Below, we review the specific role of histone modifications in expression control of *IME1*

and *FLO11*, and highlight how the disruption of chromatin homeostasis alters the maintenance of the acquired cell fates.

### Chromatin-based control of *IME1* expression, the meiotic master regulator

Sporulation occurs in three primary phases, defined as early, middle and late, which are each characterized by discrete events in meiosis and spore formation [5, 33, 36]. When the cell receives the necessary intrinsic (mating type and ploidy) and extrinsic (nutrient availability) signals (Figure 1), the transition from vegetative growth to the early phase of sporulation is marked by the rapid induction of *IME1*, the transcription factor which drives early gene expression required for entry into meiotic S phase [37–39]. The promoter of *IME1* consists of four upstream control sequences (*UCS*) which serve as the integration platform for the positive and negative signals regulating sporulation (Figure 2) [40, 41]. The action of the transcription factor Rme1, which is uniquely expressed in haploids, establishes the chromatin landscape regulating *IME1* expression. In haploid cells, *IME1* is strongly repressed by the binding of Rme1 to the *UCS* elements to prevent the aberrant activation of the sporulation program [42–44]. The mechanism by which Rme1 repressed *IME1* remained unknown for some time, however a breakthrough revealed that Rme1 promotes transcription of a long non-coding RNA (lncRNA). The transcription of this lncRNA, *IRT1*, from the upstream control sequences of *IME1* generates a repressive environment by interfering with activator recruitment to the *IME1* promoter and by the co-transcriptional recruitment of HMT and HDAC complexes (Figure 2) [45]. Specifically, *IRT1* transcription promotes recruitment of Set2 and Set3, each of which generate repressive environments through interactions of histone methyl marks and chromatin reader domains [46–48]. vanWerven [45] and colleagues propose that co-transcriptional recruitment of Set1 during *IRT1* induction deposits H3K4me2 at the 5' end of the *IME1* promoter, which stabilizes the Set3 complex and its HDAC activity [48], causing local repression. Likewise, co-transcriptional recruitment of Set2, which deposits H3K36me3, is postulated to promote association of the Rpd3(S) HDAC complex (as distinguished from Rpd3(L), a larger complex with distinct interactions) over the *IRT1* coding region [45–47]. This work therefore uncovered a key mechanism of the combinatorial role of histone modifications in the control of a critical cell fate regulator.

A second pathway reported to repress *IME1* in vegetatively-growing cells involves the Tup1/Ssn6 co-repressor complex [49, 50]. Cells lacking *TUP1* show decreased nucleosome occupancy of the *IME1* promoter, and increased recruitment of RNA pol II, consistent with constitutive expression of *IME1* [50]. However, the mechanism by which Tup1 associates with *IME1* is unclear, and whether it recruits other chromatin factors has not yet been investigated. Tup1 has been shown to play diverse roles at other nutrient-responsive promoters, such as driving the association of HDACs with *FLO1* [51, 52] and cooperating with the SWR1 complex for deposition of H2A.Z (Htz1) at the *GAL1* and *SUC2* promoters [53]. As a repressor of *IME1*, Tup1 has been linked to nutrient-sensing through PKA and TOR complex signaling networks [50], whereas Rme1 levels are responsive to mating type and ploidy [42, 54, 55]. These two DNA-binding proteins therefore represent different nodes of signal integration which promote the required chromatin state at the *IME1* promoter in response to both extrinsic and intrinsic signals.

## Histone modifications and meiotic progression

With the onset of the sporulation program, Ime1 activates a plethora of genes involved in meiotic DNA replication, chromosome remodeling and homologous recombination [5, 32]. A number of other chromatin-dependent mechanisms control the transcriptional dynamics of this cascade. Early and middle genes are also repressed by Ume6, a DNA binding protein that tethers Rpd3 and the chromatin remodeler Isw2 to promoter regions, resulting in histone deacetylation, chromatin compaction and occlusion of the TATA box [56, 57]. As Ime1 levels increase, Ume6 is targeted for degradation by the anaphase-promoting complex/cyclosome (APC/C) ubiquitin ligase activity, in a manner partially dependent on Ime1, and early gene transcription is initiated [58]. Intriguingly, Ume6 is targeted by the HAT Gcn5 for acetylation, and can be deacetylated by Rpd3 [59]. Lysine acetylation of Ume6 appears to promote its destruction, revealing that a non-histone target of Gcn5 may be a critical substrate determining its role in early gene expression, and that Rpd3 has an additional role in stabilizing Ume6 prior to the onset of the meiotic program [59]. Acetylation at additional lysines of Ume6 also appears to release it from its DNA binding sites, providing another mechanism for the rapid inactivation of Ume6 and ensuring proper timing of the transcriptional cascade [60]. These observations highlight the potential contribution of non-histone targets of chromatin-modifying enzymes to the regulation of reprogramming events.

Middle genes are repressed by the Sum1 transcriptional repressor, which binds DNA and tethers the HDAC Hst1 to middle gene promoters through an interaction with the adaptor protein Rfm1 [61–63]. Sum1-Rfm1-Hst1 and Ume6-mediated mechanisms have been reported to repress the middle sporulation gene *NDT80* [64], which encodes a master regulator transcription factor that activates a series of middle genes required for nuclear division and spore formation [63, 65]. Beyond this, we still have relatively little knowledge regarding the role of chromatin modifiers during middle, and also late-stage, gene expression. As an example, the methyltransferase Set1 has been implicated in repressing middle gene expression [66], however the mechanism by which it specifically represses this gene set has not been reported. While Set1's activity promotes meiotic progression, this is primarily thought to be due to the role of H3K4 methylation in recombination and double strand break repair [67, 68], whereas the function of Set1 in meiotic gene expression control is unclear.

In addition to the large-scale changes in gene expression that accompany the sporulation program, massive chromosomal rearrangements, which rely on critical genome integrity pathways, also occur during meiosis. While the role of chromatin in these processes has been reviewed elsewhere [69], genetic and proteomic screens focused on meiotic chromatin and sporulation have uncovered previously-unknown modifications. These approaches have revealed that phosphorylation at H4S1 is required for post-meiotic chromatin compaction and spore maturation [70], H3T11ph has been implicated in meiotic progression [71] and H4K44ac is associated with meiotic recombination [21]. These studies not only elucidate key mechanisms of meiotic chromosomal reorganization, but also underscore the notion that novel, context-dependent histone modifications may be discovered through analysis of specialized cell fates.



## Chromatin-based regulation of *FLO11*, initiator of filamentous growth

High expression levels of *FLO11*, a cell surface glycoprotein that mediates cell-cell and cell-surface adhesion, causes flocculation, haploid invasion, biofilm formation and diploid pseudohyphal, or filamentous, growth [6, 72, 73]. In *S. cerevisiae*, a family of related genes that encode adhesion proteins, *FLO1*, *FLO5*, *FLO9* and *FLO10*, are not expressed due to their proximity to telomeres and position-dependent gene silencing [74]. This gene family is similar to the *ALS* and *EPA* adhesin gene families in the human pathogens *Candida albicans* and *Candida glabrata*, respectively. In particular, the *EPA* genes of *C. glabrata* are also adjacent to telomeres and subject to metastable gene silencing, with activation of individual *EPA* genes occurring in different host environments and generating cell surface variation for the pathogen [75]. Understanding the regulatory mechanisms controlling expression of these gene families is therefore of high clinical relevance due to their role in promoting adherence to host cells, such as human epithelial cells, and biofilm formation on abiotic surfaces, including implanted medical devices [76].

Similar to *IME1*, *FLO11* expression is controlled by a very large, complex promoter that receives input from multiple nutrient-responsive signaling pathways, including the MAPK, SNF, TOR and RAS/cAMP-PKA pathways [6, 77]. Its expression levels are modulated through the coordinated activity of transcription factors, such as Flo8, which govern the contributions of histone-modifying enzymes, chromatin remodelers and lncRNAs to *FLO11* expression (Figure 3). While genetic screens have revealed potential involvement of the HAT Gcn5, the histone variant Htz1 and chromatin remodelers RSC and SWI/SNF in *FLO11* expression [78, 79], HDACs have the most well-documented roles in both repressing and activating *FLO11* expression. Epigenetic silencing of *FLO11* was first linked to the HDAC Hda1 [80], however other HDACs including Hos2 and Rpd3 have been implicated in its expression control. Interestingly, mutations of components of the Rpd3(L) complex (a larger Rpd3 complex than Rpd3(S) with unique interactions at chromatin) further repress *FLO11* [79], and combined deletion of Rpd3(L) and Hda1 demonstrates their partially overlapping functions in nucleosome depletion at the promoter and *FLO11* activation [81]. The paradoxical function of Rpd3(L) in *FLO11* activation was found to be due to the upstream expression of the lncRNA *ICR1*, in a manner dependent on the competing activity of two transcription factors [82, 83] (Figure 3). In the absence of the repressive chromatin environment generated by Rpd3(L), *ICR1* is transcribed and prevents expression of *FLO11* by occluding key *trans*-acting factors, such as TBP and the Flo8 transcriptional activator, from the promoter region [82, 83]. While these studies revealed a critical link between histone deacetylation and lncRNA-mediated repression of *FLO11*, there remain many open questions regarding the roles of other HDACs in this regulatory network. As one example, the Set3 complex, of which the HDAC Hos2 is a member, promotes filamentous growth in *Candida albicans* [84] and is required for lncRNA-mediated repression of *IME1*, yet its potential role in *FLO11* expression control is unclear.

Recent genetic studies have implicated other types of chromatin-modifying enzymes in regulating *FLO11* expression. The HAT Gcn5 has been reported to negatively regulate *ICR1* expression, thereby promoting *FLO11* transcriptional activation [85], most likely through regulation of a second ncRNA. Additionally, deletion of the demethylase *JHD2* or a



component of the CDK8 submodule of RNA pol II, *SSN8*, results in elevated H3K4me3 near the *FLO11* transcription start site (TSS) and constitutive activation of *FLO11* [86]. These data suggest a direct role for H3K4me3 in *FLO11* activation, although its potential contribution to lncRNA-dependent regulation of *FLO11* has not yet been investigated.

## Chromatin dynamics during programmed cell death

Programmed cell death (PCD) in metazoans has key developmental roles, such as promoting proper tissue architecture and the destruction of damaged cells, and aberrant regulation of PCD pathways is implicated in diseases including cancer and neurodegenerative disorders [7, 87]. Based on the lack of some of the cell death machinery, it was not clear that *S. cerevisiae* exhibited PCD until apoptosis-related phenotypes were observed in a *CDC48* mutant. Observed phenotypes included DNA fragmentation, chromatin compaction, increased reactive oxygen species and phosphatidylserine externalization [88]. It is now appreciated that budding yeast undergo multiple types of cell death, including caspase-dependent and caspase-independent intrinsic apoptosis, autophagic cell death and regulated necrosis [89]. These pathways are activated in response to a multitude of intrinsic and extrinsic signals, including oxidative and osmotic stress, acetic acid and aging, among others [7]. While PCD is clearly not required for tissue architecture in unicellular yeasts, it may act to eliminate old, damaged or nonreplicative cells, releasing nutrients to younger, replicative cells and promoting their survival [7, 89]. Interestingly, many filamentous fungi have more similar PCD mechanisms to animal cells than budding yeast, and PCD has frequently been observed to be important for pathogenic life cycles and the formation of multicellular-like structures in these organisms [90].

Histone modifications have the potential to be important for multiple components of PCD, including gene expression associated with stress responses, transcription of apoptotic regulators and the processes of chromatin condensation and DNA fragmentation. The Ste20 kinase phosphorylates H2BS10 in apoptotic chromatin in yeast [91], which is analogous to human H2BS14ph, catalyzed by the caspase-activated kinase Mst1 [92]. It was also found that deacetylation at H2BK11 by the HDAC Hos3 is a prerequisite for Ste20-mediated phosphorylation at H2BS10 [93]. This crosstalk between histone modifications appears to promote peroxide-mediated apoptosis, with H2BS10ph proposed to have a role in chromatin condensation [91, 93]. However, this mechanism has been debated [94], and it is unclear whether other substrates of Mst1 (or Ste20) may promote PCD-associated chromatin compaction [94, 95].

In a similar example of modification cross-talk, H2BK123Ub by Bre1 was reported to be an anti-apoptotic mark, with loss of the mark suspected to promote DNA damage or transcriptional deregulation of apoptotic factors [96]. H2BK123Ub precedes Set1-dependent H3K4 methylation, as well as H3K79 methylation catalyzed by Dot1 [97]. Subsequent investigation revealed that *BRE1* mutants lacking H2BK123Ub undergo PCD due to the loss of H3K4 methylation by Set1 [98]. The loss of Set1 promotes PCD in a manner dependent on both Dot1/H3K79me and the checkpoint kinase Rad9. Increased PCD in *SET1* mutants can be rescued by simultaneous inactivation of Nuc1, the yeast ortholog of endonuclease G associated with DNA fragmentation [98]. Although further experiments are required, these

data suggest that Set1 and H3K4 methylation may play a global role in promoting proper chromatin structure to prevent PCD-mediated condensation and fragmentation of DNA.

The studies conducted to date on histone modifications linked to yeast PCD have revealed that both program-specific (H2BS10ph) and general (H2BK123Ub, H3K4me) modifications make contributions to this developmental pathway and that cross-talk between modifications may serve to integrate multiple inputs from the signaling networks activating PCD. This highlights the need to comprehensively investigate the full complement of histone modifications participating in PCD. Additionally, acetylation of the autophagic protein Atg3 by the HAT Esa1 has been linked to the control of autophagy [99], indicating that there may be more diverse roles for typical chromatin-modifying enzymes that remain to be discovered in cell death pathways.

## Concluding Remarks

We have reviewed a number of significant advances in our understanding of chromatin modifications critical to yeast cell fates. In the representative cases cited here, key themes emerge and areas for further exploration are highlighted. First, while chromatin-modifying enzymes such as Rpd3, Gcn5, Set1 and Set2 have broad functions throughout the genome, they also have critical context-dependent roles, responding to specific cues and acting within distinct genomic regions to advance cell fate programs. It is also evident that chromatin modifiers integrate signals from multiple sources, and their activity is governed by complex networks that signal to transcription factors, lncRNAs, and likely other factors, to promote the appropriate changes in the local chromatin landscape.

While substantial progress has been made in understanding the functions of diverse histone modifications in meiosis and sporulation, pseudohyphal differentiation and programmed cell death pathways in yeast, it is clear that there are still many open questions. Continued investigation into the new modifications revealed by high-resolution proteomics may elucidate state-specific roles for these marks, particularly for those that are lowly abundant or enriched in certain environmental conditions. Moreover, further understanding of the role of non-histone PTMs catalyzed by enzymes classified as chromatin modifiers may uncover additional regulatory nodes critical to these cell fate programs and diversify the biological functions of these enzymes. And finally, given the role of nutritional cues in these cell fate signaling networks, it will be critical to delineate how metabolic changes within the cell affect chromatin-modifying enzymes. There is increasing evidence that metabolic changes can signal directly to these enzymes, potentially regulating their activity, localization or co-factor availability to drive cell fate changes [100, 101]. The high conservation of chromatin-modifying pathways and the breadth of knowledge regarding the genome and epigenome suggest that continued investigation of histone modifications in yeast will yield key insights into cell fate decisions in diverse organisms.

## Acknowledgments

The authors acknowledge members of the Green lab for fruitful discussions and comments on the manuscript. This work was supported in part by NIH R03AG052018 and a UMBC START award to EMG.

## Abbreviations

<b>lncRNA</b>	long non-coding ribonucleic acid
<b>DNA</b>	deoxyribonucleic acid
<b>PTM</b>	post-translational modification
<b>HAT</b>	histone acetyltransferase
<b>HDAC</b>	histone deacetylase
<b>HMT</b>	histone methyltransferase
<b>HDM</b>	histone demethylase
<b>UbT</b>	ubiquitin-protein transferase
<b>DUB</b>	deubiquinase
<b>me</b>	methylation
<b>ac</b>	acetylation
<b>ph</b>	phosphorylation
<b>MAPK</b>	mitogen activated protein kinase
<b>TOR</b>	target of rapamycin
<b>PKA</b>	protein kinase A
<b>UCS</b>	upstream control sequence
<b>APC/C</b>	anaphase-promoting complex/cyclosome
<b>PCD</b>	programmed cell death
<b>kb</b>	kilobases

## References

1. Suganuma T, Workman JL. Signals and combinatorial functions of histone modifications. *Annu Rev Biochem.* 2011; 80:473–99. [PubMed: 21529160]
2. Badeaux AI, Shi Y. Emerging roles for chromatin as a signal integration and storage platform. *Nat Rev Mol Cell Biol.* 2013; 14:211–24.
3. Bannister AJ, Kouzarides T. Regulation of chromatin by histone modifications. *Cell Res.* 2011; 21:381–95. [PubMed: 21321607]
4. Allis CD, Jenuwein T. The molecular hallmarks of epigenetic control. *Nat Rev Genet.* 2016; 17:487–500. [PubMed: 27346641]
5. Neiman AM. Sporulation in the budding yeast *Saccharomyces cerevisiae*. *Genetics.* 2011; 189:737–65. [PubMed: 22084423]
6. Cullen PJ, Sprague GF Jr. The regulation of filamentous growth in yeast. *Genetics.* 2012; 190:23–49. [PubMed: 22219507]

7. Strich R. Programmed Cell Death Initiation and Execution in Budding Yeast. *Genetics*. 2015; 200:1003–14. [PubMed: 26272996]
8. Meneghini MD, Wu M, Madhani HD. Conserved histone variant H2A.Z protects euchromatin from the ectopic spread of silent heterochromatin. *Cell*. 2003; 112:725–36. [PubMed: 12628191]
9. Zhang H, Roberts DN, Cairns BR. Genome-wide dynamics of Htz1, a histone H2A variant that poises repressed/basal promoters for activation through histone loss. *Cell*. 2005; 123:219–31. [PubMed: 16239141]
10. Meluh PB, Yang P, Glowczewski L, Koshland D, Smith MM. Cse4p is a component of the core centromere of *Saccharomyces cerevisiae*. *Cell*. 1998; 94:607–13. [PubMed: 9741625]
11. Bonasio R, Tu S, Reinberg D. Molecular signals of epigenetic states. *Science*. 2010; 330:612–6. [PubMed: 21030644]
12. Tee WW, Reinberg D. Chromatin features and the epigenetic regulation of pluripotency states in ESCs. *Development*. 2014; 141:2376–90. [PubMed: 24917497]
13. Taverna SD, Li H, Ruthenburg AJ, Allis CD, Patel DJ. How chromatin-binding modules interpret histone modifications: lessons from professional pocket pickers. *Nat Struct Mol Biol*. 2007; 14:1025–40. [PubMed: 17984965]
14. Gardner KE, Allis CD, Strahl BD. Operating on chromatin, a colorful language where context matters. *J Mol Biol*. 2011; 409:36–46. [PubMed: 21272588]
15. Musselman CA, Lalonde ME, Côté J, Kutateladze TG. Perceiving the epigenetic landscape through histone readers. *Nat Struct Mol Biol*. 2012; 19:1218–27. [PubMed: 23211769]
16. Garcia BA, Hake SB, Diaz RL, Kauer M, Morris SA, Recht J, et al. Organismal differences in post-translational modifications in histones H3 and H4. *J Biol Chem*. 2007; 282:7641–55. [PubMed: 17194708]
17. Tan M, Luo H, Lee S, Jin F, Yang JS, Montellier E, et al. Identification of 67 histone marks and histone lysine crotonylation as a new type of histone modification. *Cell*. 2011; 146:1016–28. [PubMed: 21925322]
18. Arnaudo AM, Garcia BA. Proteomic characterization of novel histone post-translational modifications. *Epigenetics Chromatin*. 2013; 6:24. [PubMed: 23916056]
19. Xie Z, Dai J, Dai L, Tan M, Cheng Z, Wu Y, et al. Lysine succinylation and lysine malonylation in histones. *Mol Cell Proteomics*. 2012; 11:100–7. [PubMed: 22389435]
20. Zhao Y, Garcia BA. Comprehensive Catalog of Currently Documented Histone Modifications. *Cold Spring Harb Perspect Biol*. 2015; 7:a025064. [PubMed: 26330523]
21. Hu J, Donahue G, Dorsey J, Govin J, Yuan Z, Garcia BA, et al. H4K44 Acetylation Facilitates Chromatin Accessibility during Meiosis. *Cell Rep*. 2015; 13:1772–80. [PubMed: 26628362]
22. Green EM, Mas G, Young NL, Garcia BA, Gozani O. Methylation of H4 lysines 5, 8 and 12 by yeast Set5 calibrates chromatin stress responses. *Nat Struct Mol Biol*. 2012; 19:361–3. [PubMed: 22343720]
23. Martín GM, King DA, Green EM, Garcia-Nieto PE, Alexander R, Collins SR, et al. Set5 and Set1 cooperate to repress gene expression at telomeres and retrotransposons. *Epigenetics*. 2014;9.
24. Zhang X, Huang Y, Shi X. Emerging roles of lysine methylation on non-histone proteins. *Cell Mol Life Sci*. 2015; 72:4257–72. [PubMed: 26227335]
25. Biggar KK, Li SS. Non-histone protein methylation as a regulator of cellular signalling and function. *Nat Rev Mol Cell Biol*. 2015; 16:5–17. [PubMed: 25491103]
26. Moore KE, Gozani O. An unexpected journey: lysine methylation across the proteome. *Biochim Biophys Acta*. 2014; 1839:1395–403. [PubMed: 24561874]
27. Moore KE, Carlson SM, Camp ND, Cheung P, James RG, Chua KF, et al. A general molecular affinity strategy for global detection and proteomic analysis of lysine methylation. *Mol Cell*. 2013; 50:444–56. [PubMed: 23583077]
28. Mazur PK, Reynoird N, Khatri P, Jansen PW, Wilkinson AW, Liu S, et al. SMYD3 links lysine methylation of MAP3K2 to Ras-driven cancer. *Nature*. 2014
29. Zhang K, Lin W, Latham JA, Riefler GM, Schumacher JM, Chan C, et al. The Set1 methyltransferase opposes Ipl1 aurora kinase functions in chromosome segregation. *Cell*. 2005; 122:723–34. [PubMed: 16143104]

30. Lin YY, Lu JY, Zhang J, Walter W, Dang W, Wan J, et al. Protein acetylation microarray reveals that NuA4 controls key metabolic target regulating gluconeogenesis. *Cell*. 2009; 136:1073–84. [PubMed: 19303850]
31. Tessarz P, Santos-Rosa H, Robson SC, Sylvestersen KB, Nelson CJ, Nielsen ML, et al. Glutamine methylation in histone H2A is an RNA-polymerase-I-dedicated modification. *Nature*. 2014; 505:564–8. [PubMed: 24352239]
32. Honigberg SM, Purnapatre K. Signal pathway integration in the switch from the mitotic cell cycle to meiosis in yeast. *J Cell Sci*. 2003; 116:2137–47. [PubMed: 12730290]
33. Chu S, DeRisi J, Eisen M, Mulholland J, Botstein D, Brown PO, et al. The transcriptional program of sporulation in budding yeast. *Science*. 1998; 282:699–705. [PubMed: 9784122]
34. Primig M, Williams RM, Winzeler EA, Tevzadze GG, Conway AR, Hwang SY, et al. The core meiotic transcriptome in budding yeasts. *Nat Genet*. 2000; 26:415–23. [PubMed: 11101837]
35. Shively CA, Eckwahl MJ, Dobry CJ, Mellacheruvu D, Nesvizhskii A, Kumar A. Genetic networks inducing invasive growth in *Saccharomyces cerevisiae* identified through systematic genome-wide overexpression. *Genetics*. 2013; 193:1297–310. [PubMed: 23410832]
36. Mitchell AP. Control of meiotic gene expression in *Saccharomyces cerevisiae*. *Microbiol Rev*. 1994; 58:56–70. [PubMed: 8177171]
37. Smith HE, Driscoll SE, Sia RA, Yuan HE, Mitchell AP. Genetic evidence for transcriptional activation by the yeast *IME1* gene product. *Genetics*. 1993; 133:775–84. [PubMed: 8462841]
38. Kassir Y, Granot D, Simchen G. *IME1*, a positive regulator gene of meiosis in *S. cerevisiae*. *Cell*. 1988; 52:853–62. [PubMed: 3280136]
39. Mandel S, Robzyk K, Kassir Y. *IME1* gene encodes a transcription factor which is required to induce meiosis in *Saccharomyces cerevisiae*. *Dev Genet*. 1994; 15:139–47. [PubMed: 8205723]
40. Granot D, Margolskee JP, Simchen G. A long region upstream of the *IME1* gene regulates meiosis in yeast. *Mol Gen Genet*. 1989; 218:308–14. [PubMed: 2674657]
41. Sagee S, Sherman A, Shenhar G, Robzyk K, Ben-Doy N, Simchen G, et al. Multiple and distinct activation and repression sequences mediate the regulated transcription of *IME1*, a transcriptional activator of meiosis-specific genes in *Saccharomyces cerevisiae*. *Mol Cell Biol*. 1998; 18:1985–95. [PubMed: 9528770]
42. Covitz PA, Mitchell AP. Repression by the yeast meiotic inhibitor *RME1*. *Genes Dev*. 1993; 7:1598–608. [PubMed: 8339935]
43. Shimizu M, Hara M, Murase A, Shindo H, Mitchell AP. Dissection of the DNA binding domain of yeast Zn-finger protein *Rme1p*, a repressor of meiotic activator *IME1*. *Nucleic Acids Symp Ser*. 1997:175–6. [PubMed: 9586056]
44. Shimizu M, Li W, Covitz PA, Hara M, Shindo H, Mitchell AP. Genomic footprinting of the yeast zinc finger protein *Rme1p* and its roles in repression of the meiotic activator *IME1*. *Nucleic Acids Res*. 1998; 26:2329–36. [PubMed: 9580682]
45. van Werven FJ, Neuert G, Hendrick N, Lardenois A, Buratowski S, van Oudenaarden A, et al. Transcription of two long noncoding RNAs mediates mating-type control of gametogenesis in budding yeast. *Cell*. 2012; 150:1170–81. [PubMed: 22959267]
46. Keogh MC, Kurdistani SK, Morris SA, Ahn SH, Podolny V, Collins SR, et al. Cotranscriptional set2 methylation of histone H3 lysine 36 recruits a repressive Rpd3 complex. *Cell*. 2005; 123:593–605. [PubMed: 16286008]
47. Carrozza MJ, Li B, Florens L, Suganuma T, Swanson SK, Lee KK, et al. Histone H3 methylation by Set2 directs deacetylation of coding regions by Rpd3S to suppress spurious intragenic transcription. *Cell*. 2005; 123:581–92. [PubMed: 16286007]
48. Kim T, Buratowski S. Dimethylation of H3K4 by Set1 recruits the Set3 histone deacetylase complex to 5' transcribed regions. *Cell*. 2009; 137:259–72. [PubMed: 19379692]
49. Mizuno T, Nakazawa N, Remgsamrarn P, Kunoh T, Oshima Y, Harashima S. The *Tup1-Ssn6* general repressor is involved in repression of *IME1* encoding a transcriptional activator of meiosis in *Saccharomyces cerevisiae*. *Curr Genet*. 1998; 33:239–47. [PubMed: 9560430]
50. Weidberg H, Moretto F, Spedale G, Amon A, van Werven FJ. Nutrient Control of Yeast Gametogenesis Is Mediated by TORC1, PKA and Energy Availability. *PLoS Genet*. 2016; 12:e1006075. [PubMed: 27272508]

51. Davie JK, Edmondson DG, Coco CB, Dent SY. Tup1-Ssn6 interacts with multiple class I histone deacetylases in vivo. *J Biol Chem.* 2003; 278:50158–62. [PubMed: 14525981]
52. Fleming AB, Beggs S, Church M, Tsukihashi Y, Pennings S. The yeast Cyc8-Tup1 complex cooperates with Hda1p and Rpd3p histone deacetylases to robustly repress transcription of the subtelomeric FLO1 gene. *Biochim Biophys Acta.* 2014; 1839:1242–55. [PubMed: 25106892]
53. Gligoris T, Thireos G, Tzamarias D. The Tup1 corepressor directs Htz1 deposition at a specific promoter nucleosome marking the GAL1 gene for rapid activation. *Mol Cell Biol.* 2007; 27:4198–205. [PubMed: 17387147]
54. Mitchell AP, Herskowitz I. Activation of meiosis and sporulation by repression of the RME1 product in yeast. *Nature.* 1986; 319:738–42. [PubMed: 3513021]
55. Covitz PA, Herskowitz I, Mitchell AP. The yeast RME1 gene encodes a putative zinc finger protein that is directly repressed by a1-alpha 2. *Genes Dev.* 1991; 5:1982–9. [PubMed: 1936989]
56. Goldmark JP, Fazzio TG, Estep PW, Church GM, Tsukiyama T. The Isw2 chromatin remodeling complex represses early meiotic genes upon recruitment by Ume6p. *Cell.* 2000; 103:423–33. [PubMed: 11081629]
57. Kadosh D, Struhl K. Repression by Ume6 involves recruitment of a complex containing Sin3 corepressor and Rpd3 histone deacetylase to target promoters. *Cell.* 1997; 89:365–71. [PubMed: 9150136]
58. Mallory MJ, Cooper KF, Strich R. Meiosis-specific destruction of the Ume6p repressor by the Cdc20-directed APC/C. *Mol Cell.* 2007; 27:951–61. [PubMed: 17889668]
59. Mallory MJ, Law MJ, Sterner DE, Berger SL, Strich R. Gcn5p-dependent acetylation induces degradation of the meiotic transcriptional repressor Ume6p. *Mol Biol Cell.* 2012; 23:1609–17. [PubMed: 22438583]
60. Law MJ, Mallory MJ, Dunbrack RL, Strich R. Acetylation of the transcriptional repressor Ume6p allows efficient promoter release and timely induction of the meiotic transient transcription program in yeast. *Mol Cell Biol.* 2014; 34:631–42. [PubMed: 24298021]
61. Xie J, Pierce M, Gailus-Durner V, Wagner M, Winter E, Vershon AK. Sum1 and Hst1 repress middle sporulation-specific gene expression during mitosis in *Saccharomyces cerevisiae*. *EMBO J.* 1999; 18:6448–54. [PubMed: 10562556]
62. McCord R, Pierce M, Xie J, Wonkatal S, Mickel C, Vershon AK. Rfm1, a novel tethering factor required to recruit the Hst1 histone deacetylase for repression of middle sporulation genes. *Mol Cell Biol.* 2003; 23:2009–16. [PubMed: 12612074]
63. Winter E. The Sum1/Ndt80 transcriptional switch and commitment to meiosis in *Saccharomyces cerevisiae*. *Microbiol Mol Biol Rev.* 2012; 76:1–15. [PubMed: 22390969]
64. Pierce M, Benjamin KR, Montano SP, Georgiadis MM, Winter E, Vershon AK. Sum1 and Ndt80 proteins compete for binding to middle sporulation element sequences that control meiotic gene expression. *Mol Cell Biol.* 2003; 23:4814–25. [PubMed: 12832469]
65. Chu S, Herskowitz I. Gametogenesis in yeast is regulated by a transcriptional cascade dependent on Ndt80. *Mol Cell.* 1998; 1:685–96. [PubMed: 9660952]
66. Sollier J, Lin W, Soustelle C, Suhre K, Nicolas A, Géli V, et al. Set1 is required for meiotic S-phase onset, double-strand break formation and middle gene expression. *EMBO J.* 2004; 23:1957–67. [PubMed: 15071505]
67. Borde V, Robine N, Lin W, Bonfils S, Géli V, Nicolas A. Histone H3 lysine 4 trimethylation marks meiotic recombination initiation sites. *EMBO J.* 2009; 28:99–111. [PubMed: 19078966]
68. Sommermeyer V, Béneut C, Chaplais E, Serrentino ME, Borde V. Spp1, a member of the Set1 Complex, promotes meiotic DSB formation in promoters by tethering histone H3K4 methylation sites to chromosome axes. *Mol Cell.* 2013; 49:43–54. [PubMed: 23246437]
69. Govin J, Berger SL. Genome reprogramming during sporulation. *Int J Dev Biol.* 2009; 53:425–32. [PubMed: 19412896]
70. Krishnamoorthy T, Chen X, Govin J, Cheung WL, Dorsey J, Schindler K, et al. Phosphorylation of histone H4 Ser1 regulates sporulation in yeast and is conserved in fly and mouse spermatogenesis. *Genes Dev.* 2006; 20:2580–92. [PubMed: 16980586]



71. Govin J, Dorsey J, Gaucher J, Rousseaux S, Khochbin S, Berger SL. Systematic screen reveals new functional dynamics of histones H3 and H4 during gametogenesis. *Genes Dev.* 2010; 24:1772–86. [PubMed: 20713519]
72. Lo WS, Dranginis AM. The cell surface flocculin Flo11 is required for pseudohyphae formation and invasion by *Saccharomyces cerevisiae*. *Mol Biol Cell.* 1998; 9:161–71. [PubMed: 9436998]
73. Reynolds TB, Fink GR. Bakers' yeast, a model for fungal biofilm formation. *Science.* 2001; 291:878–81. [PubMed: 11157168]
74. Guo B, Styles CA, Feng Q, Fink GR. A *Saccharomyces* gene family involved in invasive growth, cell-cell adhesion, and mating. *Proc Natl Acad Sci U S A.* 2000; 97:12158–63. [PubMed: 11027318]
75. De Las Peñas A, Juárez-Cepeda J, López-Fuentes E, Briones-Martín-Del-Campo M, Gutiérrez-Escobedo G, Castaño I. Local and regional chromatin silencing in *Candida glabrata*: consequences for adhesion and the response to stress. *FEMS Yeast Res.* 2015:15.
76. Nobile CJ, Johnson AD. *Candida albicans* Biofilms and Human Disease. *Annu Rev Microbiol.* 2015; 69:71–92. [PubMed: 26488273]
77. Brückner S, Mösch HU. Choosing the right lifestyle: adhesion and development in *Saccharomyces cerevisiae*. *FEMS Microbiol Rev.* 2012; 36:25–58. [PubMed: 21521246]
78. Fischer C, Valerius O, Rupprecht H, Dumkow M, Krappmann S, Braus GH. Posttranscriptional regulation of FLO11 upon amino acid starvation in *Saccharomyces cerevisiae*. *FEMS Yeast Res.* 2008; 8:225–36. [PubMed: 17999676]
79. Barrales RR, Jimenez J, Ibeas JI. Identification of novel activation mechanisms for FLO11 regulation in *Saccharomyces cerevisiae*. *Genetics.* 2008; 178:145–56. [PubMed: 18202364]
80. Halme A, Bumgarner S, Styles C, Fink GR. Genetic and epigenetic regulation of the FLO gene family generates cell-surface variation in yeast. *Cell.* 2004; 116:405–15. [PubMed: 15016375]
81. Barrales RR, Korber P, Jimenez J, Ibeas JI. Chromatin modulation at the FLO11 promoter of *Saccharomyces cerevisiae* by HDAC and Swi/Snf complexes. *Genetics.* 2012; 191:791–803. [PubMed: 22542969]
82. Bumgarner SL, Dowell RD, Grisafi P, Gifford DK, Fink GR. Toggle involving cis-interfering noncoding RNAs controls variegated gene expression in yeast. *Proc Natl Acad Sci U S A.* 2009; 106:18321–6. [PubMed: 19805129]
83. Bumgarner SL, Neuert G, Voight BF, Symbor-Nagrabska A, Grisafi P, van Oudenaarden A, et al. Single-cell analysis reveals that noncoding RNAs contribute to clonal heterogeneity by modulating transcription factor recruitment. *Mol Cell.* 2012; 45:470–82. [PubMed: 22264825]
84. Hnisz D, Majer O, Frohner IE, Komnenovic V, Kuchler K. The Set3/Hos2 histone deacetylase complex attenuates cAMP/PKA signaling to regulate morphogenesis and virulence of *Candida albicans*. *PLoS Pathog.* 2010; 6:e1000889. [PubMed: 20485517]
85. Wang LC, Montalvo-Munoz F, Tsai YC, Liang CY, Chang CC, Lo WS. The Histone Acetyltransferase Gcn5 Regulates ncRNA-ICR1 and FLO11 Expression during Pseudohyphal Development in *Saccharomyces cerevisiae*. *Biomed Res Int.* 2015; 2015:284692. [PubMed: 25922832]
86. Law MJ, Ciccaglione K. Fine-tuning of histone H3 Lys4 methylation during pseudohyphal differentiation by the CDK submodule of RNA polymerase II. *Genetics.* 2015; 199:435–53. [PubMed: 25467068]
87. Kiraz Y, Adan A, Kartal Yandim M, Baran Y. Major apoptotic mechanisms and genes involved in apoptosis. *Tumour Biol.* 2016
88. Madeo F, Fröhlich E, Fröhlich KU. A yeast mutant showing diagnostic markers of early and late apoptosis. *J Cell Biol.* 1997; 139:729–34. [PubMed: 9348289]
89. Lin SJ, Austriaco N. Aging and cell death in the other yeasts, *Schizosaccharomyces pombe* and *Candida albicans*. *FEMS Yeast Res.* 2014; 14:119–35. [PubMed: 24205865]
90. Shlezinger N, Goldfinger N, Sharon A. Apoptotic-like programmed cell death in fungi: the benefits in filamentous species. *Front Oncol.* 2012; 2:97. [PubMed: 22891165]
91. Ahn SH, Cheung WL, Hsu JY, Diaz RL, Smith MM, Allis CD. Sterile 20 kinase phosphorylates histone H2B at serine 10 during hydrogen peroxide-induced apoptosis in *S. cerevisiae*. *Cell.* 2005; 120:25–36. [PubMed: 15652479]




92. Cheung WL, Ajiro K, Samejima K, Kloc M, Cheung P, Mizzen CA, et al. Apoptotic phosphorylation of histone H2B is mediated by mammalian sterile twenty kinase. *Cell*. 2003; 113:507–17. [PubMed: 12757711]
93. Ahn SH, Diaz RL, Grunstein M, Allis CD. Histone H2B deacetylation at lysine 11 is required for yeast apoptosis induced by phosphorylation of H2B at serine 10. *Mol Cell*. 2006; 24:211–20. [PubMed: 17052455]
94. Ura S, Nishina H, Gotoh Y, Katada T. Activation of the c-Jun N-terminal kinase pathway by MST1 is essential and sufficient for the induction of chromatin condensation during apoptosis. *Mol Cell Biol*. 2007; 27:5514–22. [PubMed: 17548476]
95. Wen W, Zhu F, Zhang J, Keum YS, Zykova T, Yao K, et al. MST1 promotes apoptosis through phosphorylation of histone H2AX. *J Biol Chem*. 2010; 285:39108–16. [PubMed: 20921231]
96. Walter D, Matter A, Fahrenkrog B. Bre1p-mediated histone H2B ubiquitylation regulates apoptosis in *Saccharomyces cerevisiae*. *J Cell Sci*. 2010; 123:1931–9. [PubMed: 20460436]
97. Lee JS, Shukla A, Schneider J, Swanson SK, Washburn MP, Florens L, et al. Histone crosstalk between H2B monoubiquitination and H3 methylation mediated by COMPASS. *Cell*. 2007; 131:1084–96. [PubMed: 18083099]
98. Walter D, Matter A, Fahrenkrog B. Loss of histone H3 methylation at lysine 4 triggers apoptosis in *Saccharomyces cerevisiae*. *PLoS Genet*. 2014; 10:e1004095. [PubMed: 24497836]
99. Yi C, Ma M, Ran L, Zheng J, Tong J, Zhu J, et al. Function and molecular mechanism of acetylation in autophagy regulation. *Science*. 2012; 336:474–7. [PubMed: 22539722]
100. Li S, Swanson SK, Gogol M, Florens L, Washburn MP, Workman JL, et al. Serine and SAM Responsive Complex SESAME Regulates Histone Modification Crosstalk by Sensing Cellular Metabolism. *Mol Cell*. 2015; 60:408–21. [PubMed: 26527276]
101. Etchegaray JP, Mostoslavsky R. Interplay between Metabolism and Epigenetics: A Nuclear Adaptation to Environmental Changes. *Mol Cell*. 2016; 62:695–711. [PubMed: 27259202]
102. Briggs SD, Bryk M, Strahl BD, Cheung WL, Davie JK, Dent SY, et al. Histone H3 lysine 4 methylation is mediated by Set1 and required for cell growth and rDNA silencing in *Saccharomyces cerevisiae*. *Genes Dev*. 2001; 15:3286–95. [PubMed: 11751634]
103. Roguev A, Schaft D, Shevchenko A, Pijnappel WW, Wilm M, Aasland R, et al. The *Saccharomyces cerevisiae* Set1 complex includes an Ash2 homologue and methylates histone 3 lysine 4. *EMBO J*. 2001; 20:7137–48. [PubMed: 11742990]
104. Nagy PL, Griesenbeck J, Kornberg RD, Cleary ML. A trithorax-group complex purified from *Saccharomyces cerevisiae* is required for methylation of histone H3. *Proc Natl Acad Sci U S A*. 2002; 99:90–4. [PubMed: 11752412]
105. Krogan NJ, Dover J, Khorrami S, Greenblatt JF, Schneider J, Johnston M, et al. COMPASS, a histone H3 (Lysine 4) methyltransferase required for telomeric silencing of gene expression. *J Biol Chem*. 2002; 277:10753–5. [PubMed: 11805083]
106. Seward DJ, Cubberley G, Kim S, Schonewald M, Zhang L, Tripet B, et al. Demethylation of trimethylated histone H3 Lys4 in vivo by JARID1 JmjC proteins. *Nat Struct Mol Biol*. 2007; 14:240–2. [PubMed: 17310255]
107. Liang G, Klose RJ, Gardner KE, Zhang Y. Yeast Jhd2p is a histone H3 Lys4 trimethyl demethylase. *Nat Struct Mol Biol*. 2007; 14:243–5. [PubMed: 17310254]
108. Guillemette B, Drogaris P, Lin HH, Armstrong H, Hiragami-Hamada K, Imhof A, et al. H3 lysine 4 is acetylated at active gene promoters and is regulated by H3 lysine 4 methylation. *PLoS Genet*. 2011; 7:e1001354. [PubMed: 21483810]
109. Grant PA, Eberharter A, John S, Cook RG, Turner BM, Workman JL. Expanded lysine acetylation specificity of Gcn5 in native complexes. *J Biol Chem*. 1999; 274:5895–900. [PubMed: 10026213]
110. Kadosh D, Struhl K. Targeted recruitment of the Sin3-Rpd3 histone deacetylase complex generates a highly localized domain of repressed chromatin in vivo. *Mol Cell Biol*. 1998; 18:5121–7. [PubMed: 9710596]
111. Suka N, Suka Y, Carmen AA, Wu J, Grunstein M. Highly specific antibodies determine histone acetylation site usage in yeast heterochromatin and euchromatin. *Mol Cell*. 2001; 8:473–9. [PubMed: 11545749]

112. Fillingham J, Recht J, Silva AC, Suter B, Emili A, Stagljar I, et al. Chaperone control of the activity and specificity of the histone H3 acetyltransferase Rtt109. *Mol Cell Biol.* 2008; 28:4342–53. [PubMed: 18458063]
113. Lo WS, Duggan L, Emre NC, Belotserkovskya R, Lane WS, Shiekhattar R, et al. Snf1--a histone kinase that works in concert with the histone acetyltransferase Gcn5 to regulate transcription. *Science.* 2001; 293:1142–6. [PubMed: 11498592]
114. Hsu JY, Sun ZW, Li X, Reuben M, Tatchell K, Bishop DK, et al. Mitotic phosphorylation of histone H3 is governed by Ipl1/aurora kinase and Glc7/PP1 phosphatase in budding yeast and nematodes. *Cell.* 2000; 102:279–91. [PubMed: 10975519]
115. Kuo MH, Brownell JE, Sobel RE, Ranalli TA, Cook RG, Edmondson DG, et al. Transcription-linked acetylation by Gcn5p of histones H3 and H4 at specific lysines. *Nature.* 1996; 383:269–72. [PubMed: 8805705]
116. Strahl BD, Grant PA, Briggs SD, Sun ZW, Bone JR, Caldwell JA, et al. Set2 is a nucleosomal histone H3-selective methyltransferase that mediates transcriptional repression. *Mol Cell Biol.* 2002; 22:1298–306. [PubMed: 11839797]
117. Tu S, Bulloch EM, Yang L, Ren C, Huang WC, Hsu PH, et al. Identification of histone demethylases in *Saccharomyces cerevisiae*. *J Biol Chem.* 2007; 282:14262–71. [PubMed: 17369256]
118. Kim T, Buratowski S. Two *Saccharomyces cerevisiae* JmjC domain proteins demethylate histone H3 Lys36 in transcribed regions to promote elongation. *J Biol Chem.* 2007; 282:20827–35. [PubMed: 17525156]
119. Morris SA, Rao B, Garcia BA, Hake SB, Diaz RL, Shabanowitz J, et al. Identification of histone H3 lysine 36 acetylation as a highly conserved histone modification. *J Biol Chem.* 2007; 282:7632–40. [PubMed: 17189264]
120. Baker SP, Phillips J, Anderson S, Qiu Q, Shabanowitz J, Smith MM, et al. Histone H3 Thr 45 phosphorylation is a replication-associated post-translational modification in *S. cerevisiae*. *Nat Cell Biol.* 2010; 12:294–8. [PubMed: 20139971]
121. Han J, Zhou H, Horazdovsky B, Zhang K, Xu RM, Zhang Z. Rtt109 acetylates histone H3 lysine 56 and functions in DNA replication. *Science.* 2007; 315:653–5. [PubMed: 17272723]
122. Celic I, Masumoto H, Griffith WP, Meluh P, Cotter RJ, Boeke JD, et al. The sirtuins hst3 and Hst4p preserve genome integrity by controlling histone h3 lysine 56 deacetylation. *Curr Biol.* 2006; 16:1280–9. [PubMed: 16815704]
123. Tsubota T, Berndsen CE, Erkmann JA, Smith CL, Yang L, Freitas MA, et al. Histone H3-K56 acetylation is catalyzed by histone chaperone-dependent complexes. *Mol Cell.* 2007; 25:703–12. [PubMed: 17320445]
124. Yang B, Miller A, Kirchmaier AL. HST3/HST4-dependent deacetylation of lysine 56 of histone H3 in silent chromatin. *Mol Biol Cell.* 2008; 19:4993–5005. [PubMed: 18799617]
125. van Leeuwen F, Gafken PR, Gottschling DE. Dot1p modulates silencing in yeast by methylation of the nucleosome core. *Cell.* 2002; 109:745–56. [PubMed: 12086673]
126. Feng Q, Wang H, Ng HH, Erdjument-Bromage H, Tempst P, Struhl K, et al. Methylation of H3-lysine 79 is mediated by a new family of HMTases without a SET domain. *Curr Biol.* 2002; 12:1052–8. [PubMed: 12123582]
127. Lacoste N, Utley RT, Hunter JM, Poirier GG, Côte J. Disruptor of telomeric silencing-1 is a chromatin-specific histone H3 methyltransferase. *J Biol Chem.* 2002; 277:30421–4. [PubMed: 12097318]
128. Ng HH, Feng Q, Wang H, Erdjument-Bromage H, Tempst P, Zhang Y, et al. Lysine methylation within the globular domain of histone H3 by Dot1 is important for telomeric silencing and Sir protein association. *Genes Dev.* 2002; 16:1518–27. [PubMed: 12080090]
129. Singh RK, Kabbaj MH, Paik J, Gunjan A. Histone levels are regulated by phosphorylation and ubiquitylation-dependent proteolysis. *Nat Cell Biol.* 2009; 11:925–33. [PubMed: 19578373]
130. Cheung WL, Turner FB, Krishnamoorthy T, Wolner B, Ahn SH, Foley M, et al. Phosphorylation of histone H4 serine 1 during DNA damage requires casein kinase II in *S. cerevisiae*. *Curr Biol.* 2005; 15:656–60. [PubMed: 15823538]

131. Allard S, Utey RT, Savard J, Clarke A, Grant P, Brandl CJ, et al. NuA4, an essential transcription adaptor/histone H4 acetyltransferase complex containing Esa1p and the ATM-related cofactor Tra1p. *EMBO J.* 1999; 18:5108–19. [PubMed: 10487762]
132. Clarke AS, Lowell JE, Jacobson SJ, Pillus L. Esa1p is an essential histone acetyltransferase required for cell cycle progression. *Mol Cell Biol.* 1999; 19:2515–26. [PubMed: 10082517]
133. Parthun MR, Widom J, Gottschling DE. The major cytoplasmic histone acetyltransferase in yeast: links to chromatin replication and histone metabolism. *Cell.* 1996; 87:85–94. [PubMed: 8858151]
134. Sharma VM, Tomar RS, Dempsey AE, Reese JC. Histone deacetylases RPD3 and HOS2 regulate the transcriptional activation of DNA damage-inducible genes. *Mol Cell Biol.* 2007; 27:3199–210. [PubMed: 17296735]
135. Shia WJ, Osada S, Florens L, Swanson SK, Washburn MP, Workman JL. Characterization of the yeast trimeric-SAS acetyltransferase complex. *J Biol Chem.* 2005; 280:11987–94. [PubMed: 15659401]
136. Kimura A, Umehara T, Horikoshi M. Chromosomal gradient of histone acetylation established by Sas2p and Sir2p functions as a shield against gene silencing. *Nat Genet.* 2002; 32:370–7. [PubMed: 12410229]
137. Suka N, Luo K, Grunstein M. Sir2p and Sas2p opposingly regulate acetylation of yeast histone H4 lysine16 and spreading of heterochromatin. *Nat Genet.* 2002; 32:378–83. [PubMed: 12379856]
138. Imai S, Armstrong CM, Kaerberlein M, Guarente L. Transcriptional silencing and longevity protein Sir2 is an NAD-dependent histone deacetylase. *Nature.* 2000; 403:795–800. [PubMed: 10693811]
139. Edwards CR, Dang W, Berger SL. Histone H4 lysine 20 of *Saccharomyces cerevisiae* is monomethylated and functions in subtelomeric silencing. *Biochemistry.* 2011; 50:10473–83. [PubMed: 21985125]
140. Basnet H, Su XB, Tan Y, Meisenhelder J, Merkurjev D, Ohgi KA, et al. Tyrosine phosphorylation of histone H2A by CK2 regulates transcriptional elongation. *Nature.* 2014; 516:267–71. [PubMed: 25252977]
141. Moore JD, Yazgan O, Ataian Y, Krebs JE. Diverse roles for histone H2A modifications in DNA damage response pathways in yeast. *Genetics.* 2007; 176:15–25. [PubMed: 17028320]
142. Downs JA, Lowndes NF, Jackson SP. A role for *Saccharomyces cerevisiae* histone H2A in DNA repair. *Nature.* 2000; 408:1001–4. [PubMed: 11140636]
143. Jiang L, Smith JN, Anderson SL, Ma P, Mizzen CA, Kelleher NL. Global assessment of combinatorial post-translational modification of core histones in yeast using contemporary mass spectrometry. LYS4 trimethylation correlates with degree of acetylation on the same H3 tail. *J Biol Chem.* 2007; 282:27923–34. [PubMed: 17652096]
144. Gardner KE, Zhou L, Parra MA, Chen X, Strahl BD. Identification of lysine 37 of histone H2B as a novel site of methylation. *PLoS One.* 2011; 6:e16244. [PubMed: 21249157]
145. Robzyk K, Recht J, Osley MA. Rad6-dependent ubiquitination of histone H2B in yeast. *Science.* 2000; 287:501–4. [PubMed: 10642555]
146. Wood A, Krogan NJ, Dover J, Schneider J, Heidt J, Boateng MA, et al. Bre1, an E3 ubiquitin ligase required for recruitment and substrate selection of Rad6 at a promoter. *Mol Cell.* 2003; 11:267–74. [PubMed: 12535539]
147. Henry KW, Wyce A, Lo WS, Duggan LJ, Emre NC, Kao CF, et al. Transcriptional activation via sequential histone H2B ubiquitylation and deubiquitylation, mediated by SAGA-associated Ubp8. *Genes Dev.* 2003; 17:2648–63. [PubMed: 14563679]
148. Emre NC, Ingvarsdottir K, Wyce A, Wood A, Krogan NJ, Henry KW, et al. Maintenance of low histone ubiquitylation by Ubp10 correlates with telomere-proximal Sir2 association and gene silencing. *Mol Cell.* 2005; 17:585–94. [PubMed: 15721261]


### Highlights

- Diverse histone modifications in *S. cerevisiae* regulate cell fate decisions
- Histone modifying enzymes are targeted by intrinsic and extrinsic signaling cues to regulate transcription and chromatin reorganization
- Histone modifications play critical roles in genomic reprogramming during meiotic sporulation, filamentous growth and programmed cell death




**Figure 1. Changes in nutrient availability promote alternative differentiation pathways in diploid yeast**

Low levels of nitrogen and glucose in the presence of a nonfermentable carbon source such as acetate promote meiotic division and sporulation. When glucose levels are high but nitrogen is limiting, filamentous growth is initiated, in which pseudohyphae form and can become invasive.



**Figure 2. Regulation of *IME1* by signaling to chromatin modifiers**

Intrinsic and extrinsic signals received by the upstream control sequences 1 through 4 (*UCS1-4*) of the *IME1* regulatory region. Nutrient signaling and the transcriptional regulators  $\alpha 1/\alpha 2$  and Rme1 are essential to maintaining expression patterns of *IME1*. In diploids, the  $\alpha 1/\alpha 2$  heterodimer, encoded by the *MATa/MATα* locus releases Rme1-mediated repression of *IME1*. In haploid cells, Rme1 promotes the expression of the lncRNA *IRT1* from this region. Histone methyltransferases Set1 and Set2 are recruited co-transcriptionally and their respective methyl marks promote HDAC containing complexes Set3C and Rpd3C(L) chromatin association, repressing *IME1* expression. Tup1 is also postulated to repress *IME1*, potentially as a parallel pathway promoting HDAC recruitment, however its precise role is unknown. *IME1* expression initiates a transcriptional cascade, including inducing *NDT80*, which leads to the sporulation-specific gene expression program required for the meiotic divisions and spore formation.



**Figure 3. Chromatin-based regulation of *FLO11***  
 Low nitrogen availability regulates the *FLO11* promoter through PKA-mediated signaling to multiple transcription factors, including Flo8 (extensive signaling network regulation of *FLO11* is reviewed elsewhere [6, 77]). Transcription of the lncRNA *ICR1* approximately 3kb upstream of the *FLO11* transcription start site (TSS) occludes binding of the activator Flo8 and TBP. *ICR1* expression is repressed by Rpd3C(S) deacetylation, promoting *FLO11* activation. Deacetylation by Hda1 also promotes *FLO11* expression and dynamic regulation of H3K4me3 at the TSS by demethylase Jhd2 has been linked to *FLO11* activation.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



Histone modifications of *Saccharomyces cerevisiae*. Acetylation, methylation, phosphorylation and ubiquitination of the core histones from budding yeast are shown. For methylation, the extent of methylation is indicated in parentheses. All modifications listed have been detected using mass spectrometry and/or antibody-based methods and known writers and erasers are indicated.

**Table 1**

Histone	Residue	Modification	Writer	Eraser	References
H3	K4	Methylation (me1-3)	Set1	Jhd2	[102–107]
H3	K4	Acetylation	Gen5, Rtt109	Hst1, Sir2	[108]
H3	K9	Acetylation	Gen5 Rtt109	Rpd3	[109–112]
H3	S10	Phosphorylation	Snf1, Ipl1	Glc7	[113, 114]
H3	T11	Phosphorylation	Mek1, Pkm2	?	[71, 100]
H3	K14	Acetylation	Gen5	Rpd3	[115]
H3	K18	Acetylation	Gen5	Rpd3	[111, 115]
H3	K23	Acetylation	Gen5	Rpd3	[109, 111]
H3	K27	Acetylation	Gen5	Rpd3	[111]
H3	K36	Methylation (me1-3)	Set2	Rph1, Jhd1	[116–118]
H3	K36	Acetylation	Gen5	?	[119]
H3	T45	Phosphorylation	Cdc7	?	[120]
H3	K56	Acetylation	Rtt109	Hst3, Hst4	[121–124]
H3	K79	Methylation (me1-3)	Dot1	?	[125–128]
H3	Y99	Phosphorylation	Rad53	?	[129]
H4	S1	Phosphorylation	Cka1	?	[130]
H4	R3	Methylation (me1)	Rmt1	?	[127]
H4	K5	Methylation (me1)	Set5	?	[22]
H4	K5	Acetylation	Esa1, Hat1	Hos2, Rpd3	[111, 131–134]
H4	K8	Methylation (me1)	Set5	?	[22]
H4	K8	Acetylation	Esa1	Hos2, Rpd3	[111][131, 132, 134]
H4	K12	Methylation (me1)	Set5	?	[22]
H4	K12	Acetylation	Esa1, Hat1	Hos2, Rpd3	[111, 131, 132, 134]
H4	K16	Acetylation	Sas2	Sir2	[135–138]

Histone	Residue	Modification	Writer	Eraser	References
H4	K20	Methylation (me1)	?	?	[139]
H4	K20	Acetylation	?	?	[16]
H4	K44	Acetylation	?	?	[21]
H2A	K7	Acetylation	Esal	Rpd3	[111]
H2A	Y58	Phosphorylation	Cka1	?	[140]
H2A	Q105	Methylation	Nop1	?	[31]
H2A	S121	Phosphorylation	?	?	[141]
H2A	S128	Phosphorylation	Mec1	?	[142]
H2B	S10	Phosphorylation	Ste20	?	[91, 92]
H2B	K11	Acetylation	Gcn5	Hda1, Hos3	[93, 111]
H2B	K16	Acetylation	Gcn5, Esal	Hda1, Rpd3	[111, 143]
H2B	K37	Methylation (me2)	?	?	[144]
H2B	K123	Ubiquitination	Rad6, Bre1	Ubp8, Ubp10	[145–148]