

This work is on a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International (CC BY-NC-ND 4.0) license, <https://creativecommons.org/licenses/by-nc-nd/4.0/>. Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Complications and Patient-reported Outcomes after TRAM and DIEP Flaps: A Systematic Review and Meta-analysis

Waverley Y. He, BA*
 Leen El Eter, BS*
 Pooja Yesantharao, MS*
 Bethany Hung, BS*
 Haley Owens†
 Sarah Persing, MD, MPH*
 Justin M. Sacks, MD, MBA*‡

Introduction: Transverse rectus abdominis myocutaneous (TRAM) and deep inferior epigastric artery perforator (DIEP) flaps are the most common abdominally based breast reconstruction procedures. Each technique has its advantages and disadvantages; however, how morbidity relates to satisfaction is not well-understood. Our aim was to compare complications and patient-reported outcomes following pedicled TRAM (pTRAM), free TRAM (fTRAM), and DIEP flaps to guide flap selection.

Methods: A systematic literature search was conducted, and 2 independent reviewers identified comparative studies of abdominally based flaps. Data were extracted on patient characteristics, complications, and patient-reported outcomes. Meta-analyses were conducted using random effects modeling with the DerSimonian and Laird method.

Results: The search retrieved 5090 articles, of which 18 were included in this review. pTRAM flaps trended toward a higher risk of abdominal bulge/hernia compared with DIEP flaps, particularly in low-volume hospitals. While fTRAM flaps had a higher risk of abdominal morbidity compared with DIEP flaps, relative risk decreased when obese patients were excluded and when only muscle-sparing fTRAM flaps were compared. Muscle-sparing flaps had a higher risk of flap loss than fTRAM flaps. Compared with DIEP flaps, pTRAM flaps were associated with lower general satisfaction but comparable emotional well-being.

Conclusions: Our findings indicate that safety and satisfaction following abdominally based breast reconstruction depend on flap type and patient characteristics. When possible, DIEP or muscle-sparing fTRAM flaps should be performed for obese patients to decrease the risk of abdominal bulge/hernia. Although pTRAM flaps are associated with a greater risk of flap loss, they are still an appropriate option when microsurgery is not available. (*Plast Reconstr Surg Glob Open* 2020;8:e3120; doi: [10.1097/GOX.00000000000003120](https://doi.org/10.1097/GOX.00000000000003120); Published online 29 October 2020.)

INTRODUCTION

Although implant-based reconstruction is the most common approach to breast reconstruction,^{1,2} autologous reconstruction is also well established and is associated with increased patient satisfaction and quality of life.³ Of

women who elect to undergo breast reconstruction following mastectomy, 13 percent had autologous reconstruction using abdominally based tissue.¹³ This includes pedicled transverse rectus abdominis (pTRAM), free TRAM (fTRAM), and deep inferior epigastric (DIEP) flap reconstructions. With greater public concern surrounding implant-based reconstruction and recent FDA initiatives to explore breast implant safety,^{2,3} we anticipate a rise in autologous reconstruction in the near future. Thus, there exists a need to re-examine flap selection criteria based on contemporary literature.

From the *Department of Plastic and Reconstructive Surgery, The Johns Hopkins University School of Medicine, Baltimore, Md.; †Department of Biological Sciences, University of Maryland, Baltimore County, Baltimore, Md.; and ‡Division of Plastic and Reconstructive Surgery, Washington University School of Medicine in St. Louis, St. Louis, Mo.

Received for publication July 22, 2020; accepted July 27, 2020.

Copyright © 2020 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of The American Society of Plastic Surgeons. This is an open-access article distributed under the terms of the [Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 \(CCBY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal. [10.1097/GOX.00000000000003120](https://doi.org/10.1097/GOX.00000000000003120)

Disclosure: The authors have no financial interest to declare in relation to the content of this article. Dr. Sacks received unrestricted research funding from Vioptix and is co-founder of Lifesprout, Inc.

Related Digital Media are available in the full-text version of the article on www.PRSGlobalOpen.com.

Previous studies show that DIEP and muscle-sparing (MS) fTRAM flaps have a decreased incidence of abdominal bulge/hernia compared with pTRAM and non-muscle-sparing fTRAM flaps, suggesting that harvesting less muscle results in better donor-site outcomes.^{4,5} DIEP flaps, while sparing abdominal rectus muscle, are associated with a higher rate of flap loss and fat necrosis, presumably from a less robust blood flow compared with the fTRAM flap and technical challenges of perforator dissection.^{6–8} Despite these differences in complication rates, little is known regarding how postoperative morbidity correlates with patient-reported outcomes. While some surveys show no difference in patient satisfaction following DIEP and non-muscle-sparing fTRAM flaps,^{9–12} there is less literature comparing DIEP flaps to MS fTRAM or pTRAM flaps.

Additionally, because most flap selection criteria rely on intraoperative indications or conventional recommendations based on patient characteristics, the relationship between preoperative risk factors and complications for abdominally based flap reconstruction remains unclear. In particular, while obesity and lower hospital procedural volume status have been shown to lead to overall higher risk of postoperative morbidity,^{13–19} impact on safety outcomes has not yet been summarized according to abdominally based flap type.

To our knowledge to date, a systematic review and meta-analysis comparing complications and quality of life for all major abdominally based breast reconstruction techniques, and assessing the impact of preoperative risk factors, have not yet been performed.^{4–6,16} An improved understanding of these outcomes is critical so that patients and providers can have informed discussions about expectations for recovery during the shared decision-making process. The aim of this study was to compare safety and patient-reported outcomes following pedicled (pTRAM), free TRAM (fTRAM), and DIEP flaps, as well as to assess whether these differ on the basis of obesity and hospital procedural volume status, to guide optimal flap selection.

METHODS

Study Selection and Data Extraction

MEDLINE, EMBASE, the Cochrane Central Register of Controlled Trials, Web of Science, and Scopus were systematically queried for studies of breast reconstruction following mastectomy for cancer, according to our systematic review protocol (PROSPERO registration CRD42020147475). Covidence (Melbourne, Australia) was used to manage study selection, quality assessment, and data extraction. Study selection was based on predefined eligibility criteria in a Population, Intervention, Comparison, Outcome, Timing, and Setting (PICOTS) framework (See appendix, Supplemental Digital Content 1, which displays a PICOTS (population, interventions, comparators, outcomes, timing, and setting) for the key questions. Appendix B. Risk of bias across included studies of complications and patient-reported outcomes. <http://links.lww.com/PRSGO/B491>). Following PRISMA guidelines, 2 reviewers independently screened abstracts and full text articles to identify comparative studies of pedicled,

free, and muscle-sparing free TRAM and DIEP flap breast reconstructions published after 2000. References of included studies were searched for additional relevant titles.

Two reviewers independently assessed the risk of bias for individual studies using the Risk of Bias in Non-Randomised Studies of Interventions (ROBINS-I),²⁰ and extracted data on general study characteristics, patient characteristics, and complications and patient-reported outcomes. Differences between reviewers at any stage were resolved through consensus.

Statistical Analysis

To standardize results, we conducted the meta-analyses for complications to reflect the authors' definitions from each study. Hospital procedural volume was calculated as the number of abdominally-based autologous breast reconstructions performed per year during the study period, and institutions in the lower 50th percentile were classified as low-volume.

Meta-analyses for outcomes were conducted using a random effects model with the DerSimonian and Laird method when there were at least 2 sufficiently homogeneous studies. Associations between flap type (ie, pTRAM, fTRAM, and DIEP flaps) and complications were reported as effect sizes with 95% confidence intervals. A post-hoc sensitivity analysis was conducted using the Hartung-Knapp-Sidik-Jonkman method. Subgroup analyses were conducted using flap subtypes (ie, MS fTRAM flaps) as well as patient- and hospital-level factors. A Haldane-Anscombe correction was used for outcomes with small counts. All meta-analyses were conducted using Microsoft Excel (version 16).

RESULTS

The search retrieved 5090 unique articles, of which 4132 were excluded during abstract screening and 940 were excluded during full-text screening (Fig. 1). A total of 18 articles were ultimately included in this systematic review.^{9–11,13,14,21–32} Overall risk of bias was low for 5 and moderate for 11 studies (see Supplemental Digital Content 1, <http://links.lww.com/PRSGO/B491>). Fifteen articles assessed safety,^{9–11,13,14,21–30,32} and 8 provided patient-reported outcomes (Table 1).^{9–12,14,25,31,32} All were cohort studies, with the exception of 1 cross-sectional study.³¹

Complications

Pooled estimates demonstrated a trend toward higher abdominal bulge/hernia rates (RR = 2.82, 95% CI = 0.83–4.80) following pTRAM compared to DIEP flaps, which increased in a subgroup analysis with only low-volume hospitals (RR = 3.08, 95% CI = 0.46–5.70) (Table 2). Similarly, there were trends toward higher relative risks of flap loss, mastectomy skin flap necrosis, fat necrosis, and wound healing complications following pTRAM compared with DIEP flaps in low-volume hospitals, although these did not reach statistical significance.

Patients undergoing fTRAM were 2.87 times more likely to experience abdominal bulge/hernia (95% CI = 1.73–4.00) than patients undergoing DIEP flap




Fig. 1. Summary of the study selection process, showing the included 18 unique articles.

reconstruction (Table 3). However, relative risk decreased to 2.35 when obese patients were excluded in a subgroup analysis (95% CI = 1.49–3.21) and 2.55 when only MS fTRAM flaps were compared to DIEP flaps (95% CI = 1.42–3.67) (Fig. 2). DIEP flaps were associated with a greater risk of flap loss (ES = 2.67, 95% CI = 1.00–4.34) than fTRAM, although relative risk decreased (RR = 1.93, 95% CI = 0.26–3.61) in a comparison of DIEP with only MS fTRAM flaps. Differences in mastectomy skin flap necrosis, fat necrosis, infection, hematoma/seroma, and wound healing complications after fTRAM and DIEP flap reconstructions did not reach statistical significance.

Only one study reported on the development of hypertrophic scarring and keloids. It was found that breast scars developed after 2.3% of DIEP flap reconstructions, while none developed after pedicled or free TRAM flap reconstructions.¹⁰ A comparable, low rate of abdominal donor-site scarring was seen after DIEP (1.4%), pedicled TRAM (1.3%), and free TRAM (1.1%) flap reconstructions.

Patient-reported Outcomes

Four articles reported on psychosocial outcomes;^{9,10,14,25} 4 on satisfaction outcomes;^{9,10,31,32} 5 on physical well-being outcomes;^{9–12,14} and 1 on pain (Table 4).¹¹ Studies used the BREAST-Q^{9,10} and Short Form-36 (SF-36)^{14,25} to measure psychosocial outcomes. In the Mastectomy Reconstruction Outcomes Consortium study, BREAST-Q psychosocial and sexual well-being scores in pTRAM and fTRAM groups were not significantly different from scores in the DIEP group at both the first and second year post-reconstruction after adjusting for covariates,¹⁰ and scores improved with time regardless of reconstruction approach. Similar conclusions were observed in another multi-institution study.⁹ Psychosocial and sexual well-being scores were also

not significantly different between MS fTRAM and DIEP groups. In one study, patients receiving bilateral pTRAM had higher SF-36 energy and well-being scores compared with those receiving DIEP flaps ($P < 0.05$ and $P = 0.06$, respectively).²⁵ In another study, there were minimal differences in the SF-36 mental health composite score between MS fTRAM and DIEP flaps at long-term follow-up.¹⁴ Although obese and non-obese patients had similar preoperative scores, the SF-36 mental health composite score improved in non-obese patients and worsened in obese patients ($P = 0.02$).

Two studies measured satisfaction through a 7-item questionnaire developed for the Michigan Breast Reconstruction Outcome Study,^{31,32} and two studies measured satisfaction using the BREAST-Q.^{9,10} Momoh et al. demonstrated that recipients of DIEP flap reconstruction had significantly higher general satisfaction than recipients of pTRAM flap reconstruction ($P = 0.04$).³² This likelihood was even more pronounced in the bilateral DIEP group compared with the bilateral pTRAM group ($P = 0.0095$). In contrast, another study found that there were no significant differences in general satisfaction between pTRAM and DIEP groups.³¹ Both studies suggest similar aesthetic satisfaction rates across different approaches to reconstruction. Macadam et al. demonstrated that there was significantly higher satisfaction with long-term outcome in the DIEP group compared with the pTRAM group ($P = 0.015$) after controlling for patient characteristics.⁹ This finding did not reach significance when comparing fTRAM or MS fTRAM with DIEP flaps, and the difference in satisfaction with breasts did not vary between pTRAM, fTRAM, and DIEP groups after 1 and 2 years.^{9,10}

Physical outcomes were assessed by the BREAST-Q and SF-36. In one study, DIEP flap patients experienced

Table 1. Summary of Included Studies

Year of Publication	Author	Country	Study Design	Intervention Groups Compared	Complications	Patient-reported Outcomes	Comments
2002	Nahabedian et al.	United States	Retrospective cohort	• Free TRAM (n = 118) • DIEP (n = 17)	X		
2005	Nahabedian et al.	United States	Retrospective cohort	• Muscle-sparing free TRAM (n = 89) • DIEP (n = 88)	X		
2006	Bajaj et al.	United States	Retrospective cohort	• Muscle-sparing free TRAM (n = 124) • DIEP (n = 35)	X		
2006	Garvey et al.	United States	Retrospective cohort	• Pedicled TRAM (n = 94) • DIEP (n = 96)	X		Unilateral flaps
2007	Chen et al.	United States	Retrospective cohort	• Free TRAM (n = 114) • DIEP (n = 29)	X		
2007	Schaverien et al.	United Kingdom	Retrospective cohort	• Free TRAM (n = 30) • DIEP (n = 30)	X	X	Unilateral flaps
2010	Chun et al.	United States	Retrospective cohort	• Pedicled TRAM (n = 105) • DIEP (n = 58)	X	X	Bilateral flaps
2010	Nelson et al.	United States	Retrospective cohort	• Muscle-sparing free TRAM (n = 91) • DIEP (n = 53)	X		
2012	Momoh et al.	United States	Retrospective cohort	• Pedicled TRAM (n = 179) • DIEP (n = 167)	X	X	
2013	Fischer et al.	United States	Retrospective cohort	• Muscle-sparing free TRAM (n = 300) • DIEP (n = 105)	X		Obese patients
2014	Benditte-Klepetko et al.	Netherlands	Retrospective cohort	• Free TRAM (n = 22) • DIEP (n = 18)	X		
2015	Weichman et al.	United States and Canada	Prospective cohort	• Pedicled TRAM (n = 83) • Free TRAM (n = 91) • DIEP (n = 296)		X	
2015	Yang et al.	China	Cross-sectional	• Pedicled TRAM (n = 44) • DIEP (n = 62)		X	
2016	Knox et al.	Canada	Retrospective cohort	• Pedicled TRAM (n = 377) • DIEP (n = 130)	X		
2016	Macadam et al.	United States and Canada	Retrospective cohort	• Pedicled TRAM (n = 683) • Free TRAM (MS-0) (n = 144) • Muscle-sparing free TRAM (n = 293) • DIEP (n = 670)	X	X	
2018	Erdmann-Sager et al.	United States and Canada	Prospective cohort	• Pedicled TRAM (n = 89) • Free TRAM (n = 115) • DIEP (n = 445)		X	
2018	Xu et al.	China	Prospective cohort	• Pedicled TRAM (n = 39) • DIEP (n = 9)	X		Immediate, unilateral flaps
2019	Nelson et al.	United States	Prospective cohort	• Muscle-sparing free TRAM (n = 35) • DIEP (n = 13)	X	X	

significantly higher abdominal physical well-being compared with pTRAM flap patients after controlling for cofounders ($P < 0.001$).⁹ Differences did not reach significance when comparing the DIEP flap to fTRAM and MS fTRAM flaps. Similarly, no differences in abdominal well-being scores between MS fTRAM flaps and DIEP flaps were noted in another study.¹⁴ The Mastectomy Reconstruction Outcomes Consortium study showed higher abdomen-specific scores for DIEP compared with both pTRAM and fTRAM flaps one year postoperatively ($P = 0.078$ and $P = 0.051$, respectively) and significantly higher scores two years postoperatively ($P = 0.006$ and $P = 0.037$) after controlling for laterality, although in all groups unadjusted postoperative scores were lower than preoperative scores.¹⁰ The Mastectomy Reconstruction Outcomes Consortium study demonstrated that unadjusted physical chest well-being decreased from preoperative to 1 week postoperatively in pTRAM, fTRAM, and DIEP groups, and did not return to baseline levels at 3 months ($P < 0.001$).¹² Multivariable analysis showed there was no significant difference in physical well-being of chest scores between

groups at 1 year postoperatively.¹⁰ Interestingly, however, the pTRAM group reported significantly lower scores at 2 years postoperatively compared to the DIEP group ($P = 0.04$).

In one study, the SF-36 physical health composite score significantly improved over time for both unilateral and bilateral MS fTRAM and DIEP flaps ($P < 0.05$), although no significant differences were noted during a direct comparison of flap type or laterality.¹⁴ Obesity lead to a significant worsening in the physical health composite score over time ($P = 0.003$), while scores improved in non-obese patients. In another study, no significant differences were found between fTRAM and DIEP groups for scores on the SF-36 physical functioning or physical role limitations scales.¹¹ Patients also reported that they rarely suffered from postoperative abdominal pain for a mean duration of 1–3 months.

DISCUSSION

This review evaluates abdominally based autologous breast reconstruction techniques and provides updated

Table 2. Pooled Complication Rates for pTRAM versus DIEP Flaps

Outcomes	Studies, N	pTRAM		DIEP		Effect Size	95% CI	I ²
		No. Events	No. Patients	No. Events	No. Patients			
Total or partial flap loss	6	90	1562	76	1233	1.03	-0.81 to 2.88	36.62%
Low-volume hospitals only	4	22	774	17	505	1.15	-1.08 to 3.39	60.21%
Mastectomy skin flap necrosis	3	80	680	56	409	0.36	-1.39 to 2.11	0
Low-volume hospitals only	2	69	483	34	192	0.86	-2.30 to 4.02	0
Fat necrosis (breast)	6	342	1667	210	1291	1.43	-0.33 to 3.18	90.50%
Low-volume hospitals only	4	147	774	76	505	1.63	-0.53 to 3.80	93.00%
Infection (breast)	5	184	1457	93	1175	1.63	-0.36 to 3.62	86.59%
Low-volume hospitals only	4	78	774	51	505	1.39	-0.87 to 3.65	77.24%
Hematoma/seroma (breast)	5	96	1468	105	1137	0.76	-1.21 to 2.73	30.37%
Low-volume hospitals only	4	70	785	49	467	0.84	-1.39 to 3.07	0
Wound healing complications (breast)	4	97	774	29	505	1.49	-0.78 to 3.77	80.18%
Low-volume hospitals only	3	91	577	23	288	1.64	-1.03 to 4.31	83.62%
Abdominal bulge/hernia	5	174	1100	74	1000	2.82	0.83 to 4.80	85.70%
Low-volume hospitals only	3	55	312	15	272	3.08	0.46 to 5.70	48.08%
Wound healing complications (donor-site)	3	129	576	106	284	0.59	-1.86 to 3.04	0
Low-volume hospitals only	2	120	471	94	226	0.68	-2.30 to 3.66	0

Table 3. Pooled Complication Rates for fTRAM versus DIEP Flaps

Outcomes	Studies, N	DIEP		fTRAM		Effect Size	95% CI	I ²
		No. Events	No. Patients	No. Events	No. Patients			
Total or partial flap loss	8	73	1083	50	1401	2.67	1.00 to 4.34	87.62%
Without obese patients	7	68	978	45	1101	2.64	0.83 to 4.45	88.57%
Muscle-sparing only	4	65	899	21	632	1.93	0.26 to 3.61	87.57%

Outcomes	Studies, N	fTRAM		DIEP		Effect Size	95% CI	I ²
		No. Events	No. Patients	No. Events	No. Patients			
Mastectomy skin flap necrosis	4	154	604	58	235	0.96	-1.31 to 3.28	0
Without obese patients	3	28	304	8	130	1.03	-1.71 to 3.77	0
Fat necrosis (breast)	7	133	1430	143	1065	0.80	-0.83 to 2.44	0
Without obese patients	6	125	1130	133	960	0.89	-0.88 to 2.66	0
Muscle-sparing only	4	67	684	126	899	0.83	-1.32 to 2.98	0
Infection (breast)	5	63	1041	47	905	2.60	0.54 to 4.65	71.51%
Without obese patients	4	51	741	44	800	2.93	0.61 to 5.26	78.60%
Muscle-sparing only	2	42	560	45	364	2.10	-1.04 to 5.23	70.20%
Hematoma/seroma (breast)	5	55	1041	72	905	0.80	-1.20 to 2.80	15.91%
Without obese patients	4	48	741	68	800	0.85	-1.40 to 3.10	36.80%
Muscle-sparing only	2	25	560	41	364	0.43	-2.59 to 3.46	0
Wound healing complications, any (breast)	3	132	582	52	217	0.93	-1.74 to 3.59	0
Without obese patients	2	6	282	2	112	0.96	-2.49 to 4.40	0
Abdominal bulge/hernia	9	94	1253	40	1030	2.87	1.73 to 4.00	70.48%
Without obese patients	8	76	953	39	925	2.35	1.49 to 3.21	41.44%
Muscle-sparing only	5	56	612	36	855	2.55	1.42 to 3.67	62.83%

flap selection criteria based on complications and patient-reported outcomes. Consistent with previous reports in the literature, we found a trend toward greater risk of abdominal bulge/hernia following pTRAM compared with DIEP flaps, reinforcing the notion that losing muscle makes abdominal fascia prone to weakness.^{4,5} Vascular complications, including flap loss and fat necrosis, also trend towards greater likelihood following pTRAM, suggesting that tunneling a pedicled flap up to the chest compromises blood flow via kinking or compression of the superior epigastric vessels, or that the pedicle blood supply provides decreased perfusion.^{7,28,33} In contrast, DIEP flaps benefit from the enhanced blood supply of the inferior epigastric vessels or the flow dynamics from the internal mammary or thoracodorsal system. These relative risks are increased in low-volume hospitals, perhaps because low-volume hospitals see patients with fewer comorbidities, for

whom microsurgery is preferred.^{4,5} Another possibility is that because DIEP flap reconstructions have an increased mean surgical time and steeper learning curve,²² patients undergoing this procedure benefit from more careful intraoperative technique and postoperative monitoring. Albornoz et al. found that in higher-volume hospitals, where surgical technique benefits from greater experience,³⁴ patients undergoing microvascular reconstruction were also less likely to experience complications than those undergoing autologous reconstructions in general.¹⁸ Further studies examining the association between hospital volume and complication rates following abdominally-based breast reconstruction, and differentiating between pTRAM, fTRAM, and DIEP flaps, are necessary to create flap-specific guidelines for improving safety.

There is a significantly greater risk of abdominal bulge/hernia following fTRAM compared to DIEP and




Fig. 2. Higher relative risk of abdominal bulge and hernia shown for pTRAM and fTRAM flaps compared with that for DIEP flaps on meta-analysis, particularly for obese patients. The comparisons shown are (a) pTRAM vs DIEP, (b) fTRAM vs DIEP, (c) fTRAM vs DIEP subgroup analysis excluding 1 study using obese patients, and (d) MS fTRAM vs DIEP subgroup analysis. The horizontal lines represent the 95% confidence intervals for each study.

Table 4. Summary of Patient-Reported Outcomes Findings

Outcome	Study	Direction of Finding			Statistical Adjustment	Measure
		pTRAM versus DIEP	fTRAM versus DIEP	MS fTRAM versus DIEP		
Psychosocial outcomes	Chun et al., 2010	+			No	SF-36 energy/fatigue
	Macadam et al., 2016	=	=	=	Yes	SF-36 emotional well-being
	Erdmann-Sager et al., 2018	=	=		Yes	BREAST-Q psychosocial well-being
Satisfaction	Nelson et al., 2019			=	No	BREAST-Q sexual well-being
	Momoh et al., 2012	General satisfaction: -			No	BREAST-Q psychosocial well-being
	Yang et al., 2015	Aesthetic satisfaction: =			No	BREAST-Q sexual well-being
		=			No	SF-36 mental health composite
	Macadam et al., 2016	Satisfaction breasts: =	=	=	Yes	MBROS general satisfaction
	Erdmann-Sager et al., 2018	Satisfaction outcome: -	=		Yes	MBROS aesthetic satisfaction
Physical well-being	Schaverien et al., 2007		=		No	MBROS general satisfaction
	Weichman et al., 2015	=	=		Yes	MBROS aesthetic satisfaction
	Macadam et al., 2016	-	=	=	Yes	BREAST-Q satisfaction breasts
	Erdmann-Sager et al., 2018	PWB chest (one year): =	PWB chest: =		Yes	BREAST-Q satisfaction outcome
		PWB chest (two year): -	PWB abdomen: -			BREAST-Q satisfaction breasts
	Nelson et al., 2019	PWB abdomen: -		=	No	BREAST-Q PWB chest
Pain	Schaverien et al., 2007		=		No	SF-36 physical functioning

Bold entries indicate the studies finding statistically significant differences in patient-reported outcome scores between flaps (i.e. the direction of finding is either + or -).

+ score for first reconstruction technique higher than score for second reconstruction technique.

- score for second reconstruction technique higher than score for first reconstruction technique.

= scores equivalent for both reconstruction techniques.

SF-36, short-form 36; MBROS, Michigan Breast Reconstruction Outcome Study.

MS fTRAM flaps. Notably, relative to non-obese patients, obese patients experience a higher risk of abdominal bulge/hernia with fTRAM compared with DIEP flaps; so DIEP flaps should be performed in this population if possible. Obesity may increase the risk of abdominal bulge/hernia following fTRAM flaps due to increased tension being placed on the weakened abdominal fascia.¹⁵ These healing problems may be mitigated following DIEP flaps, which leave less of an abdominal wall defect.³⁵ Our findings also indicate that muscle conservation increases the relative risk of flap loss. Therefore, DIEPs pose a greater risk of flap loss compared with MS fTRAM flaps, while fTRAM flaps pose the least risk. As fewer perforating vessels are included with DIEP compared with free TRAM flaps, there is potential for less robust vascular supply.^{6,7}

Studies reported significant differences in patient-reported outcomes following TRAM and DIEP flap reconstructions. Patients undergoing pTRAM flaps tended to report lower general satisfaction; lower physical well-being, as measured by the BREAST-Q; and higher energy and emotional well-being, as measured by the SF-36. However, there were no differences in psychosocial and sexual well-being, or breast and aesthetic satisfaction. This indicates that although patients receiving pTRAM flap reconstruction have higher rates of abdominal morbidity

and are less satisfied with their physical recovery process,¹² they are still ultimately satisfied with their aesthetic outcome. They may even be more energetic and positive immediately postoperatively, following shorter operating times and lengths of stay.^{16,25,36} Thus, although free abdominally based flaps have a more favorable complication profile, pTRAM flaps may still be useful for certain patients, particularly when microsurgery is not an option secondary to surgeon expertise, comfort level or patient preference based on risk-weighted decisions. Patient-reported outcomes following fTRAM, MS fTRAM, and DIEP flaps were similar, with the exception of higher BREAST-Q abdominal physical well-being scores in DIEP flaps compared with fTRAM flaps. This again suggests that while surgeons are highly concerned with donor-site morbidity, patients have a more holistic perspective toward well-being.

On the basis of these findings, we outline several recommendations for preoperative flap selection (Fig. 3). To decrease risk of abdominal bulge/hernia, muscle-sparing flaps (particularly DIEP, but also MS fTRAM flaps) should be considered before fTRAM among women who are appropriate microsurgical candidates, particularly those who are obese.¹³ Based on a previous meta-analysis, raising muscle-sparing flaps in obese patients may also have the additional benefit of lowering risk for flap loss and fat




Fig. 3. Higher relative risk of total and partial flap loss shown for DIEP flaps compared with fTRAM flaps on meta-analysis. The comparisons shown are (a) DIEP vs fTRAM and (b) DIEP vs MS fTRAM subgroup analysis.

necrosis.¹⁶ Our findings with regard to DIEP flaps in obese patients contrast with conventional recommendations, suggesting that TRAM flaps should be raised for obese patients.^{27,29} Additionally, given higher rates of abdominal morbidity, the use of synthetic mesh for fascial closure may have particular utility in obese patients and should be investigated in future work.^{37,38} To prevent vascular complications such as flap loss, planned DIEP and MS fTRAM flap reconstructions should be converted to fTRAM flaps if surgeons are unable to identify perforators greater than 1.5 mm in the same intermuscular septum.^{11,23,26,27,29} If microsurgical breast reconstruction is unavailable or if the patient is too unstable to undergo a prolonged operation, pTRAM flaps should be performed despite the trend toward an increased risk of flap-related and abdominal morbidity. Finally, while this review does not explore non-clinical patient characteristics such as preoperative activity level, it is important to remember that patient goals are vital to the flap selection process. For example, a woman who desires less fatigue and emotional toll during recovery may opt to pursue pTRAM flap reconstruction despite the increased risk of abdominal morbidity.^{16,25,36}

Because the current literature is limited by a lack of randomized controlled trials and because patients undergoing different flap reconstruction techniques are often not

equivalent surgical candidates, it is particularly important for cohort studies to report patient demographic, clinical, and treatment characteristics as well as to use analysis methods that control for these characteristics. Yet more than half of the included studies were characterized by a moderate or severe bias due to confounding (see **Supplemental Digital Content 1**, <http://links.lww.com/PRSGO/B491>). Our statistical analysis attempted to control for obesity and hospital procedure volume status, and our findings regarding complication rates are consistent with those reported by consortium and national database studies,^{10,39,40} which are less likely to be biased by single-surgeon and hospital factors. Due to a lack of extractable data, we were unable to control for other preoperative risk factors. More high-quality, prospective cohort studies are therefore needed to demonstrate how additional patient characteristics impact complication rates and patient well-being.

The lack of standardization in flap harvest techniques creates another potential source of bias. Due to surgeon-specific variations in technique, MS fTRAM flaps at one institution may be coded as DIEP flaps at another institution, thereby confounding the results. This review demonstrates that increased muscle harvest is associated with increased relative risk of abdominal bulge/hernia and other complications. However, several studies do not




Fig. 4. Evidence-based decision-making algorithm for abdominally based breast reconstruction options (ie, pTRAM, fTRAM, MS fTRAM, and DIEP flaps).

report which degree of muscle-sparing surgery they perform during MS fTRAM flap reconstruction.^{9,13,14,23,26,27} As muscle-sparing flap harvest techniques become more popular and new variations are developed,⁴¹ for the sake of accurate outcomes reporting and patient counseling, it is increasingly important for surgeons to adhere to a uniform classification system of muscle wall preservation. DellaCroce et al. has expanded on the traditional classification system by offering a specific definition for the point at which muscle transection in the DIEP flap harvest equates to a muscle-sparing fTRAM flap.

The use of the abdominal wall for autologous breast reconstruction is an optimized procedure for both aesthetic and functional outcomes. Our findings demonstrate that complications and patient-reported outcomes following abdominally based breast reconstruction techniques depend on flap type as well as patient- (ie, obesity) and hospital-level factors (ie, procedure volume). When possible, DIEP and MS fTRAM reconstructions should be performed for obese patients to decrease risk of abdominal bulge/hernia. Although pTRAM flaps are associated with a greater risk of complications, particularly in low-volume hospitals, they may still be an appropriate option when microsurgery is not available. When selecting the appropriate abdominally based flap type for autologous breast reconstruction, surgeon expertise must be balanced with patient preferences, and risks of flap- and donor-site morbidity must be weighed with patient wishes and comorbidities. This systematic review is a useful guide that will inform both patient and provider in the shared decision-making process. Maintaining transparency regarding potential safety and satisfaction outcomes will allow these discussions to take place in an open and honest manner.

Justin M. Sacks, MD, MBA

Division of Plastic and Reconstructive Surgery
Washington University School of Medicine in St. Louis
St. Louis, MO 63110
E-mail: jmsacks@wustl.edu

ACKNOWLEDGMENT

We thank Stella Seal, MLS, for her assistance in developing and conducting a search strategy.

REFERENCES

1. American Society of Plastic Surgeons. 2018 Plastic Surgery Statistics Report. 2018. Available at <https://www.plasticsurgery.org/documents/News/Statistics/2018/plastic-surgery-statistics-full-report-2018.pdf>.
2. Marra A, Viale G, Pileri SA, et al. Breast implant-associated anaplastic large cell lymphoma: a comprehensive review. *Cancer Treat Rev*. 2020;84:101963.
3. U.S. Food & Drug Administration. Breast Implants – Certain Labeling Recommendations to Improve Patient Communication. 2019. Available at <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/breast-implants-certain-labeling-recommendations-improve-patient-communication>.
4. Egeberg A, Rasmussen MK, Sørensen JA. Comparing the donor-site morbidity using DIEP, SIEA or MS-TRAM flaps for breast reconstructive surgery: a meta-analysis. *J Plast Reconstr Aesthet Surg*. 2012;65:1474–1480.
5. Atisha D, Alderman AK. A systematic review of abdominal wall function following abdominal flaps for postmastectomy breast reconstruction. *Ann Plast Surg*. 2009;63:222–230.
6. Sailon AM, Schachar JS, Levine JP. Free transverse rectus abdominis myocutaneous and deep inferior epigastric perforator flaps for breast reconstruction: a systematic review of flap complication rates and donor-site morbidity. *Ann Plast Surg*. 2009;62:560–563.
7. Kroll SS. Fat necrosis in free transverse rectus abdominis myocutaneous and deep inferior epigastric perforator flaps. *Plast Reconstr Surg*. 2000;106:576–583.
8. Mulvey CL, Cooney CM, Daily FF, et al. Increased flap weight and decreased perforator number predict fat necrosis in DIEP breast reconstruction. *Plast Reconstr Surg Glob Open*. 2013;1:1–7.
9. Macadam SA, Zhong T, Weichman K, et al. Quality of life and patient-reported outcomes in breast cancer survivors: a multicenter comparison of four abdominally based autologous reconstruction methods. *Plast Reconstr Surg*. 2016;137:758–771.
10. Erdmann-Sager J, Wilkins EG, Pusic AL, et al. Complications and patient-reported outcomes after abdominally based breast reconstruction: results of the Mastectomy Reconstruction Outcomes Consortium Study. *Plast Reconstr Surg*. 2018;141:271–281.

11. Schaverien MV, Perks AG, McCulley SJ. Comparison of outcomes and donor-site morbidity in unilateral free TRAM versus DIEP flap breast reconstruction. *J Plast Reconstr Aesthet Surg*. 2007;60:1219–1224.
12. Weichman KE, Hamill JB, Kim HM, et al. Understanding the recovery phase of breast reconstructions: patient-reported outcomes correlated to the type and timing of reconstruction. *J Plast Reconstr Aesthet Surg*. 2015;68:1370–1378.
13. Fischer JP, Nelson JA, Sieber B, et al. Free tissue transfer in the obese patient: an outcome and cost analysis in 1258 consecutive abdominally based reconstructions. *Plast Reconstr Surg*. 2013;131:681e–692e.
14. Nelson JA, Tecci MG, Lanni MA, et al. Function and strength after free abdominally based breast reconstruction: a 10-year follow-up. *Plast Reconstr Surg*. 2019;143:22e–31e.
15. Chang D, Wang B, Robb G, et al. Effect of obesity on flap and donor-site complications in free transverse rectus abdominis myocutaneous flap breast reconstruction. *Plast Reconstr Surg*. 2002;105:1199–1200.
16. Lee KT, Mun GH. Effects of obesity on postoperative complications after breast reconstruction using free muscle-sparing transverse rectus abdominis myocutaneous, deep inferior epigastric perforator, and superficial inferior epigastric artery flap: a systematic review and meta-analysis. *Ann Plast Surg*. 2016;76:576–584.
17. Chattha A, Chen AD, Muste J, et al. Revisiting the relationship between hospital case volume and outcomes in abdominally based free flap breast reconstruction. *Ann Plast Surg*. 2020;85:397–401.
18. Alborno CR, Cordeiro PG, Hishon L, et al. A nationwide analysis of the relationship between hospital volume and outcome for autologous breast reconstruction. *Plast Reconstr Surg*. 2013;132:192–200.
19. Billig JJ, Lu Y, Momoh AO, et al. A nationwide analysis of cost variation for autologous free flap breast reconstruction. *JAMA Surg*. 2017;152:1039–1047.
20. Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in non-randomised studies of interventions. *BMJ*. 2016;355:i4919.
21. Benditte-Klepetko HC, Lutgendorff F, Kästenbauer T, et al. Analysis of patient satisfaction and donor-site morbidity after different types of breast reconstruction. *Scand J Surg*. 2014;103:249–255.
22. Knox AD, Ho AL, Leung L, et al. Comparison of outcomes following autologous breast reconstruction using the DIEP and pedicled TRAM flaps: a 12-year clinical retrospective study and literature review. *Plast Reconstr Surg*. 2016;138:16–28.
23. Nelson JA, Guo Y, Sonnad SS, et al. A Comparison between DIEP and muscle-sparing free TRAM flaps in breast reconstruction: a single surgeon's recent experience. *Plast Reconstr Surg*. 2010;126:1428–1435.
24. Chen CM, Halvorson EG, Disa JJ, et al. Immediate postoperative complications in DIEP versus free/muscle-sparing TRAM flaps. *Plast Reconstr Surg*. 2007;120:1477–1482.
25. Chun YS, Sinha I, Turko A, et al. Comparison of morbidity, functional outcome, and satisfaction following bilateral TRAM versus bilateral DIEP flap breast reconstruction. *Plast Reconstr Surg*. 2010;126:1133–1141.
26. Nahabedian MY, Tsangaris T, Momen B. Breast reconstruction with the DIEP flap or the muscle-sparing (MS-2) free TRAM flap: is there a difference? *Plast Reconstr Surg*. 2005;115:436–444; discussion 445.
27. Bajaj AK, Chevray PM, Chang DW. Comparison of donor-site complications and functional outcomes in free muscle-sparing TRAM flap and free DIEP flap breast reconstruction. *Plast Reconstr Surg*. 2006;117:737–746; discussion 747.
28. Garvey PB, Buchel EW, Pockaj BA, et al. DIEP and pedicled TRAM flaps: a comparison of outcomes. *Plast Reconstr Surg*. 2006;117:1711–1719; discussion 1720.
29. Nahabedian MY, Momen B, Galdino G, et al. Breast reconstruction with the free TRAM or DIEP flap: patient selection, choice of flap, and outcome. *Plast Reconstr Surg*. 2002;110:466–475; discussion 476.
30. Xu F, Sun H, Zhang C, et al. Comparison of surgical complication between immediate implant and autologous breast reconstruction after mastectomy: a multicenter study of 426 cases. *J Surg Oncol*. 2018;118:953–958.
31. Yang B, Li L, Yan W, et al. The type of breast reconstruction may not influence patient satisfaction in the Chinese population: a single institutional experience. *PLoS One*. 2015;10:1–11.
32. Momoh AO, Colakoglu S, Westvik TS, et al. Analysis of complications and patient satisfaction in pedicled transverse rectus abdominis myocutaneous and deep inferior epigastric perforator flap breast reconstruction. *Ann Plast Surg*. 2012;69:19–23.
33. Schusterman MA, Kroll SS, Weldon ME. Immediate breast reconstruction: why the free TRAM over the conventional TRAM flap? *Plast Reconstr Surg*. 1992;90:255–261.
34. Hallock GG. Is there a “learning curve” for muscle perforator flaps? *Ann Plast Surg*. 2008;60:146–149.
35. Garvey PB, Buchel EW, Pockaj BA, et al. The deep inferior epigastric perforator flap for breast reconstruction in overweight and obese patients. *Plast Reconstr Surg*. 2005;115:447–457.
36. Holoyda KA, Simpson AM, Ye X, et al. Immediate bilateral breast reconstruction using abdominally based flaps: an analysis of the nationwide inpatient sample database. *J Reconstr Microsurg*. 2019;35:594–601.
37. Banic A, Boeckx W, Greulich M, et al. Late results of breast reconstruction with free TRAM flaps: a prospective multicentric study. *Plast Reconstr Surg*. 1995;95:1195–1204; discussion 1205.
38. Rossetto LA, Ablat LE, Vidal R, et al. Factors associated with hernia and bulge formation at the donor site of the pedicled TRAM flap. *Eur J Plast Surg*. 2010;33:203–208.
39. Shubinetz V, Fox JP, Sarik JR, et al. Surgically treated hernia following abdominally based autologous breast reconstruction: prevalence, outcomes, and expenditures. *Plast Reconstr Surg*. 2016;137:749–757.
40. Wilkins EG, Hamill JB, Kim HM, et al. Complications in postmastectomy breast reconstruction. *Ann Surg*. 2018;267:164–170.
41. DellaCroce FJ, DellaCroce HC, Blum CA, et al. Myth-busting the DIEP flap and an introduction to the abdominal perforator exchange (APEX) breast reconstruction technique: a single-surgeon retrospective review. *Plast Reconstr Surg*. 2019;143:992–1008.