

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Table 2: Carbapenemase Enzymes and Antimicrobial Susceptibilities of CNSC to Novel Beta-Lactam-Beta-Lactamase Agents

Isolate	Organism	Carbapenemase*	Ceftazidime-Avibactam MIC	Meropenem-Vabrobactam MIC
RS102	<i>C. freundii</i>	-	0.5	0.06
RS189	<i>C. freundii</i>	NDM	0.5	0.015
RS226	<i>C. freundii</i>	KPC	<0.25	0.015
RS236	<i>C. freundii</i>	-	2	0.5
RS237	<i>C. freundii</i>	-	64 (R)	0.5
RS240	<i>C. freundii</i>	-	8	0.5
RS259	<i>C. freundii</i>	KPC	>256 (R)	>8 (R)
RS289	<i>C. koseri</i>	-	2	0.12
RS77	<i>C. freundii</i>	KPC	0.5	0.015
YDC582	<i>C. freundii</i>	KPC	>256 (R)	>8 (R)
YDC608	<i>C. freundii</i>	KPC	4	0.06
YDC638-3	<i>C. freundii</i>	KPC	2	0.06
YDC645	<i>C. freundii</i>	KPC	<0.25	0.043
YDC661	<i>C. freundii</i>	KPC	4	0.06
YDC667-1	<i>C. freundii</i>	KPC	0.5	0.03
YDC689-2	<i>C. koseri</i>	-	4	0.12
YDC693	<i>C. freundii</i>	KPC	1	0.03
YDC693-2	<i>C. farmeri</i>	KPC	4	0.12
YDC697-2	<i>C. farmeri</i>	KPC/OXA-48	4	0.12
YDC730	<i>C. werkmanii</i>	-	0.5	0.12
YDC849-1	<i>C. freundii</i>	NDM	>256 (R)	16 (R)

Abbreviations: R: Resistant

*Carbapenemase gene presence was evaluated by multiplex PCR. Genomic DNA from isolates was used in 50 µL reactions containing BlueTaq DNA Polymerase (Denville). Each multiplex PCR reaction contained KPC primers (forward TCGCGCTAGTCTGCTGCTTG and reverse ACAGCTCCGCCACGTCAT), NDM primers (forward ACTGGCTTGTGCTGCTT and reverse CATTAGCCGCTGCTGAT), and OXA-48 primers (forward ATGGCTGATTAGCTTATGCTG and reverse CATCTCTAACCCAGCCCAATC) at 0.5 µM concentration. Cycle parameters were as follows: 1 cycle at 94 °C for 15 min, then 30 cycles of 94 °C for 60 s, 66 °C for 90 s, and 72 °C for 90 s, followed by 1 cycle at 72 °C for 10 min.

Disclosures. All authors: No reported disclosures.

486. Epidemiology of Carbapenem-Resistant *Pseudomonas aeruginosa* Identified through the Emerging Infections Program (EIP), United States, 2016–2018

Julian E. Grass, MPH¹; Sandra N. Bulens, MPH¹; Wendy M. Bamberg, MD²; Sarah J. Janelle, MPH²; Kyle Schutz, MS²; Jesse T. Jacob, MD, MSc³; Chris W. Bower, MPH⁴; Rebekah Blakney, MS⁵; Lucy E. Wilson, MD, ScM⁵; Elisabeth Vaeth, MPH⁶; Linda Li, MPH⁶; Ruth Lynfield, MD⁷; Paula Snippes Vagnone, MT(ASCP)⁸; Ginette Dobbins, MPH⁷; Erin C. Phipps, DVM, MPH²; Emily B. Hancock, MS²; Ghinwa Dumyati, MD¹⁰; Rebecca Tsay, MPH¹⁰; P. Maureen Cassidy, MPH¹¹; Nicole West, MPH¹¹; Marion A. Kainer, MBBS, MPH, FRACP, FSHEA¹²; Jacquelyn Mounsey, RN¹²; Richard A. Stanton, PhD¹; Gillian A. McAllister, BS¹³; Davina Campbell, MS¹; Joseph D. Lutgring, MD¹; Maria Karlsson, PhD¹³ and Maroya S. Walters, PhD¹; ¹Division of Healthcare Quality Promotion, Centers for Disease Control and Prevention, Atlanta, Georgia; ²Colorado Department of Public Health and Environment, Denver, Colorado; ³Emory University, Atlanta, Georgia; ⁴Georgia Emerging Infections Program, Decatur, Georgia; ⁵University of Maryland Baltimore County, Baltimore, Maryland; ⁶Maryland Department of Health, Baltimore, Maryland; ⁷Minnesota Department of Health, Saint Paul, Minnesota; ⁸Minnesota Department of Health Laboratory, St. Paul, Minnesota; ⁹University of New Mexico, Albuquerque, New Mexico; ¹⁰New York Rochester Emerging Infections Program at the University of Rochester Medical Center, Rochester, New York; ¹¹Oregon Health Authority, Portland, Oregon; ¹²Tennessee Department of Health, Nashville, Tennessee; ¹³Centers for Disease Control and Prevention, Atlanta, Georgia

Session: 53. HAI: MDRO – GNR Epidemiology, Other

Thursday, October 3, 2019: 12:15 PM

Background. *Pseudomonas aeruginosa* is intrinsically resistant to many commonly used antimicrobials, and carbapenems are often required to treat infections. We describe the crude incidence, epidemiology, and molecular characteristics of carbapenem-resistant *P. aeruginosa* (CRPA) in the EIP catchment area.

Methods. From August 1, 2016 through July 31, 2018, we conducted laboratory- and population-based surveillance for CRPA in selected areas in eight sites. We defined a case as the first isolate of *P. aeruginosa* resistant to imipenem, meropenem, or doripenem from the lower respiratory tract, urine, wounds, or normally sterile sites identified from a resident of the EIP catchment area in a 30-day period. Patient charts were reviewed. Analysis excluded cystic fibrosis patients. A random sample of isolates was collected. Real-time PCR to detect carbapenemase genes and whole-genome sequencing are in progress.

Results. We identified 4,209 cases in 3373 patients. The annual incidence was 14.50 (95% CI, 14.07–14.94) per 100,000 persons and varied among sites from 4.89 in OR to 25.21 in NY. The median age of patients was 66 years (range: <1–101), 42.1% were female, and nearly all (97.5%) had an underlying condition. Most cases were identified from urine (42.8%) and lower respiratory tract (35.7%) cultures. Nearly all (93.3%) occurred in patients with inpatient healthcare facility stay, surgery, chronic dialysis, or indwelling devices in the prior year; death occurred in 7.2%. Among 937 isolates tested, 847 (90.4%) underwent PCR; six (0.7%) harbored a carbapenemase, from four sites (CO, MD, NY, and OR): bla_{VIM} (3), bla_{KPC} (2), and bla_{IMP} (1). Of 612 (65.3%) isolates sequenced, the most common ST types were ST235 (9.2%) and ST298 (4.9%).

Conclusion. Carbapenemases were rarely the cause of carbapenem resistance but were found at EIP sites with high and low CRPA incidence. The emergence of mobile carbapenemases in *P. aeruginosa* has the potential to increase the incidence of CRPA. Increased detection and early response to carbapenemase-producing CRPA is key to prevent further emergence.

Disclosures. All authors: No reported disclosures.

487. Prevalence of Antimicrobial Resistance in Gram-Negative Bacilli Bloodstream Infections at a Tertiary Teaching Hospital in the Dominican Republic

Alfredo J. Mena Lora, MD¹; Julia Rodriguez Abreu, MD MPH²; Claudia Blanco, MD²; Jacquelin de Lara, MS² and Susan C. Bleasdale, MD¹;

¹University of Illinois at Chicago, Chicago, Illinois; ²CEDIMAT, Santo Domingo, Distrito Nacional, Dominican Republic

Session: 53. HAI: MDRO – GNR Epidemiology, Other

Thursday, October 3, 2019: 12:15 PM

Background. Bloodstream infections (BSI) with gram-negative bacilli (GNB) are a major cause of morbidity and mortality worldwide. Sepsis due to BSI can carry a mortality rate as high as 40%, with higher mortality in developing nations. Early and appropriate empiric therapeutic selection plays an important role in survival. The rising incidence of antimicrobial resistance (AMR) limits empiric treatment options. Local susceptibility patterns can vary per region, institution or setting. Understanding local AMR may help guide empiric treatment choices. We seek to describe resistance rates for GNB BSI in the Dominican Republic (DR).

Methods. This is a retrospective review of antimicrobial susceptibility patterns from bloodstream infections in a tertiary hospital in the DR. Susceptibility data from all adult inpatient blood cultures were collected from January 1 to December 31, 2017.

Results. A total of 124 blood cultures were reported. The most common organisms were *Escherichia coli* (43%) and *Klebsiella pneumoniae* (23%). Fluoroquinolone resistance was present in 70% of *E. coli*. Phenotypic susceptibility patterns consistent with extended-spectrum β-lactamase (ESBL) producing GNB were present in 46% of isolates. Carbapenem resistance was found in 4 samples and was most common in *P. aeruginosa*. Susceptibility profile is described on Table 1.

Conclusion. AMR was high in GNB BSIs in the DR. High rates of ESBL render common cephalosporins sub-optimal for empiric treatment. PTZ retains *in vitro* susceptibilities despite cefepime resistance but clinical efficacy is controversial. CTX-M ESBLs may cause these resistance pattern *in vitro*. Further studies are needed to determine genetic mechanisms of resistance. Establishing antimicrobial stewardship programs with rapid diagnostic testing that identify mechanisms of resistance may promote judicious use of carbapenems and reduce further the risk of further development of AMR.

Table 1. Susceptibility patterns for GNB BSI (%)

Organisms	Ampicillin-Subactam	Cefazolin	Cefepime	Ceftazidime	Ceftioxone	Ciprofloxacin (CQ)	Ertapenem	Gentamicin	Imipenem	Piperacillin-tazobactam (PTZ)
<i>E. coli</i>	47	47	48	48	48	30	100	59	100	97
<i>K. pneumoniae</i>	29	37	37	37	37	52	97	50	97	82
<i>P. aeruginosa</i>	-	-	92	97		87	-	92	79	75
<i>E. cloacae</i>	-	-	90	81	81	81	100	100	100	86
<i>A. baumannii</i>	86	-	86	86		86	-	86	100	100

Disclosures. All authors: No reported disclosures.

488. Epidemiology and Outcomes for *Stenotrophomonas maltophilia* Infections at a Tertiary Care Center in Detroit, MI

Erin Goldman, DO¹; Justin Oring, DO¹; Reda Awali, MD, MPH¹ and Teena Chopra, MD, MPH²; ¹Wayne State University, Oak Park, Michigan; ²Detroit Medical Center, Wayne State University, Detroit, Michigan

Session: 53. HAI: MDRO – GNR Epidemiology, Other

Thursday, October 3, 2019: 12:15 PM

Background. *Stenotrophomonas maltophilia* is a gram-negative, biofilm-forming bacterium. The increasing use of antibiotics has allowed this bacterium to become a predominant nosocomial pathogen with inherent resistance to several antibiotics. In this study, we describe the epidemiology and outcomes for patients treated for *S. maltophilia* infections who were admitted to Detroit Medical Center from January 1, 2010 to August 31, 2018.

Methods. This was a retrospective cohort study that included *S. maltophilia* cultures isolated from sterile body sites from January 1, 2010 to August 31, 2018. Nonsterile body sites and tissue cultures were excluded, as well as cultures that were deemed to be colonization based upon clinical evaluation. Appropriate empiric antibiotic therapy was defined as a regimen administered three days prior to or four days following the *S. maltophilia* culture date. Appropriate definitive therapy was defined as antibiotic treatment administered five to fourteen days following the culture date. Patient data were extracted from the electronic medical record which included demographic information, length of stay and outcome data. Bivariate analysis was performed using SAS database.

Results. 126 patients with *S. maltophilia* infections were analyzed: 89 had bacteremia, 22 had lung infections, and 15 had other infections. The median length of stay was 16 days (IQR 6–30 days). Sixty-one patients (48%) admitted to the ICU had a median length of stay of 10 days (Table 2). Among the patients that were followed after discharge, 21 were readmitted within 30 days. Table 1 highlights the bivariate analysis of patients who died within 30 days vs. survived. Patients who received definitive antibiotic therapy had lower 30-day mortality (Table 1; CI 95%, OR=0.37, P = 0.03). In