

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

INTERSTELLAR COSMIC RAY SPECTRA FROM THE NONTHERMAL RADIO BACKGROUND FROM 0.4 TO 400 MHz

M. L. Goldstein*

University of Maryland, College Park, Maryland 20742

and

R. Ramaty and L. A. Fisk†

National Aeronautics and Space Administration, Goddard Space Flight Center, Greenbelt, Maryland 20771

(Received 6 April 1970)

We deduce the interstellar electron spectrum from the nonthermal radio background. From 200 MeV to a few GeV the spectral index is 1.8 and there is evidence for residual solar modulation. Above a few GeV, the spectrum is steeper and the intensity is similar to that observed at earth. A consistent modulation for electrons and protons of the same rigidity can be obtained by using the complete diffusion-convection energy-loss theory of solar modulation.

Alexander *et al.*^{1,2} have recently reported satellite measurements of the galactic radio background in the range 0.4 to 6.5 MHz. When their measurements in the direction of the galactic anticenter are combined with the corresponding high-frequency data,³ the radio spectrum is consistent with synchrotron emission by cosmic electrons, free-free absorption by ionized hydrogen, and a uniform interstellar mixture of the emitting and absorbing regions. The radio data in both the optically thin and optically thick regimes require that the emissivity from 0.4 MHz to about 100 MHz be of the form $\nu^{-0.4}$. The corresponding electron spectrum from about 200 MeV to several GeV is of the form $E^{-1.8}$. The observations at higher frequencies ($\gtrsim 100$ MHz) can be interpreted as a steepening in the radio spectrum, which suggests that above a few GeV, the electron spectral index is close to 2.5.

It has been suggested (e.g., by Anand, Daniel, and Stephens⁴) that it is possible to determine the residual solar modulation from a comparison of the electron spectrum inferred from radio observations with the spectrum observed at the

earth at solar minimum. However, such a comparison will give a meaningful result at low energies, where modulation is important, only if reliable low-frequency radio data are used.

From a preliminary analysis, Alexander *et al.*² obtained an interstellar electron spectrum around 200 MeV, which, on the average, is greater by a factor of 50 to 100 than that observed at the earth at solar minimum. This finding depends critically on the values assumed for the interstellar parameters, and clearly, a more detailed study is needed before firm conclusions about solar modulation can be made.

In this Letter, we present the results of extensive calculations of synchrotron emission by cosmic electrons, absorption by ionized interstellar hydrogen, and modulation in the interplanetary medium. All the radio data are consistent with an interstellar electron intensity that is significantly smaller than that given in Ref. 2. The deduced modulation agrees with measured values of the solar wind speed and the particle diffusion coefficient in interplanetary space.

The synchrotron emissivity produced by an arbitrary electron density $N(E)$ is given by⁵

$$\epsilon(\nu) = [\sqrt{3} e^3 H / 4\pi m c^2] \int_{mc^2}^{\infty} \{ (\nu/\nu_c) \int_{\nu/\nu_c}^{\infty} K_{5/3}(y) dy \} N(E) dE, \quad (1)$$

where $\nu_c = [3eH/4\pi mc] (E/mc^2)^2$ and H is the mean value of the magnetic field perpendicular to the line of sight. Possible values of H in the interstellar medium range from about 3 to 5 μG .⁶ The values we have chosen for interstellar parameters enable us to deduce a minimal interstellar cosmic electron intensity and, hence, a plausible lower limit to the solar modulation. Consequently, we used $H = 5 \mu\text{G}$.

The free-free absorption coefficient in ionized hydrogen is given by⁷

$$K(\nu) = (10^{-2} n_e^2 / \nu^2 T_e^{3/2}) [17.7 + \ln(T_e^{3/2} / \nu)], \quad (2)$$

where ν is measured in Hz, and n_e and T_e are the density and temperature of the ambient electrons. Recent studies of interstellar heating^{8,9} have indicated that a two-component model consisting of cold clouds and a hot intercloud medium is required to characterize the interstellar gas. We have used an

ambient electron density and temperature in cold clouds of 0.05 cm^{-3} and 60°K , respectively, consistent with observations of 21-cm absorption.¹⁰ We also used an intercloud electron density of 0.03 cm^{-3} , consistent with pulsar dispersion measurements.¹¹ There is no direct observational information on the temperature T_i of the intercloud medium. Because theoretical investigations indicate values ranging from 500 to 10^4°K , we assumed that the intercloud temperature is a free parameter within this range.

The average fraction of a line of sight toward the anticenter intercepted by cold clouds is about 10^{-3} to 10^{-2} .^{8,9} The radio spectrum at low frequencies is consistent with absorption by a diffuse medium and does not exhibit an exponential falloff characteristic of a compact absorber between the observer and the emitting region. We used a uniform distribution of cold clouds with a cloud diameter of 1 pc and intercloud separation of 1 kpc as a model. Furthermore, we assumed that the closest cloud is at a distance of 1 kpc, so that the radio intensity from the anticenter is given by

$$I = (\epsilon/K)[1 - \exp(-\tau_i)] \{1 - \exp[-L(\tau_c + \tau_i)]\} [1 - \exp(-\tau_c - \tau_i)]^{-1}, \quad (3)$$

where τ_c and τ_i are the optical depths corresponding to one cloud and one intercloud separation, respectively, and L is the size of the emitting region in kiloparsecs.

We evaluated Eqs. (1)-(3) for several values of T_i and L . For $T_i = 10^4^\circ$, K is small and a value of L of about 10 kpc is needed to account for all the absorption at low frequencies. But for such a large value of L , even for the minimal conditions we have used, absorption by cold clouds becomes significant below a few MHz, so that the theoretical spectrum no longer fits the data. For $T_i < 10^3^\circ$, K is large, and to fit the data, L must be less than about 1 kpc. Such a small value of L means that radiation is collected only over a small fraction of the available line-of-sight distance toward the anticenter. At several GeV, this small value of L requires an average electron flux that is ≥ 10 times larger than is observed near the earth. Solar modulation is probably negligible at these energies and hence the difference must be due to interstellar fluctuations in the cosmic electron intensity. Because fluctuations of this magnitude over a scale smaller than 1 kpc seem unlikely, values of $T_i < 10^3^\circ \text{K}$ appear untenable.

We achieved a good fit to the radio data with $T_i \approx 4000^\circ\text{K}$, which requires $L \approx 4 \text{ kpc}$. The computed radio spectra for these parameters are shown in Fig. 1, with the observations from the anticenter. Curves A, B, and C correspond to the assumed interstellar electron spectra which are shown in Fig. 2 with the electron observations at earth near solar minimum.

The spectral index for these spectra is 1.8, increasing to 2.5 above E_c , where E_c is 1, 2, and 5 GeV for A, B, and C, respectively. Because synchrotron and Compton energy losses in interstellar space are not expected to have a major effect on the spectrum below a few GeV, the elec-

tron-injection spectral index is also close to 1.8. This is consistent with observed spectral indices of radio emission from galactic nonthermal sources. These indices are centered on a value of 0.5 with an rms scatter of about 0.3 and show little evidence of an evolutionary trend.¹³ Within the framework of synchrotron theory, this means that the electron spectra in the supernova remnants have an index of 2.0 ± 0.6 , and could very well be representative of the initial injection spectrum.

Above E_c , solutions A and B agree with the measured electron intensity at the earth, and solution C requires an average intensity greater only by a factor of $\lesssim 4$. Within the uncertainties of the cosmic ray measurements and the interstellar parameters, the observed electron intensity at a few GeV does not differ significantly from that required to produce the radio emission. Hence, interstellar fluctuations do not play a major role at these energies.¹⁴




FIG. 1. Nonthermal radio background spectrum from the galactic anticenter (Refs. 2 and 3). Curves A, B, and C were computed using Eq. (3) with $T_i = 4000^\circ\text{K}$, $L = 4 \text{ kpc}$, and the electron spectra A, B, and C shown in Fig. 2.




FIG. 2. Electron observations (Ref. 12) at the earth from 0.1 to 10 GeV. Curves A, B, C are the interstellar electron spectra required to produce the radio spectra A, B, and C in Fig. 1, and are normalized to produce the same radio intensity at low frequencies. The modulated electron intensity was obtained from Curve B, using the complete diffusion-convection energy-loss theory of solar modulation.

Below E_c , all solutions require a finite solar modulation. For example, at 200 MeV, our modulating factor is about 10, which is somewhat below the lower limit obtained by Alexander et al.² However, their mean modulating factor of about 50 to 100 results from using an intercloud temperature of 10^3 °K which, as discussed above, leads to a fluctuation of ~ 10 in the cosmic electron intensity over a distance of less than 1 kpc. It appears to us that our solution with $T_i \approx 4000$ °K gives a more reasonable cosmic electron spectrum in terms of both interstellar propagation and interplanetary modulation.

Modulation has frequently been discussed in terms of a function of the form $M = \exp[\eta/f(R)\beta]$. R and β are rigidity and velocity, respectively, and

$$\eta = \int_{r_e}^{r_b} dr V/D(r),$$

where V is the speed of the solar wind, $D(r)$ is the radial dependence of the particle diffusion coefficient, and r_e and r_b are the heliocentric




FIG. 3. Proton observations (Ref. 17) at the earth from 20 MeV to 10 GeV. The modulated intensity is obtained from the interstellar spectra, with the same solar wind speed and diffusion coefficient used in Fig. 2. When adiabatic deceleration is included, the interstellar proton spectrum is approximately a power law in total energy.

distances to the earth and to the boundary of the modulating region, respectively. The results shown in Fig. 2 can be interpreted in terms of such a modulating function with $\eta \approx 0.64$ GV, and $f(R) = R$ for $R \geq R_0 = 350$ MV, and $f(R) = (RR_0)^{1/2}$ for $R < R_0$. When this modulating function is applied to protons, $M_p(R) = [M_e(R)]^{1/\beta}$, where p and e denote protons and electrons, respectively. In particular, $M_e = 10$ at 200 MeV implies that $M_p \approx 5 \times 10^4$ at ~ 20 MeV. This leads to a very large cosmic-ray energy density in interstellar space and is probably in conflict with the dynamics of the interstellar medium.⁶

A more reasonable fit for the modulation of both electrons and protons can be obtained with a detailed theory of modulation that includes the effects of adiabatic deceleration.^{15, 16} In Figs. 2 and 3, we present the results of numerical solutions of the Fokker-Planck equation that describe the modulation of electrons and protons in this

theory.¹⁸ The functional form of $f(R)$ that we used is the same as that given above, the solar wind speed is 400 km/sec, and $D(r) = 1.5 \times 10^{21} \times \exp[0.625(r-1)]$ cm²/(sec BV), where r is in A. U. These values of D and $f(R)$ are consistent with the diffusion coefficient determined from the observed radial gradient of high-energy protons¹⁹ and the measured power spectra of magnetic fluctuations.²⁰

The effects of adiabatic deceleration are most important when the modulation is large and the interplanetary spectrum is increasing with increasing energy. In this case, energy losses diminish the effects of diffusion and simple convection. Consequently, adiabatic deceleration is extremely significant for protons below ~200 MeV, but does not play an important part in the modulation of electrons. Therefore, the proton spectrum suffers a much smaller modulation than that deduced from the simple formula $M_p(R) = [M_e(R)]^{1/\beta}$, and thus there is no inconsistency with the dynamics of the interstellar medium.

The residual electron modulation can also be estimated from the comparison of the measured positron intensity and the interstellar positron flux produced by the π - μ - e process.^{21,22} The measured ratio $e^+/(e^+ + e^-)$ at the earth at 200 MeV is about 0.1.^{22,23} The positron intensity in interstellar space is $\sim 2.5 \times 10^2$ (m² sec sr GeV)⁻¹,²¹ which, when combined with the demodulated spectrum shown in Fig. 2, also gives $e^+/(e^+ + e^-) \sim 0.1$.

We conclude that the radio data between 0.4 and 400 MHz can be explained by an electron spectrum as shown in Fig. 2. Most of the absorption at low frequencies results from the intercloud medium at $\sim 4000^\circ\text{K}$. The cosmic electron spectrum is consistent with injection spectra inferred from radio observations of supernova remnants as well as with the ratio $e^+/(e^+ + e^-)$. The deduced electron modulation can be applied to cosmic protons of the same rigidity only if the complete diffusion-convection energy-loss theory is used. The resulting interstellar proton intensity is approximately a power law in total energy.

*Research supported by NASA Grant No. NGL 21-002-033.

†NAS-NASA Resident Research Associate.

¹J. K. Alexander, L. W. Brown, T. A. Clark, R. G. Stone, and R. R. Weber, *Astrophys. J. Letters* **157**, L163 (1969).

²J. K. Alexander, L. W. Brown, T. A. Clark, and

R. G. Stone, to be published.

³W. R. Webber, *Australian J. Phys.* **21**, 845 (1968).

⁴K. C. Anand, R. R. Daniel, and S. A. Stephens, *Nature* **217**, 25 (1968).

⁵V. L. Ginzburg and S. I. Syrovatskii, *The Origin of Cosmic Rays* (Pergamon, Oxford, England, 1964).

⁶E. N. Parker, *Space Sci. Rev.* **9**, 651 (1969).

⁷V. L. Ginzburg, *The Propagation of Electromagnetic Waves in Plasmas* (Pergamon, Oxford, England, 1964).

⁸G. B. Field, D. W. Goldsmith, and H. J. Habing, *Astrophys. J.* **155**, L149 (1969).

⁹R. M. Hjellming, C. P. Gordon, and K. J. Gordon, *Astron. Astrophys.* **2**, 202 (1969).

¹⁰U. Mebold, *Beitr. Radioastron.* **1**, 97 (1969).

¹¹A. H. Bridle and V. R. Venugopal, *Nature* **224**, 545 (1969).

¹²W. R. Webber and C. Chotkowski, *J. Geophys. Res.* **72**, 2783 (1967); J. L'Heureux and P. Meyer, *Can. J. Phys.* **46**, S892 (1968); J. A. M. Bleeker, J. J. Burger, A. J. M. Deerenberg, A. Scheepmaker, B. N. Swanenburg, and Y. Tanaka, *Can. J. Phys.* **46**, S522 (1968); J. L. Faselow, R. C. Hartman, R. H. Hildebrand, and P. Meyer, *Astrophys. J.* **158**, 771 (1969); J. A. Earl, D. E. Neely, and T. A. Rygg, Univ. of Maryland Technical Report No. 70-076, 1970 (unpublished).

¹³M. J. L. Kesteven, *Australian J. Phys.* **21**, 739 (1968).

¹⁴According to J. K. Alexander (private communication) the extragalactic background is not completely negligible at about 10 MHz in the direction of the anti-center. Thus, the deduced electron spectra above ~ 1 GeV may be somewhat lower, further substantiating our conclusion that there is no evidence for interstellar fluctuations of cosmic electrons at these energies.

¹⁵E. N. Parker, *Planet. Space Sci.* **13**, 9 (1965).

¹⁶L. J. Gleeson and W. I. Axford, *Astrophys. J.* **154**, 1011 (1968).

¹⁷C. Y. Fan, G. Gloeckler, and J. A. Simpson, *Phys. Rev. Letters* **17**, 329 (1966); V. K. Balasubrahmanyam, D. E. Hagge, G. H. Ludwig, and F. B. McDonald, in *Proceedings of the Ninth International Conference on Cosmic Rays, London, England, 1965* (Institute of Physics and the Physical Society, London, 1966), Vol. 1, p. 427, and *J. Geophys. Res.* **71**, 1771 (1966); C. J. Waddington and P. S. Freier, in *Proceedings of the Ninth International Conference on Cosmic Rays, London, England, 1965* (Institute of Physics and the Physical Society, London, 1966), Vol. 1, p. 339; J. F. Ormes and W. R. Webber, *ibid.*, p. 349; F. B. McDonald, *Phys. Rev. Letters* **109**, 1367 (1958).

¹⁸L. A. Fisk, dissertation, Univ. of California, 1969 (unpublished).

¹⁹L. J. Gleeson and W. I. Axford, *Can. J. Phys.* **46**, S937 (1968).

²⁰J. W. Sari and N. F. Ness, *Solar Phys.* **8**, 155 (1969).

²¹R. Ramaty and R. E. Lingenfelter, *Phys. Rev. Letters* **20**, 120 (1968).

²²K. P. Beuermann, J. Rice, E. C. Stone, and R. E. Vogt, *Phys. Rev. Letters* **22**, 412 (1969).

²³See Faselow, Hartman, Hildebrand, and Meyer, Ref. 12.