

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

**Please provide feedback**

Please support the ScholarWorks@UMBC repository by emailing [scholarworks-group@umbc.edu](mailto:scholarworks-group@umbc.edu) and telling us what having access to this work means to you and why it's important to you. Thank you.

EGU23-2131, updated on 03 Apr 2023

<https://doi.org/10.5194/egusphere-egu23-2131>

EGU General Assembly 2023

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



## Sub-seasonal variability of Asian summer monsoon transport of aerosols and CO to the UTLS in the context of recent aircraft observations in the Asia

Mian Chin<sup>1</sup>, Huisheng Bian<sup>2</sup>, Peter Colarco<sup>1</sup>, Leslie Lait<sup>3</sup>, and Gao Chen<sup>4</sup>

<sup>1</sup>NASA Goddard Space Flight Center, Code 614, Greenbelt, United States of America (mian.chin@nasa.gov)

<sup>2</sup>University of Maryland Baltimore County

<sup>3</sup>SSAI

<sup>4</sup>NASA Langley Research Center

We present our study on the sub-seasonal variability of UTLS aerosols and CO that is a result of the variability induced by the sub-seasonal variability of the Asian summer monsoon dynamics. We use the NASA global model GEOS simulations that incorporates emissions from anthropogenic, biomass burning, volcanic, and other natural sources to simulate CO, aerosols and related gases and the model experiments separating source types (anthropogenic, biomass burning, volcanic) and source locations (East Asia, South Asia). With model results and observations from recent aircraft measurements (StratoClim, ACCLIP) in the UTLS over the Asian summer monsoon regions, we will discuss (1) the sub-seasonal variability of transport pathways of surface-generated pollutants to reach UTLS, and (2) sub-seasonal variation of aerosol composition that is determined by the variability of source type originating in different locations.