

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

MIRRORING IN THE FOKKER-PLANCK COEFFICIENT FOR COSMIC-RAY PITCH-ANGLE SCATTERING IN HOMOGENEOUS MAGNETIC TURBULENCE

M. L. GOLDSTEIN* AND A. J. KLIMAS†

Laboratory for Extraterrestrial Physics, NASA-Goddard Space Flight Center

AND

G. SANDRI

Aeronautical Research Associates of Princeton, Inc.

Received 1974 May 3; revised 1974 August 20

ABSTRACT

The Fokker-Planck coefficient for pitch-angle scattering, appropriate for cosmic rays in homogeneous, stationary magnetic turbulence, is computed from first principles. No assumptions are made concerning any special statistical symmetries the random field may have. This result can be used to compute the parallel diffusion coefficient for high-energy cosmic rays moving in strong turbulence, or low-energy cosmic rays moving in weak turbulence. Because of the generality of the magnetic turbulence which is allowed in this calculation, special interplanetary magnetic field features, such as discontinuities or particular wave modes, can be included rigorously. The reduction of this result to previously available expressions for the pitch-angle scattering coefficient in random field models with special symmetries is discussed.

The general existence of a Dirac δ -function in the pitch-angle scattering coefficient is demonstrated. It is proved in this paper that this delta function is the prediction of the Fokker-Planck equation for pitch-angle scattering due to mirroring in the magnetic field. The conditions under which this δ -function contributes to pitch-angle scattering are determined, and shown to be identical to the conditions under which first-order mirroring occurs in the random field. These conditions are generally fulfilled in interplanetary and probably interstellar space. The implications of the δ -function for the validity of the Fokker-Planck equation are discussed.

Subject headings: cosmic rays — hydromagnetics — plasmas

I. INTRODUCTION

The parallel diffusion coefficient for low-energy cosmic rays in a random magnetic field has often been calculated from the small-gyroradius (or guiding center) approximation to the Fokker-Planck pitch-angle scattering coefficient (Jokipii 1966, 1967, 1968, 1971; Hall and Sturrock 1967; Hasselmann and Wibberenz 1968). But Klimas and Sandri (1973b), in a numerical calculation using the exact pitch-angle scattering coefficient, have computed a parallel diffusion coefficient which differs markedly from that predicted using the small gyroradius approximation. This calculation was limited to the special case of statistically isotropic magnetic turbulence with a Gaussian correlation function (and thus, a Gaussian power spectrum). Because of this limitation, the significance of this discrepancy was not fully appreciated because (1) in any case the small-gyroradius approximation was not expected to be accurate for Gaussian power spectra (Jokipii 1971) but was still proposed as an accurate approximation for the more familiar power-law, power spectra, and (2) the source of the discrepancy, and especially its physical interpretation, was not discernible in the numerical computation.

Recently, Fisk *et al.* (1974) and Klimas and Sandri (1973c), still within the framework of isotropic turbulence but for power-law power spectra, calculated the pitch-angle scattering coefficient without making the small-gyroradius approximation. They found that the exact result differed significantly from the approximate one, especially for pitch angles θ (measured relative to the mean field) near 90° where the Fokker-Planck coefficient contains a Dirac δ -function in $\mu = \cos \theta$. The discrepancy found by Klimas and Sandri was then completely explained. For low-energy cosmic rays in the Gaussian power spectrum, the δ -function dominated all other contributions to the parallel diffusion coefficient. Furthermore, Klimas and Sandri (1973c) in calculating the parallel diffusion coefficient showed that, for power-law spectra with any reasonable spectral index, the contribution of the δ -function still dominated the contribution of the small-gyroradius approximation. For these spectra, Fisk *et al.* (1974) through a numerical calculation found that, for $\mu \neq 0$, the pitch-angle scattering coefficient is overestimated by the small-gyroradius approximation. Thus, the δ -function was apparently the sole contributor to the parallel diffusion coefficient in realistic models of the interplanetary magnetic field.

The above conclusions concerning the importance of the δ -function depend on the particular expansion of the cosmic-ray distribution function used by Klimas and Sandri. Recently, Earl (1974) has shown that a unique relationship exists between the pitch-angle scattering coefficient and the parallel diffusion coefficient. In this

* NAS-NRC Postdoctoral Resident Research Associate.

† NAS-NRC Senior Postdoctoral Resident Research Associate.

formulation of the parallel diffusion coefficient, however, the contribution of the δ -function is not defined. Consequently, a physical interpretation of the δ -function, as well as a reexamination of the validity of the Fokker-Planck equation in light of its existence, has become necessary.

Concurrent with the above developments, a number of papers (Jones, Birmingham, and Kaiser 1973; Kaiser, Jones, and Birmingham 1973; Jones, Kaiser, and Birmingham 1973*a, b*; Kaiser 1973; Jones and Birmingham 1974; Volk 1973; Volk *et al.* 1974) were presented in which it was argued that the quasi-linear approximation used to derive the Fokker-Planck equation fails to correctly describe pitch-angle scattering near $\theta = 90^\circ$ ($\mu = 0$). For mathematical simplicity, these papers have invariably discussed an idealized model of the random magnetic field in which the $\delta(\mu)$ does not appear. We will demonstrate in this paper that the δ -function is the rule, rather than the exception, and must therefore be taken into account in studying the apparent failure of the quasi-linear approximation near $\mu = 0$.

In this paper we derive the Fokker-Planck pitch-angle scattering coefficient with no restrictions on the statistical behavior of the random magnetic field, other than the assumption that it is homogeneous and stationary over several correlation lengths. Our calculation is based on the Liouville equation for the cosmic-ray distribution function. We apply the quasi-linear and adiabatic approximations which, together, give us a Fokker-Planck equation for the *mean* cosmic-ray distribution function. The mean distribution function is assumed gyrotropic in order to focus our attention on pitch-angle scattering. This derivation is done in a way which clearly demonstrates the existence of the $\delta(\mu)$ contribution to pitch-angle scattering. But the generality of our derivation enables us to study the conditions under which the strength of the δ -function is zero or nonzero. In this way, we are able to determine that the δ -function part of the pitch-angle scattering coefficient is the prediction of the Fokker-Planck equation for the contribution of mirroring (Alfvén and Falthammar 1963; Northrop 1963) to pitch-angle scattering. It has been thought that the effects of mirroring were not contained, *a priori*, in the Fokker-Planck equation but had to be developed separately. Work along these lines has been done by Noerdlinger (1968), Quenby *et al.* (1970), Cesarsky and Kulsrud (1972), and Jokipii (1973).

In general, the interplanetary or interstellar magnetic field will have fluctuations in field magnitude which will produce particle mirroring to first order in the weak random field strength. Further efforts at constructing a correct description of the behavior of those cosmic-ray particles with pitch angles near 90° should not be based on idealized models of the random magnetic field which do not allow first-order mirroring.

II. PITCH-ANGLE SCATTERING COEFFICIENT: ARBITRARY HOMOGENEOUS TURBULENCE

We have been able to construct the Fokker-Planck pitch-angle scattering coefficient using two independent methods. The first method, which we will present here, follows directly from first principles, i.e., the Liouville equation for the cosmic-ray particle distribution function. The second method, outlined in Appendix A, is based on the standard approach (Chandrasekhar 1943; Hasselman and Wibberenz 1968; Jokipii 1971). Although these methods are similar in some respects, they also contain surprisingly dissimilar reasoning. The results of both methods are identical.

Klimas and Sandri (1973*a*) have shown that, starting from the Liouville equation for the cosmic-ray distribution function, the following *truncated master equation* for the *mean* cosmic-ray distribution function can be constructed:

$$\frac{\partial f}{\partial \tau} + \alpha \mathcal{K} f + \epsilon \mathcal{L} f = (\epsilon')^2 \langle \mathcal{L}' G_0 \mathcal{L}' \rangle f. \quad (1)$$

The right-hand side of this equation represents the leading significant term in an infinite series expansion in the small parameter, ϵ' . By truncating this expansion at this point, we in effect make the familiar quasi-linear approximation. As a consequence of this truncation, we approximate the actual trajectory of a particle in the magnetic field by the helical trajectory it would have in the mean magnetic field. We assume this helical trajectory for however long it takes the particle to travel approximately one correlation length along the mean field. Particles with pitch angles near 90° may take a long time to move this distance, however, and in this time the assumed helical approximation to the actual particle motion becomes suspect. It is this point which has led to recent modifications of the quasi-linear approximation in the region of particle phase-space near, $\mu = 0$. Our point of view here is to determine the actual predictions of the quasi-linear theory, which we may then use, in confidence, to further investigate the apparent failure of this theory and its possible modifications.

Equation (1) has been written in a dimensionless form. The time variable is given by $\tau = t(v/\lambda_c)$, where t is the dimensional time, v is the particle speed, and λ_c is the correlation length in the random field. The term $\alpha \mathcal{K} f$ is nonzero only if the distribution function depends on spatial position. Our purpose here is to fully investigate the pitch-angle scattering coefficient that this theory predicts. Consequently, we can drop the term $\alpha \mathcal{K} f$ by assuming f independent of position.

The parameters ϵ and ϵ' are defined by

$$\epsilon = \frac{\lambda_c \langle B \rangle}{P}, \quad \epsilon' = \epsilon \eta = \epsilon \left(\frac{B'_{\text{rms}}}{\langle B \rangle} \right), \quad (2)$$

where $\langle B \rangle$ and B'_{rms} are the mean and the rms magnetic field strengths, respectively, and where P is the particle rigidity. We note that Klimas and Sandri (1973a) have given a rescaling of equation (1) which is necessary when $\epsilon \geq 1$. The version given here is appropriate when $\epsilon \leq 1$. However, the results of the calculation we will present here can be applied to any range in ϵ , as long as we remember that we must have $\epsilon' \ll 1$ when $\epsilon \leq 1$, but when $\epsilon \geq 1$ we must have $\eta \ll 1$ instead.

The differential operators \mathcal{L} and \mathcal{L}' generate the effects on the distribution function of the Lorentz force on a charged particle in the mean and random components of the magnetic field, respectively. Convenient representations of \mathcal{L} and \mathcal{L}' can be given in terms of the spherical coordinate system variables: θ , the polar or pitch angle between the particle momentum and the direction of the mean magnetic field; and ϕ , the azimuth or phase angle of the momentum vector. With the definition of $\mu \equiv \cos \theta$, we are able to express these operators as

$$\mathcal{L} = -\frac{\partial}{\partial \phi} \quad (3)$$

and

$$\mathcal{L}' = \frac{\partial}{\partial \mu} (\hat{p} \cdot \Omega \cdot \beta') - \frac{\partial}{\partial \phi} (\beta' \cdot n \cdot \hat{\beta}) (1 - \mu^2)^{-1}, \quad (4a)$$

or

$$\mathcal{L}' = -(\beta' \cdot \Omega \cdot \hat{p}) \frac{\partial}{\partial \mu} + (1 - \mu^2)^{-1} (\beta' \cdot n \cdot \hat{\beta}) \frac{\partial}{\partial \phi}, \quad (4b)$$

where \hat{p} is a unit vector in the momentum direction, $\hat{\beta}$ is a unit vector in the direction of the mean field, $\beta' = B'/B'_{\text{rms}}$, and the tensors Ω and n are defined by

$$\Omega_{ij} = \epsilon_{ijk} \beta_k \quad (5)$$

and

$$n_{ij} = \delta_{ij} - p_i p_j / p^2. \quad (6)$$

A full description of the integral operator, G_0 , has been given by Klimas and Sandri (1973a). This operator, when operating on an arbitrary function of position, momentum, and time, operates as follows:

$$GA(x, p, \tau) = \int_0^\tau d\lambda \mathcal{S}_0(\epsilon, \lambda) A(x, p, \tau - \lambda) = \int_0^\tau d\lambda A(x(\lambda), p(\lambda), \tau - \lambda). \quad (7a, b)$$

The streaming operator $\mathcal{S}_0(\epsilon, \lambda)$ in equation (7a) shifts the phase-space position (x, p) to $[x(\lambda), p(\lambda)]$ in equation (7b) which is along the helical particle trajectory in the mean magnetic field with (x, p) for the starting point; i.e.,

$$x(\lambda) = x - r(\lambda) \quad (8)$$

where

$$r(\lambda) = \mathcal{D}(\epsilon, \lambda) \cdot \hat{p} \quad (9)$$

and

$$\hat{p}(\lambda) = \mathcal{C}(\epsilon, \lambda) \cdot \hat{p}, \quad (10)$$

where

$$\mathcal{C}(\epsilon, \lambda) = P + N \cos \epsilon \lambda - \Omega \sin \epsilon \lambda \quad (11)$$

and

$$\mathcal{D}(\epsilon, \lambda) = P\lambda + \frac{1}{\epsilon} [N \sin \epsilon \lambda + \Omega (\cos \epsilon \lambda - 1)]. \quad (12)$$

The skew-symmetric tensor Ω is defined above in equation (5), and the parallel and normal projection operators, P and N , are defined through

$$P_{ij} = \beta_i \beta_j \quad (13)$$

and

$$N_{ij} = \delta_{ij} - P_{ij}. \quad (14)$$

Now, we can rewrite equation (1) as

$$\begin{aligned} \frac{\partial f}{\partial \tau} - \epsilon \frac{\partial f}{\partial \phi} = (\epsilon')^2 & \left\langle \left[\frac{\partial}{\partial \mu} (\hat{p} \cdot \Omega \cdot \beta') - \frac{\partial}{\partial \phi} (\beta' \cdot n \cdot \hat{\beta}) (1 - \mu^2)^{-1} \right] \right. \\ & \times \left. \int_0^\tau d\lambda \mathcal{S}_0(\epsilon, \lambda) \left[-(\beta' \cdot \Omega \cdot \hat{p}) \frac{\partial}{\partial \mu} + (1 - \mu^2)^{-1} (\beta' \cdot n \cdot \hat{\beta}) \frac{\partial}{\partial \phi} \right] \right\rangle f(\mu, \phi, \tau - \lambda). \end{aligned} \quad (15)$$

The integro-differential operator on the right side of equation (15) is quite complicated, as it acts on an arbitrary

$f(\mu, \phi, \tau - \lambda)$. For the purpose of computing pitch-angle scattering only, we assume a gyrotropic distribution function; i.e., we assume f independent of ϕ . Then, after averaging equation (15) over ϕ , we obtain

$$\frac{\partial f}{\partial \tau} = (\epsilon')^2 \frac{1}{2} \frac{\partial}{\partial \mu} D_\mu(\tau) * \frac{\partial f}{\partial \mu}, \quad (16)$$

where

$$D_\mu(\tau) = -\frac{1}{\pi} \int_0^\tau d\lambda \int_0^{2\pi} d\phi \langle (\hat{\mathbf{p}} \cdot \boldsymbol{\Omega} \cdot \boldsymbol{\beta}'(\mathbf{x})) \mathcal{S}_0(\epsilon, \lambda) (\boldsymbol{\beta}'(\mathbf{x}) \cdot \boldsymbol{\Omega} \cdot \hat{\mathbf{p}}) \rangle \quad (17)$$

and where the star (*) in equation (16) indicates the convolution integral operation on the function of time to the right. Notice that this term, because it is assumed to be a function of μ and τ only, is not affected by the streaming operator in $D_\mu(\tau)$.

We proceed by introducing the Fourier integral transform representation of $\boldsymbol{\beta}'(\mathbf{x})$, and then allowing the streaming operator to operate on the resulting explicit dependence on the phase-space variables. We obtain

$$D_\mu(\tau) = -\frac{1}{\pi} \int_0^\tau d\lambda \int_0^{2\pi} d\phi \left(\frac{1}{2\pi} \right)^3 \int d^3k \int d^3k' \exp(i\mathbf{k} \cdot \mathbf{x}) \exp\{i\mathbf{k}' \cdot [\mathbf{x} - \mathbf{r}(\lambda)]\} [\hat{\mathbf{p}} \cdot \boldsymbol{\Omega} \cdot \langle \boldsymbol{\beta}'(\mathbf{k}) \boldsymbol{\beta}'(\mathbf{k}') \rangle \cdot \boldsymbol{\Omega} \cdot \mathcal{C}(\lambda) \cdot \hat{\mathbf{p}}] \quad (18)$$

With the assumption of homogeneous magnetic turbulence we can introduce

$$\langle \boldsymbol{\beta}'(\mathbf{k}) \boldsymbol{\beta}'(\mathbf{k}') \rangle = (2\pi)^{3/2} \mathbf{R}(\mathbf{k}) \delta(\mathbf{k} + \mathbf{k}'), \quad (19)$$

where $\mathbf{R}(\mathbf{k})$ is the Fourier integral transform of the correlation tensor,

$$\mathbf{R}(\mathbf{r}) = \langle \boldsymbol{\beta}'(\mathbf{x}) \boldsymbol{\beta}'(\mathbf{x} + \mathbf{r}) \rangle. \quad (20)$$

No other assumptions concerning $\mathbf{R}(\mathbf{k})$, or $\mathbf{R}(\mathbf{r})$, will be made.

Through the assumption of homogeneity we are able to rewrite the exponential terms appearing in equation (18) as

$$\exp(i\mathbf{k} \cdot \mathbf{x}) \exp\{i\mathbf{k}' \cdot [\mathbf{x} - \mathbf{r}(\lambda)]\} = \exp[-i\mathbf{k} \cdot \mathbf{r}(\lambda)], \quad (21)$$

where $\mathbf{r}(\lambda)$ is given by equations (9) and (12). This result can be further rewritten to bring out the explicit dependence on the phase angle ϕ . Then the averaging over phase can be done. We will proceed in a particular coordinate system. Since $D_\mu(\tau)$ is a scalar quantity, our choice of coordinate system is irrelevant. The end result of this calculation will, however, be written in an invariant form which then will apply in any coordinate system.

In a Cartesian coordinate system denoted by (1, 2, 3), we let the mean field be in the 3-direction. Then, both $\hat{\mathbf{p}}$ and \mathbf{k} have parallel and perpendicular components denoted, for example, by $(\hat{\mathbf{p}})_\parallel \equiv \mathbf{P} \cdot \hat{\mathbf{p}}$ and $(\hat{\mathbf{p}})_\perp \equiv \mathbf{N} \cdot \hat{\mathbf{p}}$ with similar definitions for \mathbf{k}_\parallel and \mathbf{k}_\perp . The magnitudes of these vector components will be denoted by $|(\hat{\mathbf{p}})_\parallel| = \mu$ and $|(\hat{\mathbf{p}})_\perp| = \mu' = (1 - \mu^2)^{1/2}$, and by k_\parallel and k_\perp . The Cartesian components of these vectors are defined through

$$\hat{p}_1 = \mu' \cos \phi, \quad \hat{p}_2 = \mu' \sin \phi, \quad \hat{p}_3 = \mu, \quad \text{and} \quad k_1 = k_\perp \cos \psi, \quad k_2 = k_\perp \sin \psi, \quad k_3 = k_\parallel. \quad (22)$$

Now, we can rewrite equation (21) as follows:

$$\begin{aligned} \exp[-i\mathbf{k} \cdot \mathbf{r}(\lambda)] &= \exp[-i\mathbf{k}_\parallel \cdot \mathbf{r}_\parallel(\lambda)] \exp[-i\mathbf{k}_\perp \cdot \mathbf{r}_\perp(\lambda)] \\ &= \exp(-ik_\parallel \mu \lambda) \sum_{m,n=-\infty}^{\infty} \exp[i(m-n)(\psi - \phi)] \exp(-i\epsilon m \lambda) J_m\left(\frac{k_\perp \mu'}{\epsilon}\right) J_n\left(\frac{k_\perp \mu'}{\epsilon}\right), \end{aligned} \quad (23)$$

where the J 's are the Bessel functions of the first kind. On substituting equations (19) and (23) into equation (18), we find

$$\begin{aligned} D_\mu(\tau) &= -\frac{(2\pi)^{-3/2}}{\pi} \int_0^\tau d\lambda \int_0^{2\pi} d\phi \int d^3k \exp(-ik_\parallel \mu \lambda) \sum_{m,n=-\infty}^{\infty} e^{i(m-n)(\psi - \phi)} \\ &\quad \times e^{-i\epsilon m \lambda} J_m\left(\frac{k_\perp \mu'}{\epsilon}\right) J_n\left(\frac{k_\perp \mu'}{\epsilon}\right) [\hat{\mathbf{p}} \cdot \boldsymbol{\Omega} \cdot \mathbf{R}(\mathbf{k}) \cdot \boldsymbol{\Omega} \cdot \mathcal{C}(\lambda) \cdot \hat{\mathbf{p}}]. \end{aligned} \quad (24)$$

Then, by introducing the tensor elements of $\mathbf{R}(\mathbf{k})$, denoted by $R_{11}(\mathbf{k})$, $R_{22}(\mathbf{k})$, etc., we are able to exhibit explicitly

the phase-angle dependence of the quantity in square brackets in equation (24). We find

$$[\] = \frac{1}{2}(1 - \mu^2)[-(R_{11} + R_{22})\cos \epsilon\lambda + (R_{11} - R_{22})\cos(\epsilon\lambda + 2\phi) + (R_{12} + R_{21})\sin(\epsilon\lambda + 2\phi) - (R_{12} - R_{21})\sin \epsilon\lambda]. \quad (25)$$

From our earlier assumption of homogeneity, we find

$$R_{ij}(\mathbf{k}) = R_{ji}(-\mathbf{k}); \quad (26)$$

and, since the magnetic field is real,

$$R_{ij}(-\mathbf{k}) = R_{ij}^*(\mathbf{k}), \quad (27)$$

where the asterisk denotes complex conjugation. We will use these symmetry properties of the tensor elements in the following development.

Upon introducing equation (25) into equation (24), we explicitly exhibit all ϕ -dependence in the integrand of equation (24). The phase-angle averaging can now be done. Because of the orthogonality of the trigonometric functions in the interval $0-2\pi$, only certain values of n , relative to m , in the infinite summation will contribute, thereby reducing the double infinite sum to a single one. The result is

$$\begin{aligned} D_\mu(\tau) = & (2\pi)^{-3/2} \frac{(1 - \mu^2)}{2} \int_0^\tau d\lambda \int d^3k \sum_{m=-\infty}^{\infty} J_m\left(\frac{k_\perp \mu'}{\epsilon}\right) \exp[-i(k_\parallel \mu + m\epsilon)\lambda] \\ & \times \left\{ J_m\left(\frac{k_\perp \mu'}{\epsilon}\right) [(R_{11} + R_{22})(e^{i\epsilon\lambda} + e^{-i\epsilon\lambda}) - i(R_{12} - R_{21})(e^{i\epsilon\lambda} - e^{-i\epsilon\lambda})] \right. \\ & - (R_{11} - R_{22}) \left[J_{m-2}\left(\frac{k_\perp \mu'}{\epsilon}\right) e^{i(2\psi + \epsilon\lambda)} + J_{m+2}\left(\frac{k_\perp \mu'}{\epsilon}\right) e^{-i(2\psi + \epsilon\lambda)} \right] \\ & \left. + i(R_{12} + R_{21}) \left[J_{m-2}\left(\frac{k_\perp \mu'}{\epsilon}\right) e^{i(2\psi + \epsilon\lambda)} - J_{m+2}\left(\frac{k_\perp \mu'}{\epsilon}\right) e^{-i(2\psi + \epsilon\lambda)} \right] \right\}. \quad (28) \end{aligned}$$

At this point, we must remember that $D_\mu(\tau)$ is actually an integral operator (see eqs. [16] and [17]). Further simplifications of this expression for $D_\mu(\tau)$ depend on the function on which $D_\mu(\tau)$ operates. We must make the adiabatic approximation at this point to obtain the Fokker-Planck pitch-angle scattering coefficient, even though there are good reasons for suspecting that this approximation may not be valid (Klimas and Sandri 1971, 1973a). Klimas and Sandri (1973b) have shown that in the special case of isotropic, homogeneous magnetic turbulence, with a cosmic-ray distribution function that is essentially isotropic, the adiabatic approximation to the parallel transport of the cosmic-ray particles can be formally constructed, but its accuracy remains in doubt. In the general case being considered here, no formal justification of the existence of the adiabatic approximation is available. However, because it is our intention to compute the Fokker-Planck coefficient as a basis for further arguments on its validity, we will proceed anyway.

The adiabatic approximation to equation (16) is the Fokker-Planck equation for pitch-angle diffusion, which can be written as

$$\frac{\partial f}{\partial \tau} = \frac{1}{2} \frac{\partial}{\partial \mu} \left(\frac{\langle (\Delta \mu)^2 \rangle}{\Delta \tau} \right) \frac{\partial f}{\partial \mu}, \quad (29)$$

where

$$\frac{\langle (\Delta \mu)^2 \rangle}{\Delta \tau} = (\epsilon')^2 D_\mu(\infty). \quad (30)$$

In Appendix B we show that, by using the symmetry properties of equations (26)–(28), we obtain

$$\begin{aligned} \frac{\langle (\Delta \mu)^2 \rangle}{\Delta \tau} = & \frac{(\epsilon')^2(1 - \mu^2)}{2\sqrt{2\pi}} \int d^3k \sum_{m=-\infty}^{\infty} \delta(\epsilon m + k_\parallel \mu) J_{m-1}\left(\frac{k_\perp \mu'}{\epsilon}\right) \\ & \times \left[(R_{11} + R_{22}) J_{m-1}\left(\frac{k_\perp \mu'}{\epsilon}\right) - (R_{11} - R_{22}) J_{m+1}\left(\frac{k_\perp \mu'}{\epsilon}\right) \cos 2\psi \right. \\ & \left. - (R_{12} + R_{21}) J_{m+1}\left(\frac{k_\perp \mu'}{\epsilon}\right) \sin 2\psi + i(R_{12} - R_{21}) J_{m-1}\left(\frac{k_\perp \mu'}{\epsilon}\right) \right]. \quad (31) \end{aligned}$$

This result is applicable in a coordinate system with the 3-direction in the direction of the mean field. However,

by using the relationships between the angle ψ and the components of \mathbf{k} exhibited in equation (22), as well as the recursion relationships, $J_{m-1}(z) + J_{m+1}(z) = (2m/z)J_m(z)$ and $J_{m-1}(z) - J_{m+1}(z) = 2J'_m(z)$, we are able to rewrite equation (31) as

$$\begin{aligned} \frac{\langle(\Delta\mu^2)\rangle}{\Delta\tau} &= \frac{(\epsilon')^2(1-\mu^2)}{(2\pi)^{1/2}} \int d^3k \sum_{m=-\infty}^{\infty} \delta(\epsilon m + k_{\parallel}\mu) \\ &\times \left\{ -J_{m+1}\left(\frac{k_{\perp}\mu'}{\epsilon}\right) J_{m-1}\left(\frac{k_{\perp}\mu'}{\epsilon}\right) \left[\frac{\mathbf{k}_{\parallel} \cdot \mathbf{R}(\mathbf{k}) \cdot \mathbf{k}_{\parallel}}{k_{\perp}^2} \right] + \left(\frac{\epsilon m}{k_{\perp}\mu'}\right)^2 J_m^2\left(\frac{k_{\perp}\mu'}{\epsilon}\right) \text{Tr}[\mathbf{N} \cdot \mathbf{R}(\mathbf{k})] \right. \\ &\quad \left. - i \left(\frac{\epsilon m}{k_{\perp}\mu'}\right) J_m\left(\frac{k_{\perp}\mu'}{\epsilon}\right) J'_m\left(\frac{k_{\perp}\mu'}{\epsilon}\right) \text{Tr}[\boldsymbol{\Omega} \cdot \mathbf{R}(\mathbf{k})] \right\} \end{aligned} \quad (32)$$

which now may be applied in any coordinate system. (The symbol $\text{Tr}[\]$ indicates that the trace of the tensor inside the square bracket should be taken.) Equation (32) can be viewed as the pitch-angle scattering coefficient which results from resonant interactions between the particles and waves in the magnetic field with wavenumbers $k_{\parallel} = \epsilon m/\mu$. Similar expressions of this scattering coefficient have appeared previously for special forms of $\mathbf{R}(\mathbf{k})$ (Hasselman and Wibberenz 1968; Volk 1973; Fisk *et al.* 1974). In the next section we will consider the reduction of this general expression of the pitch-angle scattering coefficient for *any* homogeneous magnetic turbulence to the previously available special cases.

III. ISOTROPIC, AND SLAB, RANDOM FIELD MODELS

a) Isotropic Model for Magnetic Turbulence

The magnetic turbulence is statistically isotropic if the tensor $\mathbf{R}(\mathbf{k})$ has the form

$$\mathbf{R}(\mathbf{k}) = R(k) \left(\mathbf{I} - \frac{\mathbf{k}\mathbf{k}}{k^2} \right), \quad (33)$$

where \mathbf{I} is the unit matrix, and $R(k)$ is an arbitrary scalar function of $k = |\mathbf{k}|$ which can, however, be related to the one-dimensional power spectrum for the field components parallel to the displacement vector (Klimas and Sandri 1973a) through the relation (Batchelor 1960)

$$R(k) = \frac{1}{2}k \frac{d}{dk} \left(\frac{1}{k} \frac{dP_{\parallel}(k)}{dk} \right). \quad (34)$$

By substituting equation (33) into equation (32), we find, for isotropic turbulence,

$$\begin{aligned} \frac{\langle(\Delta\mu^2)\rangle}{\Delta\tau} &= \frac{(\epsilon')^2(1-\mu^2)}{\sqrt{2\pi}} \int d^3k \sum_{m=-\infty}^{\infty} \delta(k_{\parallel}\mu + \epsilon m) R(k) \\ &\times \left[-J_{m+1}\left(\frac{k_{\perp}\mu'}{\epsilon}\right) J_{m-1}\left(\frac{k_{\perp}\mu'}{\epsilon}\right) \left(\frac{k_{\parallel}}{k}\right)^2 + \left(\frac{\epsilon m}{k_{\perp}\mu'}\right)^2 J_m^2\left(\frac{k_{\perp}\mu'}{\epsilon}\right) \left(1 + \left(\frac{k_{\parallel}}{k}\right)^2\right) \right]. \end{aligned} \quad (35)$$

This result, written in a dimensional form, has been given by Fisk *et al.* (1974).

All terms in equation (35) which correspond to $m \neq 0$ do not contribute to $\langle(\Delta\mu^2)\rangle/\Delta\tau$ at $\mu = 0$. Notice that no part of the integrand depends on ψ . Thus, the integrations over ψ and k_{\parallel} can be carried out. Then

$$\begin{aligned} \frac{\langle(\Delta\mu^2)\rangle}{\Delta\tau} \Big|_{m \neq 0} &= (\epsilon')^2 (2\pi)^{1/2} \frac{(1-\mu^2)}{|\mu|} \int_0^{\infty} dk_{\perp} k_{\perp} R \left[k_{\perp}^2 + \left(\frac{\epsilon m}{\mu}\right)^2 \right]^{1/2} \\ &\times \left[-J_{m+1}\left(\frac{k_{\perp}\mu'}{\epsilon}\right) J_{m-1}\left(\frac{k_{\perp}\mu'}{\epsilon}\right) \left(\frac{k_{\parallel}}{k}\right)^2 + \left(\frac{\epsilon m}{k_{\perp}\mu'}\right)^2 J_m^2\left(\frac{k_{\perp}\mu'}{\epsilon}\right) \left(1 + \left(\frac{k_{\parallel}}{k}\right)^2\right) \right]_{k_{\parallel} = \epsilon m/\mu}. \end{aligned} \quad (36)$$

For a continuously differentiable magnetic field which is statistically homogeneous, $R(k)$ must approach zero as k approaches infinity, at least as fast as k^{-5} (Erdélyi 1956; Batchelor 1960). Therefore, from equation (36), $\langle(\Delta\mu^2)\rangle/\Delta\tau|_{m \neq 0} = 0$ at $\mu = 0$.

The $m = 0$ term in equation (35) contributes to $\langle(\Delta\mu^2)\rangle/\Delta\tau$ at $\mu = 0$ only. In fact,

$$\frac{\langle(\Delta\mu^2)\rangle}{\Delta\tau} \Big|_{m=0} = \delta(\mu) \frac{\epsilon'^2}{(2\pi)^{1/2}} \int d^3k \frac{R(k)}{|k_{\parallel}|} \left(\frac{k_{\parallel}}{k}\right)^2 J_1^2\left(\frac{k_{\perp}}{\epsilon}\right). \quad (37)$$

Thus, it is the $m = 0$ term which contributes the $\delta(\mu)$ part of the pitch-angle scattering coefficient. Looking back at equation (32), we can see that this conclusion is completely general.

The infinite series of resonant terms ($m \neq 0$) has been investigated by Fisk *et al.* (1974), for power-law power spectra. They found that, for $1 \leq \epsilon \leq 30$, the entire series can be very well approximated by μ times the small-gyroradius approximation given by Jokipii (1971). This result applies over a wide range of values of the power-law spectral index.

b) Slab Model for Magnetic Turbulence

In the slab model we let

$$\mathbf{R}(\mathbf{k}) = NR(k_{\parallel})\delta(\mathbf{k}_{\perp}), \quad (38)$$

where $R(k_{\parallel})$ is an arbitrary even function of k_{\parallel} . Thus, the random field components are normal to the mean field, and they are functions only of distance along the mean field. The correlations in the random field are assumed cylindrically symmetric about the mean field.

This model is closely related to others that have been studied previously. In Jokipii's (1972) plane polarized model, the random field is assumed to have only one orthogonal component. We will see shortly that the pitch-angle scattering coefficient computed here is simply twice Jokipii's. Jones, Kaiser, and Birmingham (1973a, b) have used the plane-polarized field model to investigate modifications of quasi-linear theory near $\mu = 0$. Volk (1974) has also studied such modifications in a random field made up of transverse Alfvén waves. A common feature of all of these models is that the $\delta(\mu)$ does not appear in the Fokker-Planck pitch-angle scattering coefficient.

With the choice for $R(k)$ given in equation (43), we find, from equation (32), that

$$\frac{\langle(\Delta\mu)^2\rangle}{\Delta\tau} = 2 \frac{\epsilon'^2(1-\mu^2)}{(2\pi)^{1/2}} \int d^3k \delta(\mathbf{k}_{\perp}) R(k_{\parallel}) \sum_{m=-\infty}^{\infty} \delta(k_{\parallel}\mu + \epsilon m) \left(\frac{\epsilon m}{k_{\perp}\mu'}\right)^2 J_m^2\left(\frac{k_{\perp}\mu'}{\epsilon}\right). \quad (39)$$

We see immediately that there can be no $\delta(\mu)$ contribution to $\langle(\Delta\mu)^2\rangle/\Delta\tau$ since the $m = 0$ term in equation (39) is zero. In fact, because of the $\delta(\mathbf{k}_{\perp})$, only the $m = \pm 1$ terms are nonzero. All other terms in the expansion are zero because $(\epsilon m/k_{\perp}\mu')J_m(k_{\perp}\mu'/\epsilon) \rightarrow 0$ as $k_{\perp} \rightarrow 0$, and the integrand can contribute only at $k_{\perp} = 0$. Thus

$$\frac{\langle(\Delta\mu)^2\rangle}{\Delta\tau} = \epsilon'^2(2\pi)^{1/2} \left(\frac{1-\mu^2}{|\mu|}\right) R\left(k_{\parallel} = \frac{\epsilon}{\mu}\right). \quad (40)$$

(NOTE. $\delta(\mathbf{k}_{\perp})$ is normalized to 2π instead of 1.) This result is exactly twice that of Jokipii's (1972) plane-polarized field model calculation in which there is only one orthogonal field component instead of two.

IV. MIRRORING AND THE DELTA FUNCTION

The fact that the $\delta(\mu)$ contribution to $\langle(\Delta\mu)^2\rangle/\Delta\tau$ vanished in the slab model was one of the first strong indications that this contribution represents mirroring in the random magnetic field. The guiding center approximation (Alfvén and Falthammar 1963; Northrop 1963) to the motion of charged particles in the random magnetic field applies to particles whose motion along the field is so slow that they can resonate only with very short wavelengths where the power density is typically negligible. This could be the situation for a very low energy particle, or a higher energy particle moving in a weakly turbulent field with pitch angle very near 90° . The latter situation is the one being studied here.

Within the guiding center approximation, the pitch angle changes with time according to

$$p \frac{d\mu}{dt} = -M\mathbf{e} \cdot [(\mathbf{e} \cdot \nabla)\mathbf{B}], \quad (41)$$

where $M = \frac{1}{2}p_{\perp}^2/Bm$ is the magnetic moment, and \mathbf{e} is a unit vector in the direction of the local magnetic field. Of course, the pitch angle that enters into this expression is the angle between the direction of the particle momentum and the local exact field (actually the angle averaged over a gyroperiod). In Appendix C we demonstrate that, to first order in η in a weakly turbulent magnetic field, this pitch angle is identical to the pitch angle relative to the mean field averaged over a gyroperiod. By introducing $\mathbf{B} = \langle\mathbf{B}\rangle + \mathbf{B}'$, we can rewrite equation (41). To first order in η , we find,

$$\frac{d\mu}{d\tau} = -\frac{1}{2}(1-\mu^2)\eta\hat{\beta} \cdot [(\hat{\beta} \cdot \nabla)\hat{\beta}'(x)], \quad (42)$$

where $\frac{1}{2}(1-\mu^2)\eta$ is the magnetic moment, which we consider a constant of the motion in this guiding center approximation. Since the pitch-angle scattering coefficient is quadratic in $\Delta\mu$, we see that, in order to obtain an $O(\eta^2)$ contribution to that coefficient, we must find an $O(\eta)$ contribution to $d\mu/d\tau$ through equation (42). Equation (42) is nonzero only if the random field contains a component in the direction of the mean field which also has

spatial gradients in that direction. In the slab model, $\hat{\beta} \cdot \beta'(x) = 0$, and there is also no $\delta(\mu)$ contribution to $\langle(\Delta\mu)^2\rangle/\Delta\tau$. In this section we will show that *if and only if* equation (42) is nonzero, do we find a $\delta(\mu)$ in the pitch-angle scattering coefficient. Thus, only when there are $O(\eta)$ changes in μ due to guiding center motion is there a $\delta(\mu)$ in $\langle(\Delta\mu)^2\rangle/\Delta\tau$. In view of the fact that the $\delta(\mu)$ is the *only* part of $\langle(\Delta\mu)^2\rangle/\Delta\tau$ which is nonzero at $\mu = 0$, and therefore is the only part which can possibly reverse the parallel motion of the particles, we ascribe the $\delta(\mu)$ to mirroring. The fact that mirroring exhibits itself as a $\delta(\mu)$ in $\langle(\Delta\mu)^2\rangle/\Delta\tau$ can be understood from the asymptotic nature of the Fokker-Planck equation. This equation is asymptotic in small η , and in an infinitesimal random field only particles very near $\mu = 0$ could possibly be mirrored. In fact, the δ -function is probably a crude approximation to a sharply peaked function of μ .

It is interesting to note the results of a calculation which we present in Appendix D. There, we compute $\langle(\Delta\mu)^2\rangle/\Delta\tau$ from equation (42) using the standard approach (cf. Appendix A). For the position of the particle, x , along its trajectory in the field, as it appears in $\hat{\beta}'(x)$, we substitute the position of its guiding center. This approximation to the position of the particle is exact in the limit of zero particle energy ($\epsilon = \infty$). The result of this calculation is identical to the result which we will present in this section when we study the $m = 0$ term [the $\delta(\mu)$ contribution] of equation (32) in the limit $\epsilon = \infty$. Thus, the Fokker-Planck coefficient for pitch-angle scattering, computed in the guiding center approximation to the particle trajectory, also contains a $\delta(\mu)$ term which is in exact agreement with the $\delta(\mu)$ term obtained from the quasi-linear approximation to the particle trajectory, in the limit $\epsilon = \infty$.

From equation (32), with $m = 0$, we find

$$\left. \frac{\langle(\Delta\mu)^2\rangle}{\Delta\tau} \right|_{m=0} = \delta(\mu) \frac{(\epsilon')^2(1-\mu^2)}{(2\pi)^{1/2}} \int d^3k \left(\frac{\mathbf{k}_{\parallel} \cdot \mathbf{R}(\mathbf{k}) \cdot \mathbf{k}_{\parallel}}{|\mathbf{k}_{\parallel}|} \right) \left[\frac{J_1^2\left(\frac{k_{\perp}\mu}{\epsilon}\right)}{k_{\perp}^2} \right]. \quad (43)$$

A necessary property of $\mathbf{R}(\mathbf{k})$ is that it be nonnegative, i.e., the quadratic form,

$$\mathbf{V} \cdot \mathbf{R} \cdot \mathbf{V}^* \geq 0, \quad (44)$$

for any complex vector \mathbf{V} (Batchelor 1960). In particular, $\mathbf{k}_{\parallel} \cdot \mathbf{R} \cdot \mathbf{k}_{\parallel}$ must be either positive or zero for any \mathbf{k} . Notice also that the rest of the quantities in the integrand of equation (43) are nonnegative. Therefore, barring special situations which we will discuss in a moment, $R_{\parallel,\parallel}(\mathbf{k})$ must be zero everywhere in \mathbf{k} in order for the $\delta(\mu)$ to not contribute to pitch-angle scattering. But then, $R_{\parallel,\parallel}(\mathbf{r})$ is zero for all \mathbf{r} , from which we can conclude that there is no random field component parallel to the mean field (slab or plane-polarized field model). With no random field parallel to $\hat{\beta}$, we see from equation (42) that mirroring is impossible. This argument can be carried in the reverse order. It is clear that if equation (42) indicates no mirroring because $\hat{\beta} \cdot \beta'(x)$ is zero for all x , then $R_{\parallel,\parallel}(\mathbf{r}) = 0$, and finally $R_{\parallel,\parallel}(\mathbf{k}) = 0$ for all \mathbf{k} . In this case there is no $\delta(\mu)$ contribution to $\langle(\Delta\mu)^2\rangle/\Delta\tau$.

There is a special situation in which $R_{\parallel,\parallel}(\mathbf{k}) \neq 0$ everywhere in \mathbf{k} , and yet the integral in equation (43) is zero, and $\langle(\Delta\mu)^2\rangle/\Delta\tau$ does not contain a $\delta(\mu)$. It is possible that $R_{\parallel,\parallel}(\mathbf{k})$ is nonzero only at that point in \mathbf{k} where other terms in the integrand are zero. The Bessel function contains isolated zeros which, nevertheless, we will not consider because the positions of these zeros are rigidity dependent through ϵ , and $\mathbf{R}(\mathbf{k})$ does not depend on rigidity. On the other hand, we could have $\mathbf{R}(\mathbf{k})$ nonzero only when $k_{\parallel} = 0$ without contributing to the integral. But we still have

$$k_{\parallel} R_{\parallel,\parallel}(\mathbf{k}) = 0. \quad (45)$$

This statement is the Fourier transform of

$$\frac{\partial}{\partial r_{\parallel}} R_{\parallel,\parallel}(\mathbf{r}) = 0. \quad (46)$$

Barring nondifferentiable functions of the type discussed by Wiener (1933), we can conclude that the random field component in the direction of the mean field is independent of position along the mean field; i.e., $(\hat{\beta} \cdot \nabla)[\hat{\beta} \cdot \beta'(x)] = \hat{\beta} \cdot [(\hat{\beta} \cdot \nabla)\beta'(x)] = 0$. But this is exactly the condition that equation (42) be zero. This argument can also be reversed. Starting from a zero for equation (42), we can conclude equation (46), then equation (45), and then we can conclude no $\delta(\mu)$ in $\langle(\Delta\mu)^2\rangle/\Delta\tau$. In general, a random field component along the mean field is allowed, but this component must be a constant along the field; in the slab model the constant is zero.

From the above arguments, we conclude that the $\delta(\mu)$ in $\langle(\Delta\mu)^2\rangle/\Delta\tau$ represents the physical phenomenon of mirroring in the random magnetic field.

For $\epsilon \gg 1$, equation (43) can be simplified considerably. Using $J_1(k_{\perp}/\epsilon) \approx \frac{1}{2}(k_{\perp}/\epsilon)$, we find

$$\left. \frac{\langle(\Delta\mu)^2\rangle}{\Delta\tau} \right|_{m=0} \approx \tilde{\epsilon} \left(\frac{(1-\mu^2)\eta^2}{2} \right)^2 \frac{\delta(\mu)}{(2\pi)^{1/2}} \int d^3k \left(\frac{\mathbf{k}_{\parallel} \cdot \mathbf{R}(\mathbf{k}) \cdot \mathbf{k}_{\parallel}}{|\mathbf{k}_{\parallel}|} \right). \quad (47)$$

Thus, in this approximation, as the rigidity gets small, $\langle(\Delta\mu)^2\rangle/\Delta\tau$ becomes rigidity independent. However, this

approximation is valid only when $\mathbf{R}(\mathbf{k})$ approaches zero rapidly enough with increasing k_{\perp} so that the approximation for $J_1(k_{\perp}/\epsilon)$ is valid for all significant values of k_{\perp} in the integrand of equation (43). In the interplanetary magnetic field we do expect this condition to be satisfied at some large enough value of ϵ , even though it is difficult to say what that value should be. Furthermore, there is considerable evidence (Bryant *et al.* 1965; O'Gallagher 1967; Cline and McDonald 1968) that the parallel diffusion coefficient becomes rigidity independent for low-rigidity particles in the interplanetary field. A possible explanation of this observation is that the contribution of mirroring to pitch-angle scattering dominates all other contributions for low rigidities ($\epsilon \gg 1$).

As we mentioned above, equation (47) is identical to the result (given in Appendix D) of computing $\langle(\Delta\mu)^2\rangle/\Delta\tau$ within the guiding center approximation to the particle motion.

V. CONCLUSION

We have constructed the Fokker-Planck coefficient for pitch-angle scattering of cosmic rays in otherwise arbitrary, but statistically homogeneous, magnetic turbulence. Our result was obtained both from first principles and through the standard approach of Chandrasekhar (1943). The reduction of our expression to previously available scattering coefficients, calculated in special models of the random field, has been discussed.

We have shown that the pitch-angle scattering coefficient contains a Dirac $\delta(\mu)$ in $\mu = \cos \theta$, where θ is the pitch angle. We have, further, proved that this δ -function is the prediction of the quasi-linear approximation for the contribution of mirroring to pitch-angle scattering in a weakly turbulent magnetic field.

The $\delta(\mu)$ does not contribute to pitch-angle scattering when, within the guiding center approximation to the particle motion, the pitch angle is a constant of the motion of the particle to $O(\eta)$ in the random field strength. This condition is met when the vector component of the random field which lies in the direction of the mean field is independent of distance along the mean field. The slab, or plane-polarized, or linearized Alfvén wave models of the random field, all of which have no random field component along the mean direction, fall within this class. Magnetosonic waves propagating across the mean field have a random parallel field component which is independent of distance along the mean field, and fall within this class. Typically, however, the observed interplanetary field does not fall within this class. We expect mirroring in the interplanetary field, and we expect a δ -function in the appropriate Fokker-Planck pitch-angle scattering coefficient.

The fact that mirroring exhibits itself as a $\delta(\mu)$ in the Fokker-Planck pitch-angle scattering coefficient indicates that this particular pitch-angle scattering mechanism is misordered within the quasi-linear approximation. In the quasi-linear formalism, which leads to the Fokker-Planck equation, it is *assumed* that the effects of the random field on the motion of the particles is $O(\eta^2)$ in the random field strength. The appearance of the δ -function suggests that this assumption is incorrect.

Other investigators have concerned themselves with the modifications of the quasi-linear formalism which are necessary even in the special random field models in which mirroring does not occur. In this paper we have demonstrated the existence of the δ -function, as well as its connection to mirroring, in order that further discussions of the validity and/or modifications of the quasi-linear approximation can be based on its actual predictions in realistic field models.

We would like to acknowledge stimulating discussions with Drs. L. A. Fisk and T. B. Kaiser.

APPENDIX A

With our notation, the momentum equation of motion for a charged particle in a magnetic field, can be written as

$$\frac{d\hat{\mathbf{p}}}{d\tau} = \epsilon[\boldsymbol{\Omega} + \eta\boldsymbol{\Omega}'] \cdot \hat{\mathbf{p}} \quad (\text{A1})$$

where $\Omega_{ij} = \epsilon_{ijk}\beta_k$ and $\Omega'_{ij} = \epsilon_{ijk}\beta'_k$. Thus, for $\mu = \hat{\boldsymbol{\beta}} \cdot \hat{\mathbf{p}}$ we find

$$\frac{d\mu}{d\tau} = \epsilon'(\hat{\boldsymbol{\beta}} \cdot \boldsymbol{\Omega}' \cdot \hat{\mathbf{p}}), \quad (\text{A2})$$

or

$$\frac{d\mu}{d\tau} = \epsilon'(\hat{\mathbf{p}} \cdot \boldsymbol{\Omega} \cdot \hat{\boldsymbol{\beta}}'), \quad (\text{A3})$$

where, in obtaining equation (A2), we have used $\boldsymbol{\Omega} \cdot \hat{\boldsymbol{\beta}} = 0$. By formally integrating equation (A3) we can obtain

$$\langle(\Delta\mu)^2\rangle = -(\epsilon')^2 \int_0^{\Delta\tau} d\lambda \int_0^{\Delta\tau} d\lambda' \hat{\mathbf{p}}(\lambda) \cdot \boldsymbol{\Omega} \cdot \langle\boldsymbol{\beta}'(\lambda)\boldsymbol{\beta}'(\lambda')\rangle \cdot \boldsymbol{\Omega} \cdot \hat{\mathbf{p}}(\lambda'), \quad (\text{A4})$$

where $\boldsymbol{\beta}'(\lambda) \equiv \boldsymbol{\beta}'[\mathbf{x}(\lambda)]$, and $\mathbf{x}(\lambda)$ and $\hat{\mathbf{p}}(\lambda)$ are the time-dependent coordinates given by equations (8)–(12) which describe the zeroth-order helical approximation to the particle motion. Now, by introducing the Fourier integral

transform representation of $\beta'(\mathbf{x})$, as well as the assumption of homogeneity, as expressed through equation (19), we find

$$\langle(\Delta\mu)^2\rangle = -\frac{(\epsilon')^2}{(2\pi)^{3/2}} \int_0^{\Delta\tau} d\lambda \int_0^{\Delta\tau} d\lambda' \int d^3k \exp[ik_{\parallel}\mu(\lambda - \lambda')] \exp\{ik_{\perp} \cdot [x_{\perp}(\lambda) - x_{\perp}(\lambda')]\} [\hat{p}(\lambda) \cdot \Omega \cdot \mathbf{R}(k) \cdot \Omega \cdot \hat{p}(\lambda')]. \quad (\text{A5})$$

Because $\langle(\Delta\mu)^2\rangle$ is a scalar quantity, it is invariant to rotations. We also choose a special coordinate system in which to proceed here. In a Cartesian coordinate system, we let the mean field lie in the 3-direction, as in the main text, but we further rotate our coordinate system so that the perpendicular component of the momentum, (\hat{p}_{\perp}), lies in the 1-direction. Then, on introducing the expansion, given by equation (23), of the exponentials in equation (A5), we obtain

$$\begin{aligned} \langle(\Delta\mu)^2\rangle = & -\frac{(\epsilon')^2}{(2\pi)^{3/2}} \int_0^{\Delta\tau} d\lambda \int_0^{\Delta\tau} d\lambda' \int_{-\infty}^{\infty} dk_{\parallel} \exp[ik_{\parallel}\mu(\lambda - \lambda')] \int_0^{2\pi} d\psi \sum_{m,n=-\infty}^{\infty} e^{i(m-n)\psi} \\ & \times e^{in\epsilon\lambda} e^{-im\epsilon\lambda'} \int_0^{\infty} dk_{\perp} k_{\perp} J_n\left(\frac{k_{\perp}\mu}{\epsilon}\right) J_m\left(\frac{k_{\perp}\mu'}{\epsilon}\right) [\hat{p}(\lambda) \cdot \Omega \cdot \mathbf{R}(k) \cdot \Omega \cdot \hat{p}(\lambda')]. \end{aligned} \quad (\text{A6})$$

By introducing the tensor components of $\mathbf{R}(k)$, we find

$$\begin{aligned} \langle(\Delta\mu)^2\rangle = & \frac{(\epsilon')^2(1 - \mu^2)}{2(2\pi)^{3/2}} \int_0^{\Delta\tau} d\lambda \int_0^{\Delta\tau} d\lambda' \int_{-\infty}^{\infty} dk_{\parallel} \exp[ik_{\parallel}\mu(\lambda - \lambda')] \int_0^{2\pi} d\psi e^{i(m-n)\psi} \\ & \times e^{in\epsilon\lambda} e^{-im\epsilon\lambda'} \int_0^{\infty} dk_{\perp} k_{\perp} J_n\left(\frac{k_{\perp}\mu}{\epsilon}\right) J_m\left(\frac{k_{\perp}\mu'}{\epsilon}\right) \\ & \times [(R_{11} + R_{22}) \cos \epsilon(\lambda - \lambda') - (R_{11} - R_{22}) \cos \epsilon(\lambda + \lambda') \\ & - (R_{12} + R_{21}) \sin \epsilon(\lambda + \lambda') - (R_{12} - R_{21}) \sin \epsilon(\lambda - \lambda')]. \end{aligned} \quad (\text{A7})$$

Now, following the usual arguments, we must find the parts of this expression which grow as $\Delta\tau$. Consider, for example, the integral

$$I_1 \equiv \int_0^{\Delta\tau} d\lambda \int_0^{\Delta\tau} d\lambda' \exp[ik_{\parallel}\mu(\lambda - \lambda')] e^{in\epsilon\lambda} e^{-im\epsilon\lambda'} \cos \epsilon(\lambda - \lambda') \quad (\text{A8})$$

which is associated with the term $(R_{11} + R_{22})$ in equation (A7). By introducing a new variable of integration $S = \lambda' - \lambda$, for λ' , and inverting the order of integration, we find

$$I_1 = \left[\int_0^{\Delta\tau} dS \int_0^{\Delta\tau-S} d\lambda + \int_{-\Delta\tau}^0 dS \int_{-S}^{\Delta\tau} d\lambda \right] \exp[-i(k_{\parallel}\mu + \epsilon m)S \cos \epsilon S] e^{i(n-m)\epsilon\lambda} \quad (\text{A9})$$

When $n = m$, we do obtain a term which grows linearly with $\Delta\tau$, for large $\Delta\tau$, given by

$$\Delta\tau\pi\{\delta[k_{\parallel}\mu + (m+1)\epsilon] + \delta[k_{\parallel}\mu + (m-1)\epsilon]\}. \quad (\text{A10})$$

A further examination of equation (A9) reveals no other terms with the linear growth property. In order to obtain the Fokker-Planck coefficient from equation (A7), we replace I_1 , in equation (A7), with

$$\Delta\tau\delta_{m,n}\pi\{\delta[k_{\parallel}\mu + (m+1)\epsilon] + \delta[k_{\parallel}\mu + (m-1)\epsilon]\}. \quad (\text{A11})$$

This procedure can be continued for the rest of the time integrals, and the steps which follow are identical to the steps taken in the alternate derivation of $\langle(\Delta\mu)^2\rangle/\Delta\tau$ given in the main text. As we mentioned there, the results of the two calculations are identical.

The point of this Appendix is to note the difference between the arguments given here, and those given in the main text. From equation (A9), we actually find a variety of terms, some of which do not grow unbounded with $\Delta\tau$, some which grow linearly with $\Delta\tau$, and some which grow as $(\Delta\tau)^2$. From the other time integrals in equation (A7) we find similar results with, in addition, terms which grow as $\Delta\tau$ times trigonometric functions of $\Delta\tau$; these terms oscillate with amplitudes that grow unbounded in time. In the face of these unbounded oscillations, the usual argument for choosing the terms that grow as $\Delta\tau$ in $\langle(\Delta\mu)^2\rangle$, and dropping all others, becomes difficult to support. Even ignoring this problem, we still have the problem of choosing the $\Delta\tau$ terms by arguing that $\Delta\tau$ is large compared with the interaction time, and small compared with the relaxation time, when in fact these two time scales are not clearly separated for particles in a magnetic field.

In comparison, from equation (24), notice that we never face these problems in the derivation of $\langle(\Delta\mu)^2\rangle/\Delta\tau$

given there. The inner time integration in equation (A7) is replaced by the phase-averaging integration over ϕ in equation (24). The effects of carrying out the time integration here, and the phase averaging there, are identical; both procedures pick the same terms out of the infinite expansions for retention. Thus, the results of both approaches are identical, but the reasoning contained in the standard approach is much more difficult to support.

APPENDIX B

The adiabatic approximation to equation (28) permits the time integration to be done. The result is

$$\begin{aligned}
 D_\mu(\infty) = & (2\pi)^{-3/2} \left(\frac{1 - \mu^2}{2} \right) \int d^3k \sum_{m=-\infty}^{\infty} J_m \left(\frac{k_\perp \mu'}{\epsilon} \right) \\
 & \times \left(J_m \left(\frac{k_\perp \mu'}{\epsilon} \right) \left[i(R_{11} + R_{22}) \{ \zeta[\epsilon(1 - m) - k_{\parallel} \mu] + \zeta[-\epsilon(1 + m) - k_{\parallel} \mu] \} \right. \right. \\
 & \quad \left. \left. + (R_{12} - R_{21}) \{ \zeta[\epsilon(1 - m) - k_{\parallel} \mu] - \zeta[-\epsilon(1 + m) - k_{\parallel} \mu] \} \right] \right. \\
 & \quad \left. - i(R_{11} - R_{22}) \left\{ J_{m-2} \left(\frac{k_\perp \mu'}{\epsilon} \right) e^{i2\psi} \zeta[\epsilon(1 - m) - k_{\parallel} \mu] + J_{m+2} \left(\frac{k_\perp \mu'}{\epsilon} \right) e^{-i2\psi} \zeta[-\epsilon(1 + m) - k_{\parallel} \mu] \right\} \right. \\
 & \quad \left. - (R_{12} + R_{21}) \left\{ J_{m-2} \left(\frac{k_\perp \mu'}{\epsilon} \right) e^{i2\psi} \zeta[\epsilon(1 - m) - k_{\parallel} \mu] - J_{m+2} \left(\frac{k_\perp \mu'}{\epsilon} \right) e^{-i2\psi} \zeta[-\epsilon(1 + m) - k_{\parallel} \mu] \right\} \right) .
 \end{aligned} \tag{B1}$$

The function $\zeta(x)$ is the zeta function defined by (Heitler 1954)

$$\zeta(x) = -i \lim_{\tau \rightarrow \infty} \int_0^\tau d\lambda e^{i\lambda x} = \frac{P}{x} - i\pi \delta(x) = -\zeta^*(-x), \tag{B2}$$

where P/x is the principal value of $1/x$.

In the terms that multiply $\zeta[\epsilon(1 - m) - k_{\parallel} \mu]$, let $m' = -m$ and $k' = -k$, and use the relation $J_{-m}(z) = (-1)^m J_m(z)$ along with the symmetry relations (26) and (27), and (B2) to rewrite (B1) in the form

$$\begin{aligned}
 D_\mu(\infty) = & (2\pi)^{-3/2} \left(\frac{1 - \mu^2}{2} \right) \int d^3k \sum_{m=-\infty}^{\infty} J_m \{ J_m(\zeta - \zeta^*) [i(R_{11} + R_{22}) - (R_{12} - R_{21})] \\
 & - iJ_{m+2}(R_{11} - R_{22}) [e^{i2\psi} \zeta - e^{-i2\psi} \zeta^*] - J_{m+2}(R_{12} + R_{21}) [e^{i2\psi} \zeta + e^{-i2\psi} \zeta^*] \},
 \end{aligned} \tag{B3}$$

where $J_m = J_m(k_\perp \mu' / \epsilon)$, $\zeta = \zeta[\epsilon(1 + m) + k_{\parallel} \mu]$, and $\zeta^* = \zeta^*[\epsilon(1 + m) + k_{\parallel} \mu]$. The terms containing zeta functions can be rewritten as

$$\begin{aligned}
 \zeta(x) - \zeta^*(x) &= -2\pi i \delta(x), \\
 e^{i2\psi} \zeta(x) - e^{-i2\psi} \zeta^*(x) &= 2i \left\{ \sin 2\psi \left(\frac{P}{x} \right) - \pi \cos 2\psi [\delta(x)] \right\}, \\
 e^{i2\psi} \zeta(x) + e^{-i2\psi} \zeta^*(x) &= 2 \left\{ \cos 2\psi \left(\frac{P}{x} \right) + \pi \sin 2\psi [\delta(x)] \right\}.
 \end{aligned} \tag{B4}$$

Now consider terms of the form

$$\int d^3k \sum_{m=-\infty}^{\infty} J_m J_{m+2} \left\{ \frac{(R_{11} - R_{22}) \sin 2\psi}{(R_{12} + R_{21}) \cos 2\psi} \right\} \frac{P}{\epsilon(1 + m) + k_{\parallel} \mu} \tag{B5}$$

and let $n = m + 1$, so that equation (B5) becomes

$$\int d^3k \sum_{n=-\infty}^{\infty} J_{n-1} J_{n+1} \left\{ \frac{(R_{11} - R_{22}) \sin 2\psi}{(R_{12} + R_{21}) \cos 2\psi} \right\} \frac{P}{\epsilon n + k_{\parallel} \mu}. \tag{B6}$$

If we let $n \rightarrow -n$ and $k \rightarrow -k$ in equation (B6), then equation (B6) is equal to minus itself and all terms in (B3) which contain principal value contributions are identically equal to zero. Equation (31c) follows immediately.

APPENDIX C

Just for the purpose of developing this argument, in this Appendix we will introduce a special notation which is different from that contained in the rest of this paper. Let

$$\mu(\tau) = (\mathbf{B}/B) \cdot \hat{\mathbf{p}}(\tau) \quad (\text{C1})$$

be the cosine of the pitch angle relative to the local (exact) field. Within the guiding center approximation to the particle motion, we assume (\mathbf{B}/B) constant over a gyroperiod, and assume that the particle moves in a helical trajectory in the field. Thus,

$$\mu(\tau) = \left(\frac{\mathbf{B}}{B}\right) \cdot [\bar{\mathbf{P}} + \bar{N} \cos \epsilon\tau - \bar{\Omega} \sin \epsilon\tau] \cdot \hat{\mathbf{p}}(0) \quad (\text{C2})$$

where the projection matrices, $\bar{\mathbf{P}}$, \bar{N} , and $\bar{\Omega}$, are identical to those defined by equations (5), (13), and (14) except that they are based on the local, rather than the mean, field. We now introduce the notation $\langle \mu \rangle \equiv \hat{\beta} \cdot \hat{\mathbf{p}}$, for the cosine of the pitch angle relative to the mean field. The quantity which enters equation (41) is $\{\mu(\tau)\}$,¹ which is the average over one gyroperiod, of $\mu(\tau)$:

$$\{\mu(\tau)\} = (\mathbf{B}/B) \cdot \bar{\mathbf{P}} \cdot \hat{\mathbf{p}}(0) \equiv \mu. \quad (\text{C3})$$

The time dependence of $\langle \mu \rangle$ is given by

$$\langle \mu \rangle = \hat{\beta} \cdot [\bar{\mathbf{P}} + \bar{N} \cos \epsilon\tau - \bar{\Omega} \sin \epsilon\tau] \cdot \hat{\mathbf{p}}(0) \quad (\text{C4})$$

and its average over one gyroperiod is,

$$\langle \mu \rangle = \hat{\beta} \cdot \left[\frac{\hat{\beta} + \eta \beta'}{[1 + 2\eta(\hat{\beta} \cdot \beta') + \eta^2(\beta' \cdot \beta')]^{1/2}} \right] \mu. \quad (\text{C5})$$

Thus,

$$\{\langle \mu \rangle\} = \mu + O(\eta^2). \quad (\text{C6})$$

To $O(\eta)$ in the random field strength, $\langle \mu \rangle$ and μ are identical when averaged over a gyroperiod. This averaging is implied in equation (41) since it follows from the guiding center approximation. In the rest of this paper we therefore simply use the symbol μ to stand for the cosine of any of the relevant pitch angles.

APPENDIX D

By introducing the Fourier transform representation of the random field into equation (42), we find

$$\frac{d\mu}{d\tau} = -\frac{1}{2}i(1 - \mu^2)\eta \left(\frac{1}{2\pi}\right)^{3/2} \int d^3k \hat{\beta} \cdot [(\hat{\beta} \cdot \nabla) \exp [ik \cdot \mathbf{x}(\tau)] \beta'(k)], \quad (\text{D1})$$

or

$$\frac{d\mu}{d\tau} = -\frac{1}{2}(1 - \mu^2)\eta \left(\frac{1}{2\pi}\right)^{3/2} \int d^3k [\hat{\beta} \cdot \beta'(k)] (\hat{\beta} \cdot k) \exp [ik \cdot \mathbf{x}(\tau)]. \quad (\text{D2})$$

Now, on remembering that, in this notation, $(1 - \mu^2)\eta/2$ is the magnetic moment of the particle which is an adiabatic invariant of the motion of the particle in the guiding center approximation, we can formally integrate equation (D2) and form its square to find

$$\begin{aligned} \langle (\Delta\mu)^2 \rangle &= - \left[\frac{(1 - \mu^2)\eta}{2} \right]^2 \left(\frac{1}{2\pi} \right)^3 \int_0^{\Delta\tau} d\lambda \int_0^{\Delta\tau} d\lambda' \int d^3k \int d^3k' \exp [ik \cdot \mathbf{x}(\lambda)] \exp [ik' \cdot \mathbf{x}(\lambda')] \\ &\quad \times [(k \cdot \hat{\beta}) \hat{\beta} \cdot \langle \beta'(k) \beta'(k') \rangle \cdot \hat{\beta} (\hat{\beta} \cdot k')]. \end{aligned} \quad (\text{D3})$$

If the magnetic turbulence is statistically homogeneous, we have equation (19), which can be inserted into equation (D3) to obtain

$$\langle (\Delta\mu)^2 \rangle = \left[\frac{(1 - \mu^2)\eta}{2} \right]^2 \left(\frac{1}{2\pi} \right)^{3/2} \int_0^{\Delta\tau} d\lambda \int_0^{\Delta\tau} d\lambda' \int d^3k \exp \{ik \cdot [\mathbf{x}(\lambda) - \mathbf{x}(\lambda')]\} [k_{\parallel} \cdot \mathbf{R}(k) \cdot k_{\parallel}]. \quad (\text{D4})$$

¹ In this Appendix the curly braces denote an average over one gyroperiod.

As explained in the main text, we approximate the position of the particle, by the position of its guiding center. Thus, $\mathbf{k} \cdot [\mathbf{x}(\lambda) - \mathbf{x}(\lambda')] = \mathbf{k}_{\parallel} \mu (\lambda - \lambda')$. Following the usual argument (see Appendix A) for obtaining the part of equation (D4) which grows linearly with $\Delta\tau$, when $\Delta\tau$ is large, we obtain

$$\frac{\langle (\Delta\mu)^2 \rangle}{\Delta\tau} = \left[\frac{(1 - \mu^2)\eta}{2} \right]^2 \frac{\delta(\mu)}{(2\pi)^{1/2}} \int d^3k \left[\frac{\mathbf{k}_{\parallel} \cdot \mathbf{R}(\mathbf{k}) \cdot \mathbf{k}_{\parallel}}{|\mathbf{k}_{\parallel}|} \right]. \quad (\text{D5})$$

This result is identical to that given by equation (47) which was obtained, in the low-energy limit, from the usual quasi-linear approximation to the particle motion.

REFERENCES

- Alfvén, H., and Fälthammar, C. 1963, *Cosmical Electrodynamics* (Oxford: Clarendon Press).
- Batchelor, G. K. 1960, *The Theory of Homogeneous Turbulence* (1st ed.; Cambridge: Cambridge University Press).
- Bryant, D. A., Cline, T. L., Desai, U. D., and McDonald, F. B. 1965, *Ap. J.*, **141**, 478.
- Cesarsky, C. F., and Kulsrud, R. M. 1972, Princeton Univ. Plasma Physics Lab. Rept. PPL-AP53.
- Chandrasekhar, S. 1943, *Rev. Mod. Phys.*, **15**, 1.
- Cline, T. L., and McDonald, F. B. 1968, *Solar Phys.*, **5**, 507.
- Earl, J. A. 1974, Tech. Rpt. 74-078, Univ. of Maryland, College Park.
- Erdélyi, A. 1956, *Asymptotic Expansions* (New York: Dover).
- Fisk, L. A., Goldstein, M. L., Klimas, A. J., and Sandri, G. 1974, *Ap. J.*, **190**, 417.
- Hall, D. E., and Sturrock, P. A. 1967, *Phys. Fluids*, **10**, 2620.
- Hasselmann, K., and Wibberenz, G. 1968, *Zs. f. Geophys.*, **34**, 353.
- Heitler, W. 1954, *The Quantum Theory of Radiation* (3d ed.; London: Oxford University Press).
- Jokipii, J. R. 1966, *Ap. J.*, **146**, 480.
- . 1967, *ibid.*, **149**, 405.
- . 1968, *ibid.*, **152**, 671.
- . 1971, *Rev. Geophys. and Space Phys.*, **9**, 27.
- . 1972, *Ap. J.*, **172**, 319.
- . 1973, *Proceedings, Solar Terrestrial Relations Conf.*, Calgary, Alberta, Canada, p. 463.
- Jones, F. C., and Birmingham, T. J. 1974, NASA Preprint X-602-74-9, Goddard Space Flight Center, Greenbelt, Md.
- Jones, F. C., Birmingham, T. J., and Kaiser, T. B. 1973, *Ap. J. (Letters)*, **181**, L139.
- Jones, F. C., Kaiser, T. B., and Birmingham, T. J. 1973a, *Phys. Rev. Letters*, **31**, 485.
- . 1973, *Conference Papers, 13th Int. Conf. Cosmic Rays*, **2**, 669.
- Kaiser, T. B. 1973, Tech. Rpt. 74-033, Univ. of Maryland, Dept. of Physics and Astronomy, College Park.
- Kaiser, T. B., Jones, F. C., and Birmingham, T. J. 1973, *Ap. J.*, **180**, 239.
- Klimas, A. J., and Sandri, G. 1971, *Ap. J.*, **169**, 41.
- . 1973a, *ibid.*, **180**, 937.
- . 1973b, *ibid.*, **184**, 955.
- . 1973c, *Conference Papers, 13th Int. Conf. Cosmic Rays*, **2**, 659.
- Noerdlinger, P. D. 1968, *Phys. Rev. Letters*, **20**, 1513.
- Northrop, T. G. 1963, *The Adiabatic Motion of Charged Particles* (New York: Wiley).
- O'Gallagher, J. J. 1967, *Ap. J.*, **150**, 675.
- Quenby, J. J., Balogh, A., Engel, A. R., Elliot, H., Hedgecock, P. C., Hynds, R. J., Sear, J. R. 1969, *Acta Physica Hungaricae*, **29**, 445.
- Volk, H. J. 1973, *Ap. and Space Sci.*, **25**, 471.
- Volk, H., Morfill, G., Alpers, W., and Lee, M. A. 1974, *Ap. and Space Sci.*, **26**, 403.
- Wiener, N. 1933, *The Fourier Integral and Certain of its Applications* (Cambridge: Cambridge University Press; reprinted, New York: Dover), p. 151.

M. L. GOLDSTEIN and A. J. KLIMAS: Code 692/Laboratory for Extraterrestrial Physics, NASA/Goddard Space Flight Center, Greenbelt, MD 20771

G. SANDRI: Aeronautical Research Associates of Princeton, Inc., Princeton, NJ 08540