

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Properties of magnetohydrodynamic turbulence in the solar wind as observed by Ulysses at high heliographic latitudes

B. E. Goldstein and E. J. Smith

Jet Propulsion Laboratory, Pasadena, California

A. Balogh and T. S. Horbury

Imperial College, London, England

M. L. Goldstein and D. A. Roberts

NASA Goddard Space Flight Center, Greenbelt, Maryland

Abstract. The Ulysses mission provides an opportunity to study the evolution of magnetohydrodynamic (MHD) turbulence in pure high-speed solar wind streams. The absence at high heliocentric latitudes of the strong shears in solar wind velocity generally present near the heliocentric current sheet allows investigation of how fluctuations in the magnetic field and plasma relax and evolve in the radially expanding solar wind. We report results of an analysis of the radial and latitudinal variation of the turbulence properties of the fluctuations, especially various plasma-field correlations, in high latitude regions. The results constrain current theories of the evolution of MHD turbulence in the solar wind. Compared to similar observations at 0.3 AU by Helios, we find spectra that are similar in having a large frequency band with an f^{-1} power spectrum in the outward traveling component of the waves, followed at higher frequencies by a steeper spectrum. Ulysses observations establish that at high latitudes the turbulence is less evolved (i.e., has a smaller inertial range) than it is in the ecliptic at the same heliocentric distance, apparently due to the absence of strong velocity shear. Once Ulysses is in the polar coronal hole, properties of the turbulence appear to be determined by the heliocentric distance of the spacecraft rather than by its helio-latitude.

Introduction

Studies of the solar wind during the past two decades have revealed a dynamic and turbulent medium containing hot plasma with twisted magnetic fields (see *Tu and Marsch* [1995] and *Goldstein et al.* [1995a,b]). Understanding how energetic particles propagate through these magnetic fields, requires a description of both the large-scale structure of the solar wind magnetic field and the three-dimensional characteristics of the fluctuating magnetic and velocity fields.

It is now clear that the fluctuating solar wind magnetic and velocity fields are determined in large part by two competing influences: the initial conditions in the solar atmosphere and corona, and the physics controlling the subsequent dynamical evolution in the heliosphere. Conditions in the solar corona are known to be the source of the outward propagating Alfvénic fluctuations observed in the solar wind [*Belcher and Davis*, 1971; *Roberts et al.*, 1987a,b]. Simulations suggest [*Roberts et al.*, 1991; 1992] that, at least near the ecliptic plane, turbulent evolution is driven by velocity shear arising either from the

interaction of fast and slow solar wind streams or by the striated structure of the solar atmosphere [*Woo*, 1995; *Woo and Goldstein*, 1994].

Detailed analyses of plasma and magnetic field data from the Helios and Voyager spacecraft, especially near solar minimum, suggest that in the absence of strong velocity gradients, the evolution of solar wind turbulence is slowed: the degree of "Alfvénicity" of the fluctuations as determined by the correlation between the fluctuating magnetic and velocity fields (the cross helicity) decreases rapidly with distance when velocity gradients are large [*Bavassano and Bruno*, 1989; *Roberts et al.*, 1987a], but can remain large when velocity gradients are small, even out to 8 or 9 AU [*Roberts et al.*, 1987b]. Another indicator of turbulent evolution is that the f^{-1} spectrum which characterizes much of the power spectrum of magnetic field fluctuations in the inner heliosphere at low frequencies extends to higher frequencies at larger heliocentric distances when velocity shears are small [*Roberts et al.*, 1991; 1992]. Conversely, when large velocity shears are present, an inertial range spectrum of the turbulence, which we take to be defined by the $f^{-5/3}$ power spectrum, begins at relatively low frequencies and extends over several decades in frequency.

These analyses have produced a picture of solar wind turbulence which can be tested and refined using data from the unique out-of-the-ecliptic orbit of Ulysses. In addition, several predictions and expectations concerning the nature of the solar wind out of the ecliptic have been described in the literature [*Jokipii and Kóta*, 1989; *Roberts*, 1990; *Goldstein et al.*, 1995a,b]. We report here results of an analysis of the radial and latitudinal variation of the turbulence properties of the solar wind at high latitudes, and compare those results both with Helios data and with previous expectations.

Analyses

In this analysis we used a conjoint plasma-magnetic data set at 4 minute resolution. The magnetic field, which is generally available at significantly higher time resolution than is the plasma data, was sampled at the time of the plasma measurement. The plasma data included the proton and alpha particle densities, the proton velocity, and the Alfvén speed computed including the measured temperature anisotropies. The intervals we chose were characterized by relatively steady conditions ($V_{sw} \approx 783$ and 754 km/s for the high latitude 2 and 4 AU intervals, respectively) and small velocity gradients. The first interval was DOY 299–312, 1993 after Ulysses had encountered the last evidence of corotating stream interactions on its way to high southern latitudes. The spacecraft was near 4 AU and close to -40° southern latitude. The second interval (DOY 229–292, 1994) was during the maximum southern latitude pass. For comparison with in-ecliptic

Copyright 1995 by the American Geophysical Union.

Paper number 95GL03183

0094-8534/95/95GL-03183\$03.00

data that was similarly steady, we used 10 days of Helios data obtained during days 119–129 of 1978 when that spacecraft was near 0.3 AU and the solar wind speed was relatively constant.

The Ulysses data were digitally filtered and decimated to 8 minute time resolution to remove the effects of aliasing apparent in the 4 minute data. The magnetic field data were then converted to velocity units by $\mathbf{b} = \mathbf{B} V_A / B_0$ where V_A is the Alfvén speed determined from local measurements of the solar wind density including alpha-particles, and B_0 is the magnitude of the average magnetic field. Power spectra were computed using fast Fourier transforms.

When dealing with highly Alfvénic turbulence, it is useful to replace the magnetic field and velocity vectors by the Elsässer variables defined by $\mathbf{z}^\pm = \delta\mathbf{v} \pm \delta\mathbf{b}$ [Elsässer, 1950; 1956], where \mathbf{z}^+ and \mathbf{z}^- refer to waves propagating “outward” and “inward” with respect to \mathbf{B}_0 and $\delta\mathbf{v}$ and $\delta\mathbf{b}$ are the fluctuations in the velocity and magnetic fields, respectively. The advantage of using these variables is that for nondissipative and incompressible magnetofluids they are exact solutions of the MHD equations. Because the Reynolds number of the solar wind is a very high and because the wind often behaves quasi-incompressibly, the Elsässer formalism shows the extent to which initially outward propagating Alfvén waves couple to inward propagating fluctuations and other nonAlfvénic disturbances.

Results of the Analyses

In Figure 1a–c we show power spectra of the Elsässer variables during Helios and Ulysses data intervals. In all cases, the power in the outward propagating Alfvénic fluctuations, the \mathbf{z}^+ -power, is the upper curve and the inward propagating Alfvénic fluctuations, the \mathbf{z}^- -power, is the lower curve. The Helios spectrum was constructed from one hour averages of the magnetic field and velocity data and, therefore, only extends to just above 10^{-4} Hz in contrast to the two Ulysses spectra which were constructed from 8 minute resolution data.

The Ulysses high latitude interval and the Helios interval both have much higher Alfvénicity than does the 4 AU interval; i.e. the power in \mathbf{z}^+ greatly exceeds that in \mathbf{z}^- over much of the frequency range. In addition, at low frequencies, the Helios spectra are flat to high frequencies ($>10^{-3}$ Hz), the 2 AU Ulysses spectrum to considerably lower frequency, and the 4 AU spectrum to even lower frequency (also see, *Horbury et al.* [1995a,b]). One should also note that in the vicinity of 10^{-4} Hz, the amplitude of the high latitude \mathbf{z}^+ spectrum at ~2 AU is ~2 times higher than at 4 AU, which is the ratio expected if the fluctuations obey a “WKB” scaling [Heinemann and Olbert, 1980; Roberts, 1990]. Similarly, the Helios \mathbf{z}^+ spectrum is approximately 6 times higher than the 2 AU spectrum over the entire frequency range (Fig-

ure 1c), which again reflects a “WKB-like” scaling ($\langle \delta B^2 \rangle / \rho \propto r^{-3}/r^{-2}$). *Jokipii et al.* [1995], however, report a radial variation more like a quasi-static solution, so that more study is needed before the issue can be resolved.

Other interesting aspects of the differences in Alfvénicity between the 2 AU and 4 AU data are evident from the normalized cross helicity defined by $\sigma_c = 2 H_c / E$, where $H_c = 1/2 \langle \delta\mathbf{v} \cdot \delta\mathbf{b} \rangle$ and E is the magnetic plus kinetic energy. In Figures 2a and b we show $\sigma_c(k)$ for those two intervals. In addition to $\sigma_c(f)$, the plots include $\sigma_{\perp}(f)$ and $\sigma_{\parallel}(f)$, calculated from the radial and transverse components of E and H_c . At 2 AU, $\sigma_c(f)$ is very close to unity (more Alfvénic). Of particular interest is the fact that the low values of $\sigma_c(f)$ below 10^{-6} Hz reflect the sharp decline in $\sigma_{\perp}(f)$. The transverse components $\delta\mathbf{v}_{\perp}$ and $\delta\mathbf{b}_{\perp}$ are still highly correlated, suggesting that quasi-planar Alfvénic fluctuations still dominate below 10^{-6} Hz. Because Alfvénic fluctuations are not expected at such low frequencies [Heinemann and Olbert, 1980], it is possible that spacecraft motion across field lines in the corotating frame is aliasing high frequency Alfvénic fluctuations into the observed time series at lower frequencies. The high Alfvénicity in the transverse components at large scales implies that the convergence of \mathbf{z}^+ and \mathbf{z}^- spectra at low frequencies is due entirely to variations in the radial velocity sampled as Ulysses crosses field lines. By 4 AU $\delta\mathbf{v}_{\perp}$ and $\delta\mathbf{b}_{\perp}$ are relatively uncorrelated, reflecting a general decay in Alfvénicity with distance across the entire frequency band.

Another quantity from which turbulent evolution can be deduced is the spectrum of the Alfvén ratio, defined as the ratio r_A of kinetic energy to magnetic energy. The importance of this quantity arises from the Alfvén effect first described by *Kraichnan* [1965]. Kraichnan’s prediction was that the magnetic and kinetic energy (per unit wave number) in an incompressible turbulent magnetofluid should be equal, on average, in the inertial range. As shown in Figure 3, r_A is approximately 1/2 for both the 2 and 4 AU intervals. For reasons not understood, the Alfvén ratio in the solar wind rarely equals 1 as predicted. The component energy ratios plotted in Figure 3 also show that $r_{A\perp}$ remains ~1/2 back to below 10^{-6} Hz, consistent with the high Alfvénicity of $\delta\mathbf{v}_{\perp}$ and $\delta\mathbf{b}_{\perp}$ at those scales.

Discussion and Conclusions

The most obvious characteristic of the power spectra of the high latitude Ulysses data is the relatively slow turbulent evolution evident, especially when compared with in-ecliptic observations at comparable radial distances. Whereas by 2 AU in the ecliptic the cross helicity generally shows a significant reduction from its highest values [Roberts et al. 1987a,b], the values measured by Ulysses are ~0.8




Figure 1. Trace of the power spectrum of the outward propagating (\mathbf{z}^+ , upper curve) and inward propagating (\mathbf{z}^- , lower curve) Alfvénic fluctuations; (a) for a 13 day period while Ulysses was at approximately 4 AU; (b) for a 63 day period while Ulysses was at high southern latitudes and a radial distance of approximately 2 AU; and, (c), similar to (a), but for a 10-day period during 1978 when Helios was near 0.3 AU.




Figure 2. The normalized cross helicity $\sigma_c(f)$ calculated from the trace of the power spectral densities of z^\pm and the radial and transverse normalized cross helicities computed from the radial and transverse components of z^\pm , (a) for the 2 AU Ulysses interval, and (b), for the 4 AU Ulysses interval.

(Figure 2a). That a significant percentage of the fluctuations between 10^{-5} – 10^{-3} Hz are outward propagating and highly Alfvénic, is also evident from Figure 1b. The low frequency results for $\sigma_{c\perp}(f)$ and $r_{A\perp}(f)$ shown in Figures 2 and 3 suggest that longitudinal sampling of Alfvénic fluctuations as Ulysses crosses field lines may account for the apparent presence of such fluctuations at frequencies well below the regime in which they are thought to propagate. By 4 AU, however, the Alfvénicity of the transverse fluctuations in the 10^{-6} – 10^{-5} Hz band has decreased markedly, suggesting dynamical evolution.

Another indication of the slow rate of turbulent evolution in the high latitude Ulysses data is that the f^{-1} portion of the spectrum, which dominates the low wave number, large spatial scales, extends to relatively high frequencies ($\sim 10^{-4}$ Hz) (Figure 1b), whereas, in the ecliptic near 1 AU, the f^{-1} spectrum ends closer to 10^{-5} Hz [Matthaeus and Goldstein, 1982b], except in the most Alfvénic regions. The frequency at which the spectral breakpoint changes from f^{-1} to the $f^{-5/3}$ moves to progressively lower frequencies with increasing heliocentric distance (see also Horbury *et al.* [1995c]). This evolution is more rapid in the inner heliosphere as can be seen most clearly in Helios data [Bavassano *et al.*, 1982]. Beyond 1 AU the evolution in the breakpoint is slower and less obvious [Burlaga and Goldstein, 1984; Goldstein *et al.*, 1984; Klein *et al.*, 1992].

Evidence for turbulent evolution by 2 AU is found in the spectral changes seen in Figure 1. Also, highly Alfvénic Helios data do show $r_A(f) \approx 1$ (as predicted by Kraichnan) near 0.3 AU, but $r_A(f) \approx 0.5$ by 1 AU [Goldstein *et al.*, 1995a], suggesting evolution from a nearly equipartitioned state to one in which magnetic energy dominates. Equipartition has been reported in two-dimensional simulations [Fyfe and Montgomery, 1976], but solar wind observations most often show inertial range values of $r_A(f)$ close to 0.5 [Roberts *et al.*, 1987a,b; 1990; 1992; Matthaeus and Goldstein 1982a; Roberts, 1992]. Goldstein *et al.* [1995c] have suggested that the pressure anisotropy of pickup ions may act in some circumstances to reduce r_A .

Comparison of the intensities of the power spectra in Figure 1 at 0.3 (Helios) and Ulysses at 2 and 4 AU suggests that the scaling of the z^\pm amplitudes with distance satisfies WKB at 10^{-6} Hz, *i.e.*, the power is expected to decrease as $1/r$ [Heinemann and Olbert, 1980] for the magnetic field in Alfvén speed units. The observed change in power between 2 and 4 AU is approximately a factor of 2. While this dependence is generally expected in the inertial range, Jokipii and Kóta [1989], noting that the polar magnetic fields would be radial, predicted no fall-off with radial distance of B (in Alfvén speed units) while Roberts [1990] argued that the effects of turbulence would cause the variation with distance to follow the WKB-scaling. The linear




Figure 3. The spectrum of $r_A(f)$ calculated from the trace of the power spectral densities of z^\pm and the radial and transverse ratios of kinetic and magnetic energy computed from the radial and transverse components of z^\pm , (a), for the 2 AU Ulysses interval, and (b), for the 13 day interval while Ulysses was at approximately 4 AU.

scaling argument of Jokipii and Kóta [1989], when applied to velocity fluctuations, predicts that r_A would approach 0 as $1/r^2$ ($\rho \delta v^2 \propto 1/r^4$), which is not observed. Jokipii et al. [1995] have recently analyzed the variation in the variance of the magnetic field between 1.5–4 AU and concluded that it fits their expectations. The addition of data from 1.5–2 AU, which is not included here, contributes significantly to their conclusions. Work in progress by Balogh et al. (private communication, 1995) on the radial dependence of the low frequency fluctuations finds a low frequency variation for the variance of roughly $r^{-2.2}$ (preliminary results) at frequencies of about 3×10^{-6} Hz (i.e., slower decrease than WKB). The variations of and correlations with velocity were not addressed in these works. The apparent difference in radial variation at low frequencies between our results and those just quoted requires further investigation.

We conclude that the observed evolution of the magnetic and velocity fields between 2 and 4 AU at high latitudes appears consistent with previous expectations derived from in-ecliptic studies and numerical simulations [Roberts, 1990; Roberts et al., 1991; 1992]. The turbulence is less evolved than at corresponding distances in the ecliptic, probably because large-scale velocity gradients are absent. The results below 10^{-5} Hz may be due to sampling of features in longitude; the extent to which spatial versus temporal fluctuations contribute in this frequency regime needs further investigation.

Acknowledgements. Work conducted at the Jet Propulsion Laboratory and the Goddard Space Flight Center was supported by the NASA; work at Imperial College was supported by the U.K. SERC.

References

- Bavassano, B., and R. Bruno, Large-scale solar wind fluctuations in the inner heliosphere at low solar activity, *J. Geophys. Res.*, **94**, 168, 1989.
- Bavassano, M., M. Dobrowolny, G. Fanfoni, F. Mariani, and N. F. Ness, Statistical properties of MHD fluctuations associated with high-speed streams from Helios-2 observations, *Solar Phys.*, **78**, 373, 1982.
- Belcher, J. W., and L. Davis, Large-amplitude Alfvén waves in the interplanetary medium, *2, J. Geophys. Res.*, **76**, 3534, 1971.
- Burlaga, L. F., and M. L. Goldstein, Radial variations of large-scale magnetohydrodynamic fluctuations, *J. Geophys. Res.*, **89**, 6813, 1984.
- Elsässer, W. M., The hydromagnetic equations, *Phys. Rev.*, **79**, 183, 1950.
- Elsässer, W. M., Hydromagnetic dynamo theory, *Rev. Mod. Phys.*, **18**, 135, 1956.
- Fyfe, D., and D. Montgomery, High beta turbulence in two-dimensional magnetohydrodynamics, *J. Plasma Phys.*, **29**, 181, 1976.
- Goldstein, M. L., L. F. Burlaga, and W. H. Matthaeus, Power spectral signatures of interplanetary corotating and transient flows, *J. Geophys. Res.*, **89**, 3747, 1984.
- Goldstein, M. L., D. A. Roberts, and W. H. Matthaeus, Magnetohydrodynamic turbulence in cosmic winds, in *Cosmic Winds and the Heliosphere*, edited by J. R. Jokipii, and C. P. Sonett, in press, Univ. of Arizona Press, Tucson, 1995a.
- Goldstein, M. L., D. A. Roberts, and W. H. Matthaeus, Magnetohydrodynamic turbulence in the solar wind, in *Annual Review of Astronomy and Astrophysics*, in press, 1995b.
- Goldstein, B. E., M. Neugebauer, and E. J. Smith, Alfvén waves, alpha particles, and pickup ions in the solar wind, *Geophys. Res. Lett.*, in this issue, 1995c.
- Heinemann, M., and S. Olbert, Non-WKB Alfvén waves in the solar wind, *J. Geophys. Res.*, **85**, 1311, 1980.
- Horbury, T. S., A. Balogh, and R. J. Forsyth, Magnetic field signatures of unevolved turbulence in solar polar flows, *J. Geophys. Res.*, **100**, in press, 1995a.
- Horbury, T. S., A. Balogh, R. J. Forsyth, and E. J. Smith, Ulysses magnetic field observations of fluctuations within polar coronal flows, *Annales Geophysicae*, **13**, 105, 1995b.
- Horbury, T. S., A. Balogh, R. J. Forsyth, and E. J. Smith, Observations of evolving turbulence in the polar solar wind, *Geophys. Res. Lett.*, **22**, in this issue, 1995c.
- Jokipii, J. R., and J. Kóta, The polar heliospheric magnetic field, *Geophys. Res. Lett.*, **16**, 1, 1989.
- Jokipii, J. R., J. Kóta, J. Giacalone, T. S. Horbury, and E. J. Smith, Interpretation and consequences of large-scale magnetic variances observed at high heliographic latitude, *Geophys. Res. Lett.*, **22**, in this issue, 1995.
- Klein, L. W., W. H. Matthaeus, D. A. Roberts, and M. L. Goldstein, Evolution of spatial and temporal correlations in the solar wind: Observations and interpretation, in *Proceedings of Solar Wind 7, COSPAR Collq. Ser.*, edited by E. Marsch, and R. Schwenn, p. 197, Pergamon, Oxford, Goslar, Germany, 1992.
- Kraichnan, R. H., Inertial range of hydromagnetic turbulence, *Phys. Fluids*, **8**, 1385, 1965.
- Matthaeus, W. H., and M. L. Goldstein, Measurement of the rugged invariants of magnetohydrodynamic turbulence, *J. Geophys. Res.*, **87**, 6011, 1982a.
- Matthaeus, W. H., and M. L. Goldstein, Stationarity of magnetohydrodynamic fluctuations in the solar wind, *J. Geophys. Res.*, **87**, 10347, 1982b.
- Roberts, D. A., Turbulent polar heliospheric fields, *Geophys. Res. Lett.*, **17**, 567, 1990.
- Roberts, D. A., Observation and simulation of the radial evolution and stream structure of solar wind turbulence, in *Proceedings of Solar Wind 7, COSPAR Collq. Ser.*, vol. 3, edited by E. Marsch, and R. Schwenn, p. 533, Pergamon, New York, 1992.
- Roberts, D. A., S. Ghosh, M. L. Goldstein, and W. H. Matthaeus, MHD simulation of the radial evolution and stream structure of solar wind turbulence, *Phys. Rev. Lett.*, **67**, 3741, 1991.
- Roberts, D. A., M. L. Goldstein, and L. W. Klein, The amplitudes of interplanetary fluctuations: Stream structure, heliocentric distance, and frequency dependence, *J. Geophys. Res.*, **95**, 4203, 1990.
- Roberts, D. A., M. L. Goldstein, L. W. Klein, and W. H. Matthaeus, Origin and evolution of fluctuations in the solar wind: Helios observations and Helios-Voyager comparisons, *J. Geophys. Res.*, **92**, 12023, 1987a.
- Roberts, D. A., L. W. Klein, M. L. Goldstein, and W. H. Matthaeus, The nature and evolution of magnetohydrodynamic fluctuations in the solar wind: Voyager observations, *J. Geophys. Res.*, **92**, 11021, 1987b.
- Roberts, D. A., M. L. Goldstein, W. H. Matthaeus, and S. Ghosh, Velocity shear generation of solar wind turbulence, *J. Geophys. Res.*, **97**, 17115, 1992.
- Tu, C.-Y., and E. Marsch, *MHD Structures, Waves and Turbulence in the Solar Wind: Observations and Theories*, Kluwer Academic, Boston, 1995.
- Woo, R., Solar wind speed structure in the inner corona at 3–12 R_\odot , *Geophys. Res. Lett.*, **22**, 1393, 1995.
- Woo, R., and R. Goldstein, Latitudinal variation of speed and mass flux in the acceleration region of the solar wind inferred from spectral broadening measurements, *J. Geophys. Res.*, **21**, 85, 1994.

Received: June 29, 1995; revised: September 11, 1995;
accepted September 28, 1995