

This work is on a Creative Commons Attribution 3.0 Unported (CC BY 3.0) license, <https://creativecommons.org/licenses/by/3.0/>. Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing [scholarworks-group@umbc.edu](mailto:scholarworks-group@umbc.edu) and telling us what having access to this work means to you and why it's important to you. Thank you.



PAPER • OPEN ACCESS

## Geometric quantum speed limits: a case for Wigner phase space

To cite this article: Sebastian Deffner 2017 *New J. Phys.* **19** 103018

View the [article online](#) for updates and enhancements.

### Related content

- [Quantum speed limits: from Heisenberg's uncertainty principle to optimal quantum control](#)
- [The initial-state dependence of the quantum speed limit](#)
- [Detecting quantum speedup in closed and open systems](#)

### Recent citations

- [Kibble-Zurek scaling in quantum speed limits for shortcuts to adiabaticity](#)  
Ricardo Puebla *et al*
- [Effects of coherence on quantum speed limits and shortcuts to adiabaticity in many-particle systems](#)  
Tian-Niu Xu *et al*
- [Quantum speed limit via the trajectory ensemble](#)  
Xianghong Hu *et al*



## PAPER

## Geometric quantum speed limits: a case for Wigner phase space

## OPEN ACCESS

RECEIVED  
11 April 2017REVISED  
2 June 2017ACCEPTED FOR PUBLICATION  
3 August 2017PUBLISHED  
20 October 2017Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/4.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Sebastian Deffner

Department of Physics, University of Maryland Baltimore County, Baltimore, MD 21250, United States of America

E-mail: [deffner@umbc.edu](mailto:deffner@umbc.edu)

Keywords: quantum speed limit, Heisenberg uncertainty, Wigner function

## Abstract

The quantum speed limit is a fundamental upper bound on the speed of quantum evolution. However, the actual mathematical expression of this fundamental limit depends on the choice of a measure of distinguishability of quantum states. We show that quantum speed limits are qualitatively governed by the Schatten- $p$ -norm of the generator of quantum dynamics. Since computing Schatten- $p$ -norms can be mathematically involved, we then develop an alternative approach in Wigner phase space. We find that the quantum speed limit in Wigner space is fully equivalent to expressions in density operator space, but that the new bound is significantly easier to compute. Our results are illustrated for the parametric harmonic oscillator and for quantum Brownian motion.

## 1. Introduction

It has recently been argued that already the first generation of real-life quantum computers will be able to perform certain tasks exponentially faster than classical computers [1]. This so-called ‘quantum supremacy’ [2] rests in the fact that loosely speaking quantum state space is exponentially larger than the classical computational space, and hence significantly less operations are necessary to perform the same computation. However, the working principles of quantum computers and classical computers are fundamentally different, which makes it not immediately clear how to quantify the ‘quantum speed-up’ [3]. To make matters even more involved, in the theory of quantum computation ‘time’ is not actually a physical time, but rather a synonym for the ‘number of computational operations’ [3]. The more practical question is, however, how fast a quantum computer could actually operate.

To address this issue a somewhat opposite approach has been developed in quantum dynamics, where the notion of a *quantum speed limit* has found wide-spread prominence. Whereas in the theory of quantum computation one is after characterizing quantum speed-ups—the quest for faster and faster computations with less and less single operations—the quantum speed limit sets the ultimate, maximal speed with which any quantum system can evolve. This means, in particular, that every single quantum operation takes a finite, minimal time to be accomplished—and thus even quantum computers will not be able to achieve any arbitrary speed-ups. This quantum speed limit originates in the Heisenberg indeterminacy principle [4, 5],  $\Delta E \Delta t \gtrsim \hbar/2$ . However, more than just being another expression of quantum indeterminacy the quantum speed limit is a fundamental property of quantum dynamics as highlighted by famous debates between Einstein and Bohr [6].

The first rigorous treatment was proposed by Mandelstam and Tamm [7], who showed that the minimal time a quantum system needs to evolve between orthogonal states is bounded from below by the variance of the energy,  $\tau_{\text{QSL}} = \pi \hbar / 2 \Delta E$ , where  $\Delta E = (\langle H^2 \rangle - \langle H \rangle^2)^{1/2}$ . Since, however, the variance of an operator is not necessarily a good quantifier for dynamics [8], Margolis and Levitin [9] revisited the problem and derived a second bound on the quantum evolution time in terms of the average energy  $E = \langle H \rangle - E_g$  over the ground state with energy  $E_g$ ,  $\tau_{\text{QSL}} = \pi \hbar / 2 E$ . It was eventually realized that these two bounds are not independent, and that only the unified bound is tight [10].

Nowadays, it has been established in virtually all areas of quantum physics [11–16] that the quantum speed limit [17–20], sets a fundamental upper bound on the speed of any quantum dynamics. In particular in recent

years, the quantum speed limit has been extensively studied and generalized for isolated [21–26] and open [27–39] quantum systems. The renewed and concentrated interest was inspired by three letters [27–29], which broadened the scope of the quantum speed limit beyond unitary dynamics. In particular, [27] showed that the maximal speed of quantum evolution is given by the time averaged norm of the generator of the dynamics [27], which only in the case of unitary dynamics and for orthogonal states reduces to the average energy  $E$  [25].

All treatments of the quantum speed limit have in common that the analyses start with a *choice* of a measure of the distinguishability of quantum states. For instance, [28] studies the relative purity, [27] starts with the Bures angle between initially pure states and time-evolved states, and [40] even defines an entirely new metric. Finally, Pires *et al* [41] derived a whole family of quantum speed limits, which is based on a family of contractive Riemannian metrics.

Thus, two natural questions arise: (i) are all of these treatments of quantum speed independent, or can the quantum speed limit be universally characterized—independently of the chosen measure of distinguishability? (ii) Most of the expressions for the quantum speed limit are mathematically rather involved. Hence, can one find a mathematically simple bound that is computable and nevertheless captures the universal behavior?

In the following we will answer both questions. First, we will show that the quantum speed is universally governed by any Schatten- $p$ -norm of the generator of the quantum dynamics. In the second part of the analysis, we will derive a new quantum speed limit in terms of the Wasserstein norm of the rate of change of the Wigner function. We will argue that the quantum speed limit in Wigner phase space captures the same qualitative behavior as the speed limits derived in density operator space. However, we will also see that the new quantum speed limit is significantly easier to compute, for pure as well as mixed states, and for isolated as well as open dynamics. As an illustrative example we will discuss the semi-classical, high temperature limit, and we will confirm that the quantum speed limit is a pure quantum feature, i.e., that classical systems do not experience a fundamental bound on their rates of change.

## 2. Quantum speed and the geometric approach

We begin by briefly reviewing the main results of [27] and by establishing notions and notations. Consider a quantum master equation,

$$\dot{\rho}_t = L_t(\rho_t), \quad (1)$$

where the dot denotes a derivative with respect to time. For isolated systems the Liouvillian superoperator,  $L_t$ , reduces to the von-Neumann equation,  $L_t(\rho) = [H_t, \rho]/i\hbar$ , but we explicitly allow for any open systems dynamics—Markovian as well as non-Markovian. Note that [27] theoretically predicted that non-Markovian environments can speed up quantum dynamics. This was experimentally verified in cavity QED [35].

In geometric quantum mechanics [42] it has proven useful to quantify the distinguishability of quantum states in terms of the Bures angle [43],

$$\mathcal{L}(\rho_0, \rho_t) = \arccos(\sqrt{F(\rho_0, \rho_t)}) = \arccos(\text{tr}\{\sqrt{\sqrt{\rho_0} \rho_t \sqrt{\rho_0}}\}), \quad (2)$$

where we further introduced the quantum fidelity  $F(\rho_0, \rho_t)$  [44]. To obtain an upper bound on the speed of evolution one then considers the magnitude of the geometric speed,  $|\dot{\mathcal{L}}|$ , and it is easy to see that we have [27]

$$2 \cos(\mathcal{L}) \sin(\mathcal{L}) \dot{\mathcal{L}} \leq |\dot{F}(\rho_0, \rho_t)|. \quad (3)$$

For initially pure states,  $\rho_0 = |\psi_0\rangle\langle\psi_0|$ , equation (2) can be further simplified and it can be shown that [27]

$$2 \cos(\mathcal{L}) \sin(\mathcal{L}) \dot{\mathcal{L}} \leq |\langle\psi_0|\dot{\rho}_t|\psi_0\rangle| \leq \min\{\|\dot{\rho}_t\|_p \text{ for } p \in \{1, 2, \infty\}\}. \quad (4)$$

Here  $\|A\|_p$  denotes the Schatten- $p$ -norm of an operator,  $O$ , which is defined as

$$\|O\|_p \equiv (\text{tr}\{|O|^p\})^{1/p} = \left(\sum_k o_k^p\right)^{1/p} \quad (5)$$

and the  $o_k$  are the singular values of  $O$ , i.e. the eigenvalues of the Hermitian operator  $|O| \equiv \sqrt{O^\dagger O}$ . The more familiar trace, Hilbert–Schmidt, and operator norms correspond respectively to  $p = 1, 2$ , and  $\infty$ . Equation (4) can then be used to define the quantum speed limit  $v_{\text{QSL}}$ ,

$$\dot{\mathcal{L}} \leq \frac{v_{\text{QSL}}}{2 \cos(\mathcal{L}) \sin(\mathcal{L})} \equiv \frac{1}{2 \cos(\mathcal{L}) \sin(\mathcal{L})} \min\{\|\dot{\rho}_t\|_p \text{ for } p \in \{1, 2, \infty\}\}. \quad (6)$$

Note that in contrast to [16] we did not include the denominator into the definition of  $v_{\text{QSL}}$ . The reason for this choice will become obvious shortly.

Although useful for theoretical predictions of experimental outcomes [35] equation (4) also left several questions unaddressed. Probably the most immediate one is, how the above treatment would have to be

generalized to initially mixed states. Generally this is a mathematically involved problem, since the quantum fidelity,  $F(\rho_0, \rho_t)$ , and its derivatives are non-trivial to handle. A comprehensive analysis of this issue was proposed by Pires *et al* [41]. As proven by the Morozova–Čencov–Petz theorem, there is an infinite number of contractive metrics, which are equivalent for pure states. One particular case with an interesting analytic expression is the Wigner–Yanase information,  $A(\rho_0, \rho_t)$ ,

$$\mathcal{L}^{\text{WY}}(\rho_0, \rho_t) = \arccos(A(\rho_0, \rho_t)) = \arccos(\text{tr}\{\sqrt{\rho_0}\sqrt{\rho_t}\}), \quad (7)$$

which can be used to derive quantum speed limits instead of the Bures angle (2). Pires *et al* [41] then analyze an infinite family of bounds to identify computable bounds. However, similarly to equation (6) the quantum speed is bounded by a norm of the dynamics,  $\|\dot{\rho}_t\|$ , and only the ‘prefactor’ depends on the choice of the starting point, whether it be equation (2) or (7). Thus, the analysis of [41] makes the second question even more obvious: namely, formulating quantum speed limits seems to be somewhat arbitrary, since every single treatment starts with a *choice* of a measure of distinguishability in order to define the geometric speed. Hence, it would be desirable to find a universal measure of quantum speed, and define the quantum speed limit exclusively in terms of this measure.

Both, [27] and [41] make the strong case that the choice has to be a contractive, Riemannian metric on quantum state space. This choice is justified if one would like to find tight bounds on the quantum speed [27]. If one is only interested in the qualitative behavior, however, other measures might be more convenient to work with. For instance, instead of choosing the Bures angle,  $\mathcal{L}(\rho_0, \rho_t)$ , we also could have worked with the Bures distance [42],

$$\mathcal{L}^{\text{D}}(\rho_0, \rho_t) = \sqrt{2(1 - \sqrt{F(\rho_0, \rho_t)})}. \quad (8)$$

One easily convinces oneself that a such defined geometric speed,  $\dot{\mathcal{L}}^{\text{D}}$ , leads to an inequality similar to equation (3). Therefore, we observe already here that all of these treatments have in common that eventually the quantum speed is characterized by a Schatten- $p$ -norm,  $\|\dot{\rho}_t\|_p = \|L(\rho_t)\|_p$ .

### 3. Quantum speed in density operator space

We have seen above that typically quantum speed is characterized by the dynamical behavior of the quantum fidelity,  $F(\rho_0, \rho_t)$ . Since  $F(\rho_0, \rho_t)$  is mathematically rather involved several continuity bounds have been derived. For instance, we have [42]

$$1 - \sqrt{F(\rho_0, \rho_t)} \leq \frac{1}{2}\mathcal{L}_1(\rho_t, \rho_0) \leq \sqrt{1 - F(\rho_0, \rho_t)}, \quad (9)$$

where  $\mathcal{L}_1$  denotes the trace distance, i.e., the Schatten-1-distance

$$\mathcal{L}_1(\rho_t, \rho_0) = \|\rho_t - \rho_0\|_1 \equiv \text{tr}\{|\rho_t - \rho_0|\}. \quad (10)$$

The obvious question is whether a quantum speed limit can be derived starting with the trace distance  $\mathcal{L}_1$ . Note again that the here considered quantum speed is precisely the geometric speed as given by the rate of change of the distance,  $\dot{\mathcal{L}}_1$ , which is bounded by the above defined quantum speed limit  $v_{\text{QSL}}$  (6). Therefore, we consider

$$\dot{\mathcal{L}}_1(\rho_t, \rho_0) = \text{tr}\{|\rho_t - \rho_0|^{-1}(\rho_t - \rho_0)\dot{\rho}_t\}, \quad (11)$$

which can be bounded from above by with the triangle inequality for operators,  $|\text{tr}\{O\}| \leq \text{tr}\{|O|\}$ , as

$$\dot{\mathcal{L}}_1(\rho_t, \rho_0) \leq |\dot{\mathcal{L}}_1(\rho_t, \rho_0)| \leq \text{tr}\{|\dot{\rho}_t|\} = \|\dot{\rho}_t\|_1. \quad (12)$$

We immediately conclude that whether we choose the Bures angle (2) or the trace distance (9) only determines the functional dependence of the geometric speed on the choice of the metric (4). The dynamics and, hence, the actual quantum speed limit, however, is fully characterized by the trace norm of the rate with which the quantum state changes.

We can go even one step further, and consider any Schatten- $p$ -norm as a starting point of the derivation,

$$\mathcal{L}_p(\rho_t, \rho_0) = \|\rho_t - \rho_0\|_p \equiv (\text{tr}\{|\rho_t - \rho_0|^p\})^{\frac{1}{p}}, \quad (13)$$

for which the geometric speed is bounded from above by

$$\dot{\mathcal{L}}_p(\rho_t, \rho_0) \leq \|\dot{\rho}_t\|_p. \quad (14)$$

A proof of the latter result can be found in appendix A. In conclusion we have that the actual quantum speed limit is given by

$$v_{\text{QSL}} \equiv \min\{\|\dot{\rho}_t\|_p \text{ for } p \in [0, \infty)\}. \quad (15)$$

Equation (15) constitutes our first main result. Derivations of geometric quantum speed limits depend on a rather arbitrary, although well-motivated choice of a measure of distinguishability of quantum states. The final expression will be functionally dependent on this choice, see for instance [28] for the relative purity, [27] for the Bures angle, and [41] for the Wigner–Yanase information. However, since all of these measures fulfill continuity inequalities [42] (see also equation (9)), the actual quantum speed limit  $v_{\text{QSL}}$  is given by the smallest Schatten- $p$ -norm.

We would like to emphasize that equation (15) only qualitatively characterizes the quantum dynamics. Generally, the maximal speed of a quantum process depends on the (i) initial state, (ii) the dynamical manifold, and (iii) the kind of quantum dynamics [45]. State independent bounds could now be obtained by, for instance, maximizing  $v_{\text{QSL}}$  over all possible initial states. Since this, however, poses a formidable mathematical task, we continue by deriving a mathematical more tractable bound on the quantum speed.

#### 4. Quantum speed in Wigner phase space

The expression of the quantum speed limit,  $v_{\text{QSL}}$  in equation (15), is a powerful expression that can be used to obtain physical insight into the dynamical properties of quantum systems [35]. However, computing the tightest bound, i.e., the operator norm [27, 46] is far from being a trivial task. Imagine, for instance, we want to study a driven, open quantum system such as in quantum Brownian motion [47]. In this case, the dynamics is typically solved in a computationally convenient and continuous basis [48]. Extracting the singular values from such a representation of the time-dependent density operator is computationally expensive, if it is at all feasible. Thus, it would be desirable to find an alternative expression for  $v_{\text{QSL}}$  which gives the same qualitative information, but which is also much easier to compute.

Especially in the treatment of open quantum systems [48] as well as to study the semi-classical limit [49] it has proven useful to express quantum states in their Wigner representation

$$W(x, p) = \frac{1}{\pi\hbar} \int dy \langle x + y | \rho | x - y \rangle \exp\left(-\frac{2ip y}{\hbar}\right). \quad (16)$$

If we want to derive a quantum speed limit in Wigner phase space, we now need to *choose* a measure of distinguishability. To this end, consider the *total variation distance*, which is given by the Wasserstein-1-distance [50],

$$\mathcal{D}(W_t, W_0) = \|W_t - W_0\|_1 \equiv \int d\Gamma |W(\Gamma, t) - W_0(\Gamma)|, \quad (17)$$

with  $\Gamma = (x, p)$ . The Wasserstein-1-distance can be regarded as a generalization of the trace distance to (semi-) probability distributions.

In complete analogy to above, we now consider the geometric speed

$$\dot{\mathcal{D}}(W_t, W_0) = \int d\Gamma \frac{W(\Gamma, t) - W_0(\Gamma)}{|W(\Gamma, t) - W_0(\Gamma)|} \dot{W}(\Gamma, t) \quad (18)$$

which can again be bounded with the help of the triangle inequality

$$\dot{\mathcal{D}}(W_t, W_0) \leq |\dot{\mathcal{D}}(W_t, W_0)| \leq \int d\Gamma |\dot{W}(\Gamma, t)|. \quad (19)$$

Comparing equations (12) and (19) we immediately see that we have obtained an analogous expression for the quantum speed limit,  $v_{\text{QSL}}$ , in Wigner space. In appendix B we show that if we consider the more general case of any Wasserstein- $p$ -distance [50],

$$\mathcal{D}_p(W_t, W_0) = \|W_t - W_0\|_p \equiv \left( \int d\Gamma |W(\Gamma, t) - W_0(\Gamma)|^p \right)^{1/p}, \quad (20)$$

we find

$$\dot{\mathcal{D}}_p(W_t, W_0) \leq \|\dot{W}_t\|_p. \quad (21)$$

Hence we conclude that we have for the quantum speed limit in phase space  $v_{\text{QSL}}^W$ ,

$$v_{\text{QSL}}^W \equiv \min \{ \|\dot{W}_t\|_p \text{ for } p \in [0, \infty) \}. \quad (22)$$

Equation (22) has the same functional form as the quantum speed limit derived in density operator space (15). However, equation (22) is significantly easier to compute, since it only involves the absolute value of an imaginary valued function, instead of the singular values of a high-dimensional operator.

What remains to verify is that  $v_{\text{QSL}}$  (15) and  $v_{\text{QSL}}^W$  (22) contain the same physical information and that  $v_{\text{QSL}}$  and  $v_{\text{QSL}}^W$  behave qualitatively similarly.

## 5. Qualitative comparison of the two approaches

In a mathematical sense, the Weyl–Wigner transform (16) is a well-defined, invertible integral transform between the phase-space and operator representations of quantum states [51]. Therefore, one would expect  $v_{\text{QSL}}$  (15) and  $v_{\text{QSL}}^W$  (22) to be fully equivalent.

That this is, indeed, the case we will now illustrate by computing  $v_{\text{QSL}}$  (15) and  $v_{\text{QSL}}^W$  (22) for a solvable example. For the sake of simplicity we restrict ourselves to unitary dynamics induced by the parametric harmonic oscillator with Hamiltonian,

$$H = \frac{P^2}{2M} + \frac{1}{2}M\omega_t^2 x^2. \quad (23)$$

It can be shown that the dynamics is fully analytically solvable [52]. For systems initially starting in the ground state,  $\rho_0 = |\psi_0\rangle\langle\psi_0|$ , with

$$\psi_0(x) = \left(\frac{M\omega_0}{\pi\hbar}\right)^{1/4} \exp\left(-\frac{m\omega_0 x^2}{2\hbar}\right) \quad (24)$$

the time-dependent density operator can be written as [53]

$$\begin{aligned} \rho(x, y, t) = & \sqrt{\frac{M\omega_0}{\pi\hbar} \frac{1}{Y_t^2 + \omega_0^2 X_t^2}} \\ & \times \exp\left(-\frac{M\omega_0}{2\hbar} \frac{1}{Y_t^2 + \omega_0^2 X_t^2} [x^2 + y^2 + i(x^2 - y^2)(\omega_0^2 \dot{X}_t X_t + \dot{Y}_t Y_t)]\right). \end{aligned} \quad (25)$$

Here,  $X_t$  and  $Y_t$  are the solutions of the force free harmonic oscillator,  $\ddot{X}_t + \omega_t^2 X_t = 0$ , with the boundary conditions  $X_0 = 0$ ,  $\dot{X}_0 = 1$ ,  $Y_0 = 1$ , and  $\dot{Y}_0 = 0$ .

It is then a simple exercise to numerically obtain the quantum speed limits,  $v_{\text{QSL}}$  (15) and  $v_{\text{QSL}}^W$  (22). Without loss of generality, we computed  $v_{\text{QSL}}$  (15) from the trace norm (12), and  $v_{\text{QSL}}^W$  (22) from the Wasserstein-1-norm (19). This is sufficient, since for pure states both the Schatten- $p$ -norms as well as the Wasserstein- $p$ -norms are monotonic in  $p$  [46].

Specifically,  $v_{\text{QSL}}$  can be obtained from a numerical singular value decomposition of  $\rho(x, y, t)$  (25), which is a computationally rather expensive task. On the other hand,  $v_{\text{QSL}}^W$  is obtained directly from the corresponding Wigner function (16). In figure 1 we plot the numerical results for a linear quench,  $\omega_t^2 = \omega_0^2 - (\omega_0^2 - \omega_1^2)t/\tau$ , for several values of the quench time  $\tau$ . For the ease of comparison, we further normalized the quantum speed limits  $v_{\text{QSL}}$  (15) and  $v_{\text{QSL}}^W$  (22) by their maximal value during the time interval  $t \in [0, \tau]$ . We observe perfect agreement of  $v_{\text{QSL}}$  (15) and  $v_{\text{QSL}}^W$  (22). Hence, we conclude that our initial expectation is indeed verified, namely we find that  $v_{\text{QSL}}$  and  $v_{\text{QSL}}^W$  are fully equivalent. In particular for the present case  $v_{\text{QSL}}$  and  $v_{\text{QSL}}^W$  only differ by a factor that is determined by their maximal value during the quench time  $\tau$ . It is worth emphasizing again that the quantum speed limit is significantly easier to compute in Wigner phase space, since the computationally expensive task of having to determine the singular values can be fully avoided.

## 6. Quantum speed in the semiclassical limit



We conclude the analysis by briefly studying the high-temperature, semi-classical limit. Within the geometric approach to quantum speed it often proves useful to define the quantum speed limit time,  $\tau_{\text{QSL}}$ , which is given by the inverse of the time-averaged quantum speed limit,  $v_{\text{QSL}}$ , [27]. Note that  $\tau_{\text{QSL}}$  is *not* a physical time, but rather a characteristic of the internal dynamics [7, 9].

We can thus define the quantum speed limit time in Wigner space as

$$\tau_{\text{QSL}}^W \equiv \frac{\mathcal{D}(W_t, W_0)}{1/\tau \int_0^\tau dt v_{\text{QSL}}^W}. \quad (26)$$

For unitary dynamics it is easy to see that  $\tau_{\text{QSL}}^W$  is proportional to  $\hbar$ , and hence,  $\tau_{\text{QSL}}^W$  can be understood as an expression of the Heisenberg indeterminacy principle for energy and time. For open systems, however, the interpretation is less obvious [27]. Since the quantum speed limit  $v_{\text{QSL}}^W$  is fully determined by the metric properties of the dynamics (22), it is not *a priori* obvious how  $\tau_{\text{QSL}}^W$  behaves in the semiclassical, high-temperature limit  $k_B T \gg \hbar\gamma$ , where  $T$  is the temperature and  $\gamma$  the damping coefficient.

To address this question we turn again to the harmonic oscillator. Since we are now interested merely the behavior of  $\tau_{\text{QSL}}^W$  in the limit of high-temperatures, we now consider an un-driven system with potential,  $V(x) = 1/2 M\omega_0^2 x^2$ . In this case the exact master equation in Wigner space can be written as [54, 55]



$$\partial_t W(x, P, t) = \left[ -\frac{P}{M} \partial_x + V'(x) \partial_P + \partial_P (\gamma P + D_{PP} \partial_P) + D_{xP} \partial_{xP}^2 \right] W(x, P, t), \quad (27)$$

where  $D_{PP} = M\gamma/\beta + M\beta\gamma\hbar^2(\omega_0^2 - \gamma^2)/12$  and  $D_{xP} = \beta\gamma\hbar^2/12$ . Equation (27) is a quantum analog of the Klein–Kramers equation, which describes the dynamics of a harmonic oscillator in contact with a bosonic bath [54]. In figure 2 we plot the resulting quantum speed limit,  $v_{\text{QSL}}^{\text{W}}$  (22), together quantum speed limit time,  $\tau_{\text{QSL}}^{\text{W}}$  (26), again for  $p = 1$ . As initial state we chose a narrow Gaussian

$$W_0(x, P) = \frac{1}{2\pi \sigma_x \sigma_P} \exp\left(-\frac{(x - \mu_x)^2}{2\sigma_x^2}\right) \exp\left(-\frac{(P - \mu_P)^2}{2\sigma_P^2}\right). \quad (28)$$

As expected, the quantum speed limit time  $\tau_{\text{QSL}}^{\text{W}}$  vanishes in the classical, high-temperature limit. This observation confirms that the quantum speed limit is a purely quantum phenomenon also in open systems.

## 7. Concluding remarks

What is the ultimate limit on how fast a quantum system can evolve? At the verge of the age of quantum computing this century old question is more topical than ever. In the present work we highlighted that the maximal quantum speed can be fully characterized by the Schatten- $p$ -norms of the generator of quantum dynamics. We further showed that equivalent expressions can be found in Wigner phase space, where the computationally expensive operator norm is replaced by the absolute value of an imaginary valued function. The utility of the novel approach to quantum speed was illustrated by comparing the quantum speed limits in density operator space and Wigner phase space for the parametric harmonic oscillator. As a consistency check we finally verified that the bound is a pure quantum phenomenon, and that the dynamics of open classical systems is not restricted by the quantum bound. Therefore, we imagine that our results could prove useful for practical consequences and applications of the quantum speed limit, since for any situation the new bound is significantly easier to compute than the operator norm of high-dimensional density matrices.

## Acknowledgments

The author would like to thank Steve Campbell for insightful discussions. This work was supported by the US National Science Foundation under Grant No. CHE-1648973.

## Appendix A. Quantum speed from Schatten distance

This appendix is dedicated to a proof of equation (14). Consider the Schatten- $p$ -distance

$$\ell_p(\rho_t, \rho_0) = \|\rho_t - \rho_0\|_p \equiv (\text{tr}\{|\rho_t - \rho_0|^p\})^{1/p}, \quad (\text{A.1})$$

where  $p$  is an arbitrary, positive, real number,  $p \in [0, \infty)$ . Then the geometric speed can be written as

$$\dot{\ell}_p(\rho_t, \rho_0) = (\text{tr}\{|\rho_t - \rho_0|^p\})^{\frac{1}{p}-1} \text{tr}\{[(\rho_t - \rho_0)^2]^{\frac{p-1}{2}}(\rho_t - \rho_0)\dot{\rho}_t\}. \quad (\text{A.2})$$

The latter expression looks rather involved, but it can be simplified by using,  $\dot{\ell}_p(\rho_t, \rho_0) \leq |\dot{\ell}_p(\rho_t, \rho_0)|$  and employing the triangle inequality for operators,  $|\text{tr}\{O\}| \leq \text{tr}\{|O|\}$ , to read

$$\dot{\ell}_p(\rho_t, \rho_0) \leq (\text{tr}\{|\rho_t - \rho_0|^p\})^{\frac{1}{p}-1} \text{tr}\{|\rho_t - \rho_0|^{p-1}|\dot{\rho}_t|\}. \quad (\text{A.3})$$

Equation (A.3) can be further simplified with the help of Hölder's inequality [56]

$$\text{tr}\{|OB|\} \leq (\text{tr}\{|O|^{q_1}\})^{1/q_1} (\text{tr}\{|B|^{q_2}\})^{1/q_2} \quad (\text{A.4})$$

which is true for all  $1/q_1 + 1/q_2 = 1$ . Now choosing  $B = \dot{\rho}_t$  and  $q_1 = p/(p-1)$ , for which  $q_2 = p$ , we finally obtain the desired result (14)

$$\dot{\ell}_p(\rho_t, \rho_0) \leq \|\dot{\rho}_t\|_p. \quad (\text{A.5})$$

## Appendix B. Quantum speed from Wasserstein distance

Finally, we proof the expression for the quantum speed limit in Wigner phase space (22). To this end, we start with the Wasserstein- $p$ -distance [50], which is given by

$$\mathcal{D}_p(W_t, W_0) = \|W_t - W_0\|_p \equiv \left( \int d\Gamma |W(\Gamma, t) - W_0(\Gamma)|^p \right)^{1/p}. \quad (\text{B.1})$$

Accordingly we have

$$\begin{aligned} \dot{\mathcal{D}}_p(W_t, W_0) &= \left( \int d\Gamma |W(\Gamma, t) - W_0(\Gamma)|^p \right)^{\frac{1}{p}-1} \\ &\quad \times \int d\Gamma |W(\Gamma, t) - W_0(\Gamma)|^{p-1} \frac{W(\Gamma, t) - W_0(\Gamma)}{|W(\Gamma, t) - W_0(\Gamma)|} \dot{W}(\Gamma, t), \end{aligned} \quad (\text{B.2})$$

which can be simplified again with the help of the triangle inequality to read

$$\begin{aligned} \dot{\mathcal{D}}_p(W_t, W_0) &\leq |\dot{\mathcal{D}}_p(W_t, W_0)| \\ &\leq \left( \int d\Gamma |W(\Gamma, t) - W_0(\Gamma)|^p \right)^{\frac{1}{p}-1} \int d\Gamma |W(\Gamma, t) - W_0(\Gamma)|^{p-1} |\dot{W}(\Gamma, t)|. \end{aligned} \quad (\text{B.3})$$

In complete analogy to the derivation in density operator space, we now consider again Hölder's inequality [56]

$$\int dx |f(x)g(x)| \leq \left( \int dx |f(x)|^{q_1} \right)^{1/q_1} \left( \int dx |g(x)|^{q_2} \right)^{1/q_2}, \quad (\text{B.4})$$

which holds for all  $1/q_1 + 1/q_2 = 1$ . Once again choosing  $q_1 = p/(p-1)$  and  $g = \dot{W}_t$  we finally obtain

$$\dot{D}_p(W_t, W_0) \leq \|\dot{W}_t\|_p. \quad (\text{B.5})$$

## ORCID iDs

Sebastian Deffner  <https://orcid.org/0000-0003-0504-6932>

## References

- [1] Boixo S, Isakov S V, Smelyanskiy V N, Babbush R, Ding N, Jiang Z, Martinis J M and Neven H 2016 Characterizing quantum supremacy in near-term devices (arXiv:1608.00263)
- [2] Preskill J 2012 Quantum computing and the entanglement frontier (arXiv:1203.5813)
- [3] Ronnow T F, Wang Z, Job J, Boixo S, Isakov S V, Wecker D, Martinis J M, Lidar D A and Troyer M 2014 Defining and detecting quantum speedup *Science* **345** 420
- [4] Heisenberg W 1927 Über den anschaulichen Inhalt der quantentheoretischen kinematik und mechanik *Z. Phys.* **43** 172
- [5] Heisenberg W 1944 *Die Physikalischen Prinzipien der Quantumtheorie* 5th edn (Stuttgart: Hirzel)
- [6] Hilgevoord J 1998 The uncertainty principle for energy and time: II *Am. J. Phys.* **66** 396
- [7] Mandelstam L and Tamm I 1945 The uncertainty relation between energy and time in nonrelativistic quantum mechanics *J. Phys.* **9** 249
- [8] Uffink J 1993 The rate of evolution of a quantum state *Am. J. Phys.* **61** 935
- [9] Margolus N and Levitin L B 1998 The maximum speed of dynamical evolution *Physica D* **120** 188
- [10] Levitin L B and Toffoli Y 2009 Fundamental limit on the rate of quantum dynamics: the unified bound is tight *Phys. Rev. Lett.* **103** 160502
- [11] Bekenstein J D 1981 Energy cost of information transfer *Phys. Rev. Lett.* **46** 623–6
- [12] Lloyd S 2000 Ultimate physical limits to computation *Nature* **406** 1047
- [13] Deffner S and Lutz E 2010 Generalized clausius inequality for nonequilibrium quantum processes *Phys. Rev. Lett.* **105** 170402
- [14] Giovannetti V, Lloyd S and Maccone L 2011 Advances in quantum metrology *Nat. Photon.* **5** 222
- [15] Santos A C and Sarandy M S 2015 Superadiabatic controlled evolutions and universal quantum computation *Sci. Rep.* **5** 15775
- [16] Campbell S and Deffner S 2017 Trade-off between speed and cost in shortcuts to adiabaticity *Phys. Rev. Lett.* **118** 100601
- [17] Bhattacharyya K 1983 Quantum decay and the Mandelstam-Tamm-energy inequality *J. Phys. A: Math. Gen.* **16** 2993
- [18] Pfeifer P 1993 How fast can a quantum state change with time? *Phys. Rev. Lett.* **70** 3365
- [19] Giovannetti V, Lloyd S and Maccone L 2004 The speed limit of quantum unitary evolution *J. Opt. B* **6** 807
- [20] Frey M R 2016 Quantum speed limits: primer, perspectives, and potential future directions *Quantum Inf. Process.* **15** 3919
- [21] Barnes E 2013 Analytically solvable two-level quantum systems and Landau-Zener interferometry *Phys. Rev. A* **88** 013818
- [22] Poggi P M, Lombardo F C and Wisniacki D A 2013 Quantum speed limit and optimal evolution time in a two-level system *Europhys. Lett.* **104** 40005
- [23] Hegerfeldt G C 2013 Driving at the quantum speed limit: optimal control of a two-level system *Phys. Rev. Lett.* **111** 260501
- [24] Andersson O and Heydari H 2014 Quantum speed limits and optimal Hamiltonians for driven systems in mixed states *J. Phys. A: Math. Theor.* **47** 215301
- [25] Deffner S and Lutz E 2013 Energy-time uncertainty relation for driven quantum systems *J. Phys. A: Math. Theor.* **46** G5302
- [26] Richerme P, Gong Z-X, Lee A, Senko C, Smith J, Foss-Feig M, Michalakis S, Gorshkov A V and Monroe C 2014 Non-local propagation of correlations in long-range interacting quantum systems *Nature* **511** 198
- [27] Deffner S and Lutz E 2013 Quantum speed limit for non-Markovian dynamics *Phys. Rev. Lett.* **111** 010402
- [28] del Campo A, Egusquiza I L, Plenio M B and Huelga S F 2013 Quantum speed limits in open system dynamics *Phys. Rev. Lett.* **110** 050403
- [29] Taddei M M, Escher B M, Davidovich L and de Matos Filho R L 2013 Quantum speed limit for physical processes *Phys. Rev. Lett.* **110** 050402
- [30] Deffner S 2014 Optimal control of a qubit in an optical cavity *J. Phys. B: At. Mol. Opt. Phys.* **47** 145502
- [31] Zhang Y-J, Han W, Xia Y-J, Cao J-P and Fan H 2014 Quantum speed limit for arbitrary initial states *Sci. Rep.* **4** 4890
- [32] Mukherjee V, Carlini A, Mari A, Caneva T, Montangero S, Calarco T, Fazio R and Giovannetti V 2013 Speeding up and slowing down the relaxation of a qubit by optimal control *Phys. Rev. A* **88** 062326
- [33] Xu Z-Y, Luo S, Yang W L, Liu C and Zhu S 2014 Quantum speedup in a memory environment *Phys. Rev. A* **89** 012307
- [34] Xu Z-Y and Zhu S-Q 2014 Quantum speed limit of a photon under non-Markovian dynamics *Chin. Phys. Lett.* **31** 020301
- [35] Cimmarusti A D, Yan Z, Patterson B D, Corcos L P, Orozco L A and Deffner S 2015 Environment-assisted speed-up of the field evolution in cavity quantum electrodynamics *Phys. Rev. Lett.* **114** 233602
- [36] Marvian I and Lidar D A 2015 Quantum speed limits for leakage and decoherence *Phys. Rev. Lett.* **115** 210402
- [37] Hou L, Shao B, Wei Y-B and Zou J 2015 Quantum speed limit in a qubit-spin-bath system *J. Phys. A: Math. Theor.* **48** 495302
- [38] Mondal D, Datta C and Sazim S 2016 Quantum coherence sets the quantum speed limit for mixed states *Phys. Lett. A* **380** 689
- [39] Marvian I, Spekkens R W and Zanardi P 2016 Quantum speed limits, coherence, and asymmetry *Phys. Rev. A* **93** 052331
- [40] Mondal D and Pati A K 2016 Quantum speed limit for mixed states using an experimentally realizable metric *Phys. Lett. A* **380** 1395
- [41] Pires D P, Cianciaruso M, Céleri L C, Adesso G and Soares-Pinto D O 2016 Generalized geometric quantum speed limits *Phys. Rev. X* **6** 021031
- [42] Bengtsson I and Życzkowski K 2007 *Geometry of Quantum States: An Introduction to Quantum Entanglement* (Cambridge: Cambridge University Press)
- [43] Bures D 1969 An extension of Kakutani's theorem on infinite product measures to the tensor product of semifinite  $w^*$ -algebras *Trans. Amer. Math. Soc.* **135** 199

- [44] Jozsa R 1994 Fidelity for mixed quantum states *J. Mod. Opt.* **41** 2315
- [45] Deffner S and Campbell S 2017 Quantum speed limits: from Heisenberg's uncertainty principle to optimal quantum control *J. Phys. A: Math Theor.* in press
- [46] Simon B 1979 *Trace Ideals and Their Applications* vol 35 (Cambridge: Cambridge University Press)
- [47] Hu B L, Paz J P and Zhang Y 1992 Quantum Brownian motion in a general environment: exact master equation with nonlocal dissipation and colored noise *Phys. Rev. D* **45** 2843
- [48] Ford G W and O'Connell R F 2001 Exact solution of the Hu-Paz-Zhang master equation *Phys. Rev. D* **64** 105020
- [49] Dittrich T, Viviescas C and Sandoval L 2006 Semiclassical propagator of the wigner function *Phys. Rev. Lett.* **96** 070403
- [50] Wasserstein L N 1969 Markov processes over denumerable products of spaces, describing large systems of automata *Problems Inf. Transm.* **5** 47
- [51] Schleich W P 2011 *Quantum Optics in Phase Space* (New York: Wiley)
- [52] Husimi K 1953 Miscellanea in elementary quantum mechanics: II *Prog. Theo. Phys.* **9** 381
- [53] Deffner S and Lutz E 2013 Thermodynamic length for far-from-equilibrium quantum systems *Phys. Rev. E* **87** 022143
- [54] Dillenschneider R and Lutz E 2009 Quantum Smoluchowski equation for driven systems *Phys. Rev. E* **80** 042101
- [55] Deffner S 2013 Quantum entropy production in phase space *Europhys. Lett.* **103** 30001
- [56] Hölder O 1889 Ueber einen Mittelwerthssatz *Nachr. von Königlichen Ges. Wiss. und der Georg-Augusts-Universität zu Göttingen* **1889** 38