

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.



Opinion: Aerosol Remote Sensing Over The Next Twenty Years

Lorraine A. Remer¹, Robert C. Levy², J. Vanderlei Martins³

¹Goddard Earth Sciences Technology and Research 2 (GESTAR 2), University of Maryland Baltimore County, Baltimore MD, 21228, USA

5 ²NASA Goddard Space Flight Center, Greenbelt MD, 20771, USA

³Department of Physics, University of Maryland Baltimore County, Baltimore MD, 21250, USA

Correspondence to: Lorraine A. Remer (rem@umbc.edu)

Abstract. Twenty years ago, aerosol remote sensing underwent a revolution with the launch of the
10 Terra and Aqua satellites, followed by additional launches carrying new passive and active sensors. Capable of retrieving information about aerosol loading, rudimentary particle properties and in some cases aerosol layer height, the satellite view of Earth's aerosol system came into focus. Today we see trends developing in the aerosol remote sensing and modeling communities that allow us to speculate about the future and how the community will approach aerosol remote sensing twenty years from now.
15 We anticipate technology that will replace today's standard multi-wavelength radiometers with hyperspectral and/or polarimetry all viewing in multiple angles. These will be supported by advanced active sensors with the ability to measure profiles of aerosol extinction in addition to backscatter. The result will be greater insight into aerosol particle properties. Algorithms will move from being primarily physically-based to include an increasing degree of Machine Learning methods, but physically-based
20 techniques will not go extinct. However, the concept of applying algorithms to a single sensor will no longer exist. Retrieval algorithms will encompass multiple sensors and all available ground measurements into a unifying framework, and these inverted products will be ingested directly into assimilation systems, becoming "cyborgs": half observations, half model. In twenty years we will see a true democratization in space with nations large and small, private organizations and commercial
25 entities of all sizes launching space sensors. With this increasing amount of data and aerosol products available, there will be a lot of bad data. User communities will organize to set standards and the large national space agencies will lead the effort to maintain quality by deploying and maintaining validation ground networks and focused field experiments. Through it all, interest will remain high in the global aerosol system and how that systems affects climate, clouds, precipitation and dynamics, air quality,
30 transport of pathogens and fertilization of ecosystems, and how these processes are adapting to a changing climate.

1 NASA initiates a revolution with the Terra and Aqua missions

As NASA's Terra and Aqua satellites launched at the end of 1999 and in mid-2002, respectively, they ushered in a new era of aerosol remote sensing. These satellites carrying the MODerate resolution
35 Imaging Spectroradiometer (MODIS) (Salomonson et al., 1989) and the Multiangle Imaging



SpectroRadiometer (MISR, on Terra only)(Diner et al., 1998) offered an unprecedented view of the global aerosol system (Kaufman et al., 2002). The products derived from MODIS and MISR observations defined an aerosol climatology (Remer et al., 2008, Voss and Evan, 2020), identified regional trends (Yu et al., 2020, Zhang and Reid, 2010), advanced understanding of aerosol direct radiative forcing (Christopher and Zhang, 2002, Zhang and Christopher, 2003, Yu et al., 2006, Remer and Kaufman, 2006, Bellouin et al. 2008), discovered associations between aerosols, clouds and precipitation that shifted our perception of the water cycle (Kaufman et al., 2005, Koren et al., 2005, 2008, Yuan et al., 2011ab), proved the importance of aerosol intercontinental transport and nutrient deposition (Gao et al., 2001, Yu et al., 2013, 2019), and gave us important new tools to determine aerosol plume heights (Kahn et al., 2007, 2008, Val Martin et al., 2018) and air quality monitoring and mitigation (Engel-Cox, 2004, Van Donkelaar et al., 2006, 2016, Gupta and Christopher, 2009ab).

At the same time deriving these products from basic satellite observations pushed the field of aerosol remote sensing forward. The MODIS and MISR algorithms had to run in an operational environment. The algorithms had to work for all types of aerosols, in all types of environments and the products required validation. The MODIS and MISR algorithms are rooted in the physics of radiative transfer and the mathematics of inversions, but they also incorporate empirical assumptions that tie results to the real world (Kaufman et al., 1997, Tanré et al. 1997, Martonchik et al., 1998, 2002, Kahn et al., 1998). Assumptions are necessary because of the fundamental lack of information content in the MODIS and MISR observations (Tanré et al. 1996, Kahn et al., 1998). MODIS provides a multi-spectral look at each Earth scene at a single viewing geometry. The original MODIS aerosol algorithms, Dark Target (Remer et al., 2005) and Deep Blue (Hsu et al. 2006), make use of no more than six wavelengths for their retrieval, although they use additional wavelengths for cloud masking and other peripheral needs. The MODIS Dark Target over ocean retrieval makes use of the most spectral bands, six. These six bands contain only 2-3 pieces of independent aerosol information (Tanré et al., 1996). Over land, fewer bands are used and the information content available to characterize aerosol is even less (Levy et al., 2007). MISR employs fewer wavelengths, only four from 0.486 μm to 0.867 μm , but adds significant additional information by viewing each scene from 9 different angles. MISR's angular information has shown to be immensely valuable in constraining aerosol type and in determining aerosol plume height (Kahn et al., 2007, 2008, Val Martin et al., 2018, Kahn and Gaitley, 2015).

MODIS and MISR created a revolution in aerosol remote sensing when they were launched in 1999, but the reasons for the revolution involved more than new satellite technology married to state-of-the-science algorithms. A combination of factors came together at just the right time. These factors include:



- A policy for open data access,
- An investment in infrastructure for processing algorithms and distributing data,
- A commitment to long-term science teams for maintenance, improvement and validation of data products including on-orbit sensor calibration,
- The vision and continuous support of the AERONET program for product validation (Holben et al., 1998),
- The overall growth of the international aerosol community and expanded interest in the global aerosol system as part of Earth system science.

75 We find the first bullet point of particular interest for the celebration of the twentieth anniversary of the journal *Atmospheric Chemistry and Physics*, which itself propelled a revolution in open access and information sharing at roughly the same time.

85 Thus, the lessons learned from the previous 20 years in aerosol remote sensing is that technical innovation leads to more observable information that leads to more complete aerosol characterization at higher accuracy. But that chain of advancement will only seed new scientific endeavors if the aerosol products are accessible, validated and part of a larger data effort to characterize not only the global aerosol system but the Earth system as a whole. The infrastructure created for MODIS and MISR (LAADS DAAC, 2023, ASDC, 2023) continued to absorb and produce data products derived from
90 other sensors and missions over the past two decades: the Aura mission, Cloud-Aerosol Lidar with Orthogonal Polarization (CALIOP) data from Cloud-Aerosol Lidar and Infrared Pathfinder Satellite Observations (CALIPSO), the Polarization and Anisotropy of Reflectances for Atmospheric Sciences coupled with Observations from a Lidar (PARASOL) mission. Other agencies adopted NASA's open access policies for their missions and products: Visible Infrared Imaging Radiometer Suite (VIIRS),
95 Himawari, Sentinels, Geostationary Ocean Color Instrument (GOCI) etc. Today, a user needing local or global aerosol characterization may feel overwhelmed with the number of aerosol products available to use, both those derived directly from satellite observations and those inferred from models that use satellite information in assimilation systems (Zhang et al., 2008, Benedetti et al., 2009, Gelaro et al., 2015). The perspective from 2023 is that of an apparently data rich era that could hardly have been
100 imagined 23 years ago when Terra launched.

But what will aerosol remote sensing and data products look like over the next 20 years?

2 New technology leads to MORE, MORE, MORE information

105 The future of aerosol remote sensing will be propelled by advances in satellite payload technology. Instead of imaging multi-wavelength radiometers, like MODIS, we will expect the additional angular



information pioneered by MISR plus the advantages of polarization. Imaging multi-wavelength multi-angle polarimeters were introduced to the community with POLARization and Directionality of the Earth's Reflectances (POLDER)/PARASOL (Deuzé et al., 2000, 2001) with the benefits of the added
110 information content only apparent recently (Chen et al., 2020). Future imaging polarimeters will improve upon the POLDER/PARASOL technology with better polarization accuracy, hyper spectral and hyper angle capabilities, decreases in pixel size, increase in the number of wavelengths with polarization and still allow for imaging and enough view angles to optimize aerosol characterization. The Multi-viewing, Multi-channel, Multi-polarisation imaging mission (3MI) is the European follow-on
115 to POLDER/PARASOL that will fly on EUMETSAT's 2nd generation polar-orbiting Meteorological Observing satellite series (MetOp SG) (Fougnie et al. 2018). The first of this series will be launched in the 2023-2024 time frame. Other instruments such as the HyperAngle Rainbow Polarimeter 2 (HARP2) and the Spectro-polarimeter for Planetary Exploration One (SPEXone) will fly on NASA's Plankton, Aerosol, Clouds, ocean Ecosystem (PACE) mission scheduled to launch in 2024 (Werdell et al. 2019),
120 and the Multi Angle Imager for Aerosols (MAIA) will be launched in the early 2020s (Diner et al., 2018). Similar instruments with extended wavelength ranges are part of the planning for NASA's Atmospheric Observing System (AOS) with launch of its first observatory before the end of the current decade with observations expected throughout the 2030s. Beyond AOS we expect more hyperspectral capability, broader wavelength ranges including the thermal infrared, finer spatial resolution, broader
125 swaths and more angles. Twenty years from now these imaging multi-angle and hyperspectral polarimeters will be the norm for passive aerosol characterizing sensors and simple single-view multi-wavelength radiometers will be antiques.

In the thermal infrared even single-view instruments will still have a role (DeSouza-Machado, 2010,
130 Klüser et al. 2011). Hyperspectral thermal infrared spectrometers such as the Atmospheric Infrared Sounder (AIRS) have been used to detect and quantify dust AOD and layer height beginning with the Aqua mission (DeSouza-Machado et al., 2006) and advancing to other hyperspectral sounders in this spectral range such as the Infrared Atmospheric Sounding Interferometer (IASI) (Clarisse et al., 2019). The community has recently directed greater focus to dust's contribution to Earth energy balance
135 through its perturbation of the radiation field at infrared wavelengths (Zheng et al., 2022, Song et al., 2022). New technology for hyperspectral thermal infrared spectrometers will improve signal, reduce spatial footprints and are our best hope to achieve mineral speciation of dust from space.

Along with advances in passive remote sensing imagers will be new capabilities in active lidars. The
140 straightforward backscatter lidar in space (CALIOP) has demonstrated the unique aerosol information that only a lidar can provide. By 2043 standard aerosol missions will include lidars and these lidars will surpass CALIOP with additional wavelengths, High Spectral Resolution Lidar (HSRL) or Raman capability and multiple beams to expand coverage. Ground-based and airborne lidars with these enhanced capabilities are demonstrating that like polarimetry, information content from enhanced lidar
145 design allows for aerosol characterization in ways previously unattainable (Müller et al., 2014). Lidars on the books for the future include: a 2-wavelength lidar with one HSRL channel for NASA's AOS mission (AOS, 2023), and the ATmospheric backscatter LIDar (ATLID) a one-wavelength in the ultraviolet lidar with HSRL capability for ESA's EARTHCARE mission (Groß et al., 2015). By 2043,



150 these modest steps into lidar capability expansion will grow to achieve the $3\beta + 2\alpha + 1\delta$ ideal for aerosol
characterization (Müller et al., 2014). This means that there will be at least one space payload offering 3
wavelengths of backscattering (β), two wavelengths of extinction profiles (α) using HSRL technology
and one wavelength providing depolarization (δ). Aerosol scientists have been advocating for this array
of lidar payload capability for more than a decade, and despite the next set of planned aerosol missions
falling short, we will see the dream become reality in 20 years.

155 The lessons we learned from the past twenty years in terms of the advantages of formation flying will
guide mission planning far into the future. Like the A-Train of the 2000s and 2010s, missions will be
constructed to purposely match passive polarimeters with active lidars in ways that promote synthesized
data processing, algorithms, science and applications. The workhorse sun-synchronous polar orbiting
160 mission that has been the mainstay for aerosol characterization will be supplemented with inclined
orbits that have been more typical of precipitation-centered missions but have much to offer for
aerosols. Inclined orbits allow for a multitude of geometries and time-of-day sampling of the aerosol
system over the course of weeks as well as multiple coincidences a day with other satellites looking
simultaneously at the same target at the ground. This provides a peek into aerosol situations never
165 encountered by MODIS, MISR or their ilk. Part of this expanding constellation of aerosol observations
from satellite will be the geosynchronous imagers (GEO). Already capability from GEO platforms has
evolved from the two or three broad uncalibrated wavelengths that were ubiquitous at the time of Terra
launch into multi-wavelength moderate resolution sensors that mimic most of MODIS's capabilities. By
2043, these modern-day Advanced Baseline Imagers (ABI) and Advanced Himawari Imager (AHI) will
170 be upgraded with spectrometers, as per the NASA/NOAA joint Geostationary Extended Observations
(GEO-XO) program (Frost et al., 2020). At that time we expect studies in progress to adapt polarimeters
and a scanning backscatter lidar to GEO orbit. With the equator crowded with observatories in GEO
orbit, overlap between sensors provides both a means to cross-calibrate and also multi-angle looks at
each scene to increase information content for aerosol retrieval (Bian et al., 2021).

175 The tag line for technology in 2043 will be MORE, MORE, MORE! Each sensor will be collecting
orders of magnitude more data than does MODIS today. Polarized measurements require three times the
data per wavelength than a simple radiometer to account for the three angles of polarization. Multiply
that increase by the number of view angles beyond the single MODIS-like view and we are likely now
180 producing 15X to 60X the amount of information being produced today. The implementation of
hyperspectral sensors or improvements in pixel size while maintaining broad swaths to view the entire
globe further increases data volume. Advancing lidars with additional wavelengths, HSRL, multiple
beams and quicker pulses also magnifies data volume quickly. As part of technology evolution will be
the need for spacecraft and especially spacecraft electronics, data systems and transmission to keep up
185 with data volumes. Laser transmission will replace radio frequencies and infrastructure in space will
serve as way stations to stage downlinking to overworked ground stations. On Earth, computing power
will be fully cloud-based and ever expanding to hold the proliferation of collected data.



3 Evolution of Algorithms

190 Today the majority of aerosol retrieval algorithms interpret the signal measured by the satellite sensor in terms of radiative transfer through the atmosphere, governed by this equation,

$$\rho_{TOA} = \rho_a + \frac{T\rho_s}{(1 - s\rho_s)}$$

195 Where ρ_{TOA} is the solar radiance reflected by the Earth system measured at the top of the atmosphere by the satellite sensor, ρ_a is the contribution by the atmosphere alone, ρ_s is the reflectance of Earth's surface that transmits through the atmosphere with transmittance, T , and s is the spherical albedo with the $(1 - s\rho_s)$ term representing multiple scattering between the atmosphere and Earth's surface. Each term is for a specific wavelength, sun, and sensor geometries, but that notation has been omitted for
200 simplicity. The terms depend on surface properties (ρ_s and s) and the amount and optical properties of the gases and aerosols in the atmosphere (ρ_a and T). By making physically-based assumptions on some surface, gas and aerosol properties and calculating ρ_{TOA} , the calculated ρ_{TOA} can be compared with the measured ρ_{TOA} and assumptions adjusted until the calculated and measured ρ_{TOA} are the same to within an acceptable error.

205 The simplest physically-based algorithms use one wavelength, one geometry at a time. These simple retrievals hold all assumptions constant except for the optical measure of aerosol loading, the aerosol optical depth (AOD). Thus, the retrievals use one piece of information to retrieve one parameter, the AOD. By minimizing the difference between measured and calculated ρ_{TOA} simultaneously for more
210 than one wavelength, more information is introduced and additional assumptions can become free parameters leading to retrievals of aerosol parameters beyond AOD. Likewise, additional geometries and/or polarization states can be introduced, freeing even more parameters for retrieval. The simplest algorithms have employed radiative transfer models run in the forward direction with assumed aerosol, gases, and surface parameters resulting in a Look Up Table (LUT) stored for future use. The retrieval
215 then can move quickly through the LUT, interpolating between entries, and return the retrieved parameters in an operational setting.

Today, as information content increases and computer power grows to meet demand, LUTs are being replaced by other methods to solve the radiative transfer equation. Cost-function minimization methods
220 and other multivariate tools are being used to simultaneously retrieve multiple aerosol, gases, and surface parameters. These more flexible algorithms produce more parameters and require fewer



assumptions than the old LUT algorithms, but they are still physically-based in radiative transfer. As we march towards 2043, the jettison of the LUT methods will become complete, even as radiative transfer becomes more sophisticated with better representation of aerosol optical properties including irregularly shaped particles with complex composition. Algorithms will be designed to ingest data from multiple collocated sensors, especially combinations of passive and active sensors, and combinations of space-borne and ground-based sensor observations. We expect joint retrievals of aerosol and trace gases, aerosol and land properties, aerosol and ocean color properties, and aerosol/gas/surface retrievals. Eventually there will be sufficient information by 2043 for retrieved properties to include vertical profiles of aerosol extinction and particle number concentration, retrievals over clouds and over all land surface types including snow and ice. Characterization will include particle spectral absorption and make progress towards constraining particle composition.

Development of Machine Learning (ML) retrieval algorithms will accelerate, reach a plateau of usefulness and then become incorporated within physically-based algorithms. In 2043 there will be algorithms that produce a full array of aerosol characteristics by pumping satellite-measured quantities through ML-derived models. These models have no physical basis and do not rely on the radiative transfer equation (Eq. 1). Instead, the models are created by training on a formulation data set that matches satellite-measured radiances with measured aerosol characteristics. For example, today we can train a ML model using satellite measurements collocated with AERONET observations (Lary et al., 2009, Di Noia and Hasekamp, 2018, Kang et al., 2022). In this way we can obtain aerosol characteristics such as AOD that match AERONET with great accuracy and precision (Lary et al., 2009), but we lose all physical understanding of how that AOD is retrieved. The challenge for exclusive ML methods like these cited will be the identification of accurate measurements of aerosol characteristics, other than AOD, in sufficient quantity to train the ML models.

There are other uses of ML in aerosol remote sensing besides directly obtaining aerosol parameters from satellite observations. Within a physically-based algorithm there are many assumptions that can be aided by ML methods. For example, running a radiative transfer model is computationally expensive. A ML model derived from radiative transfer runs will be more accurate than current interpolations of LUTs and save time in an operational setting (Gao et al., 2021, Ukkonen 2022). In other work, ML methods are used to classify specific aerosol types, such as dust by training on other satellite data sets (CALIPSO) (Lee et al., 2021). ML is used to bias-correct aerosol products derived using standard methods (Lipponen et al. 2022), and it can help with interpolation, constraining non-retrievable aerosol optical properties, cloud or snow masking, and surface reflectance determination. In 2043 almost all



aerosol retrieval algorithms will continue to be physically-based, but when examined carefully will contain many elements of indispensable ML within.


4 The rise of the cyborg: Blends of measurements and models

260 The biggest noticeable change over the next twenty years will be the gradual extinction of “satellite
aerosol products”. The entire purpose of what we know now as a satellite data product will be to feed
major assimilation systems that produce a complete representation of the global aerosol system within
the context of the complex Earth system (Zhang et al., 2008, Benedetti et al., 2009, Gelaro et al., 2015).
We already see the widespread acceptance of assimilation products, as noted by the number of citations
265 earned by key papers describing each system. For example, using Clarivate’s Web of Science data base
(Clarivate, referenced January 2023)) and searching on “satellite AND aerosol” we find the first
returned reference to be Holben et al. (1998), the seminal AERONET paper with 5265 citations
accumulated over 24 years. Next up is Gelaro et al. (2015), the fundamental reference for the Modern-
Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2) assimilation
270 products, with 3270 citations accumulated over only nine years. We understand that not all citations to
MERRA2 will involve applications to MERRA2 *aerosol* products and that having a single fundamental
reference concentrates users’ citations to a single paper, but the rate of acceptance of MERRA2 and
other systems’ assimilation products over the past five years is astounding. See Figure 1.

275 Assimilation systems are analogous to the science fiction concept of the “cyborg”, defined by the
Oxford Dictionary of Science Fiction as “A hybrid being: half human, half machine (a contraction of
‘cybernetic organism’)” (Prucher, 2006). In our context, the assimilation systems are a hybrid of half
observations, half models. Satellites measure electromagnetic radiation that is processed through aerosol
retrieval algorithms to determine aerosol parameters. Assimilation systems intercept parameters from
280 the retrieval algorithms during processing to ingest that data into a global model that adjusts model
representation of aerosol parameters to agree with the observations. Some assimilation systems draw
from the aerosol retrieval algorithms close to the satellite observations, making use of radiance or
reflectance directly (Randles et al., 2017). Some systems ingest the final retrieval products such as AOD
(Zhang et al., 2008). In all cases the global model is drawn closer to actual measurements at regular
285 time steps and is expected to represent the actual aerosol system better than a model that is initialized
and run forward from aerosol source inputs. The advantage of assimilation aerosol products is that users
can access aerosol parameters for any time period, at any location, at any altitude. The parameters will
have the benefit of ALL available data from ALL satellites simultaneously. User’s will not need to



patch together products from different algorithms. The model does that for you and presents the results
in an Earth system governed by physical restraints. The assimilation systems will run in the cloud and
will be called on to produce the cyborg data at user's will.



295 **5 Satellites, satellites everywhere**

A gold rush to space has begun and will intensify over the next 20 years as space becomes crowded with a variety of sensors in various Earth orbits. During this period we will realize a true democracy in



space as non-traditional players such as the space agencies from the developing world and from a wide range of commercial sector companies join traditional national agencies in blanketing the skies with satellites and sensors. While only a fraction of these satellite sensors will be Earth-viewing and aerosol-capable, the result will yield a tremendous volume of data useful for aerosol retrieval.

Satellites will come in all sizes. Traditional agencies will still launch large, expensive, robust flagship missions with multiple platforms and multiple sensors on each platform. The flagship missions will continue demonstrating lower risk, higher accuracy, greater reliability and longevity. The flagship missions will be supplemented by and integrated with constellations of smaller satellites and nano-sized cubesats. The smaller satellites will be orders of magnitude less expensive, and thus riskier and with more limited capability than the larger observatories. However, the multitude of the smaller units renders the loss of any particular unit as insignificant to the greater objectives. With so many satellites flying, observing Earth and returning measurements for aerosol retrieval, each satellite will be seeking a particular niche. All orbit types will be covered including polar-orbiting, inclined orbits and geosynchronous, and near real time observations will be achieved.

The challenge of so many data sources is the maintenance of data quality. There will be a lot of data, and there will be a lot of bad data. Eventually the scientific and user communities will organize to set standards and protocols to regulate data quality. Such standards will rely on the cohesion of the data communities for enforcement. Not all data will be free and open access. Commercialization of data products will aid in enforcement of standards, as different entities will be competing to sell their products based on many characteristics including quality. Commercial entities that launch satellites intend to produce specialized data products for specific customers. Sometimes these customers will be national agencies looking for input to their assimilation systems. Other times commercial entities will be producing their own assimilation system cyborg products and selling those results to their customers.

6 Validation systems will need to keep up

With the democratization of space, the leading national space agencies will no longer hold a monopoly on space access. Ironically, instead these government agencies that pioneered space will take the lead on *suborbital* measurements to support the validation side of producing high quality aerosol products. Government-sponsored field experiments and ground networks will provide the basis for the entire international community, including the commercial sector, to evaluate their products and improve their algorithms



Today's basis for validation of aerosol data sets is the ground-based AERONET network (Holben et al., 1998, Giles et al. 2019). The AERONET concept will continue for decades into the future, but the technology itself will evolve. Instruments will move from finite wavelength bands to hyperspectral spectrometers and will continue AERONET's current trend into the shortwave infrared. Originally, AERONET instruments had invested in polarimetry, which turned out to be a capability ahead of its time. The community has caught up and AERONET will expand the polarimetric capabilities of their networks and incorporate polarimetry into their retrievals. Data capacity, storage and transmission will expand, as will the network itself. AERONET's subsidiary, the Marine Aerosol Network (MAN) will continue to collect data on ocean-going cruises and ships of opportunity. The MAN instruments will also upgrade their technology and data capabilities without losing accuracy. The AERONET retrieval that produces reliable aerosol characterization in addition to AOD, including particle size distribution, single scattering albedo, complex refractive indices, and non-sphericity, will become a more important asset over time as aerosol satellite remote sensing advances in information content, requiring validation of a wide range of aerosol parameters.

The AERONET concept of a widespread, high quality, aerosol observing ground-network will proliferate, as we are already witnessing with the Interagency Monitoring of Protected Visual Environments (IMPROVE: Malm and Hand, 2007), the European Aerosol Research Lidar Network (EARLINET: Pappalardo et al., 2014), the Aerosol, Cloud and Trace gases Research Initiative (ACTRIS), the Surface Particulate Matter Network (SPARTAN: Snider et al., 2015) and other networks. The different networks offer different types of aerosol characterization necessary to keep up with the expansion of retrieved aerosol parameters from space. In-situ measurements of aerosol aerodynamic properties will continue to be essential for the assessment of the aerosol health impact and to link aerodynamic to optical measurements, which form the basis of satellite products. Ground LIDAR networks (Pappalardo et al., 2014) will also expand to high temporal characterization of the aerosol vertical profile. As satellite retrieval algorithms upgrade their radiative transfer, refine their assumptions, incorporate ML and become cyborgs, so will the ground network retrieval algorithms. More and more often validation data sets will include the resulting aerosol characterization obtained by simultaneous retrieval of several collocated instruments. Ideally this will include passive measurements with collocated lidar and ground-level or balloon-borne in situ measurements.

Field experiments featuring aircraft-borne measurements will be indispensable for validating the satellite observations at the radiance/polarization level (Level 1) and the retrieved aerosol particle properties in regions and situations of high scientific interest or societal relevance. Aircraft fly in three dimensions, sampling situations between network stations and providing vertical profiles of parameters



where lidar is absent. Field experiments will also be essential for confirming ground-based validation sites, as now these sites will be producing a suite of retrieved parameters that require their own validation. We will find that field experiment measurements will be the only truly independent data set for validation. With the abundance of suborbital data, quality again becomes an issue, and national
370 agencies will be the only community players with the resources to maintain high quality, long-term validation data sets.

At the opposite end of the spectrum will be the proliferation of low-cost citizen science networks. These commercial networks will be everywhere and will contribute to characterizing the global aerosol system
375 simply by the abundance of available observations. However, there will not be the same oversight to maintain quality of these data sets and users will need to calibrate the data from citizen scientist measurements with government-maintained and quality-assured sources.

7 Unanswered science questions will drive the new systems

We have described a vibrant global measurement system focused on aerosol remote sensing as we move
380 forward into the next twenty years. This picture assumes a continuing need for aerosol characterization, which will be driven by pressing science questions and societal priorities. What will be the aerosol science questions of the coming decades?

Quantifying changes to the aerosol system will be essential as we continue to struggle to catch up to
385 climate change. By 2043 we will have 70 years of Total Ozone Measuring System (TOMS), 60 years of AVHRR and 40 years of MODIS + VIIRS aerosol time series. While most of these time series offer insight into global and regional changes to aerosol loading, inferring changes to aerosol types and particle properties will be especially important.

390 We will still be working towards understanding aerosol-cloud-precipitation processes, how these have changed in time and how do these processes interact with atmospheric dynamics.

The role of aerosols in transporting nutrients around the globe and depositing them into terrestrial and oceanic ecosystems will continue to be a pressing issue in the coming decades. Likewise, the role of
395 aerosols in transporting pathogens will still be under investigations in the coming years.

Maintaining healthy air quality for all the world's populations will continue to call for understanding of global aerosol distributions and characterization in a way that will require satellite remote sensing.



400 Underlying these issues will be the goal of using the characterization of the global aerosol system to make our planet healthier and environmentally just.

8 So what will 2043 look like?

Despite today's assessment that we are "data rich", by 2043 we will look back at 2023 as a primitive data desert. Satellite-derived aerosol products that are actually cyborgs produced by assimilation
405 systems will represent the global aerosol, producing near real time representation of aerosol fields at fine temporal and spatial scales. Aerosol product users will forget that clouds (real clouds not data clouds) exist, as all aerosol fields will be interpolated through traditionally cloudy scenes by the assimilation models. Likewise, there will be no gaps due to difficult surfaces such as sun glint, snow or ice. However, the challenge will be the quality checking and synthesis of so much data. There will be
410 commercial aerosol data ambassadors who, for a fee, will create custom data packages from the overwhelming supply of public and private data product sources.

Today's retrieval algorithms will appear quaint to aerosol remote sensing scientists in 2043. The days of a single inversion applied to a single sensor will be long gone. Even physically-based algorithms will
415 be designed to invert the observations from multiple sensors and platforms simultaneously. This includes joint inversions between satellite and ground-based sensors to produce full characterization of atmospheric constituents (aerosols and gases), as well as complete characterization of the surface beneath whether that surface be a cloud, the ocean or a snow field.

420 Few people will care about aerosol optical depth. The assimilation systems will offer aerosol characterizations in forms most useful for individual users. Air quality managers will automatically obtain particulate mass concentration. Aerosol-cloud-precipitation scientists will obtain vertically-resolved number concentration or heating rates due to aerosol absorption. If anybody is still interested in aerosol radiative forcing, those numbers in Wm^{-2} will come automatically out of some assimilation
425 system. There will be attempts to produce deposition mass flux, but the error bars will still be large. Likewise, attempts to reliably retrieve chemical composition directly from remote sensing likely will fail, although aerosol type will be constrained very well. The only interest in aerosol optical depth will be users working towards continuity with the old sensors for long-term trend analysis.

430 In 2043 validation networks must be in place and major research efforts will switch from algorithm development to designing and implementing a strategy that reserves sufficient independent data for



validation. If the assimilation is producing a cyborg vertically-resolved distribution of particle number concentration near or in clouds, then there needs to be a validation strategy to confirm and put error bars on that product. Investment will be needed to develop the instruments and platforms needed to
435 implement a validation strategy. True validation of some difficult parameters such as vertically-resolved particle number validation may be measured only infrequently during dedicated field campaigns. In some ways validation in 2043 may look like a step backward from what aerosol scientists have come to expect today. Instead of 1 million collocations that we achieve now when we validate AOD with AERONET, for some retrieved aerosol parameters, we will be satisfied with 10 to 50 validation points
440 hard-earned in the field.

In summary, 2043 will be an exciting time for aerosol remote sensing. There will be plenty of data for answering science questions and plenty of work to do to assure the integrity of that data. We are sorry to say, though, that we cannot promise flying hover boards as seen in the *Back To The Future* movies
445 (Zemeckis and Gale, 1989, Schildhouse, 2014), even in 2043.

References

- 450 ACTRIS (Aerosol, Cloud and Trace gases Research Initiative) <https://www.actris.eu>, referenced 29 May 2023
- AOS (Atmospheric Observation System) <https://aos.gsfc.nasa.gov/mission.htm>, referenced 05 February 2023.
- 455 ASDC (Atmospheric Science Data Center) <https://asdc.larc.nasa.gov/about>, referenced 05 February 2023
- Bellouin, N., Jones, A., Haywood, J., and Christopher, S. A., Updated estimate of aerosol direct radiative forcing from satellite observations and comparison against the Hadley Centre climate model, J. Geophys. Res., 113, D10205, doi:10.1029/2007JD009385, 2008.
- 460 Benedetti, A., Morcrette, J.J., Boucher, O., Dethof, A., Engelen, R.J., Fisher, M., Flentje, H., Huneeus, N., Jones, L., Kaiser, J.W., and Kinne, S.: Aerosol analysis and forecast in the European centre for medium-range weather forecasts integrated forecast system: 2. Data assimilation. J. Geophys. Res. Atmos., 114, D13205, doi:10.1029/2008JD011115, 2009.



- 465 Bian, Q., Kreidenweis, S., Chiu, J.C., Miller, S.D., Xu, X., Wang, J., Kahn, R.A., Limbacher, J.A.,
Remer, L.A. and Levy, R.C.: Constraining Aerosol Phase Function Using Dual-View Geostationary
Satellites. *J. Geophys. Res. Atmos.*, 126, doi:10.1029/2021JD035209, 2021.
- Chen, C., Dubovik, O., Fuertes, D., Litvinov, P., Lapyonok, T., Lopatin, A., Ducos, F., Derimian, Y.,
Herman, M., Tanré, D., Remer, L. A., Lyapustin, A., Sayer, A. M., Levy, R. C., Hsu, N. C., Descloitres,
470 J., Li, L., Torres, B., Karol, Y., Herrera, M., Herreras, M., Aspetsberger, M., Wanzelboeck, M.,
Bindreiter, L., Marth, D., Hangler, A., and Federspiel, C.: Validation of GRASP algorithm product from
POLDER/PARASOL data and assessment of multi-angular polarimetry potential for aerosol
monitoring. *Earth Syst. Sci. Data*, 12, 3573–3620, doi.org/10.5194/essd-12-3573-2020, 2020.
- Clarivate. <https://clarivate.com/webofsciencegroup/solutions/web-of-science/>
- 475 Deuzé, J.L., Goloub, P., Herman, M., Marchand, A., Perry, G., Susana, S. and Tanré, D.: Estimate of
the aerosol properties over the ocean with POLDER. *J. Geophys. Res. Atmos.*, 105, 15329-15346,
doi.org/10.1029/2000JD900148, 2000.
- Deuzé, J.L., Bréon, F.M., Devaux, C., Goloub, P.H., Herman, M., Lafrance, B., Maignan, F., Marchand,
A., Nadal, F., Perry, G. and Tanré, D.: Remote sensing of aerosols over land surfaces from POLDER-
480 ADEOS-1 polarized measurements. *J. Geophys. Res. Atmos.*, 106, 4913-4926,
doi.org/10.1029/2000JD900364, 2001.
- Christopher, S.A. and Zhang, J.: Shortwave aerosol radiative forcing from MODIS and CERES
observations over the oceans. *Geophys. Res. Lett.*, 29, 6-1, doi.org/10.1029/2002GL014803, 2002.
- Clarisse, L., Clerbaux, C., Franco, B., Hadji-Lazaro, J., Whitburn, S., Kopp, A. K., et al.: A decadal
485 data set of global atmospheric dust retrieved from IASI satellite measurements. *J. Geophys. Res.*
Atmos., 124, 1618– 1647. doi.org/10.1029/2018JD029701, 2019.
- DeSouza-Machado, S. G., Strow, L. L., Hannon, S. E., and Motteler, H. E.: Infrared dust spectral
signatures from AIRS, *Geophys. Res. Lett.*, 33, L03801, doi:10.1029/2005GL024364, 2006.
- DeSouza-Machado, S.G., Strow, L.L., Imbiriba, B., McCann, K., Hoff, R.M., Hannon, S.E., Martins,
490 J.V., Tanré, D., Deuzé, J.L., Ducos, F. and Torres, O.: Infrared retrievals of dust using AIRS:
Comparisons of optical depths and heights derived for a North African dust storm to other collocated
EOS A-Train and surface observations. *J. Geophys. Res. Atmos.*, 115, doi.org/10.1029/2009JD012842,
2010.
- 495 Diner, D.J., Boland, S.W., Brauer, M., Bruegge, C., Burke, K.A., Chipman, R., Di Girolamo, L., Garay,
M.J., Hasheminassab, S., Hyer, E. and Jerrett, M.: Advances in multiangle satellite remote sensing of
speciated airborne particulate matter and association with adverse health effects: from MISR to
MAIA. *J. Appl. Rem. Sens.*, 12, 042603-042603, doi.org/10.1117/1.JRS.12.042603, 2018.
- 500 Diner, D.J., Beckert, J.C., Reilly, T.H., Bruegge, C.J., Conel, J.E., Kahn, R.A., Martonchik, J.V.,
Ackerman, T.P., Davies, R., Gerstl, S.A. and Gordon, H.R.: 1998. Multi-angle Imaging



- SpectroRadiometer (MISR) instrument description and experiment overview. *IEEE Tran. Geosci. Rem. Sens.*, 36, 1072-1087, 10.1109/36.700992, 1998.
- 505 Di Noia, A. and Hasekamp, O.P.: Neural networks and support vector machines and their application to aerosol and cloud remote sensing: A review. *Springer Series in Light Scattering: Volume 1: Multiple Light Scattering, Radiative Transfer and Remote Sensing*, 279-329, doi.org/10.1007/978-3-319-70796-9_4, 2018.
- 510 Engel-Cox, J.A., Holloman, C.H., Coutant, B.W., and Hoff, R.M.: Qualitative and quantitative evaluation of MODIS satellite sensor data for regional and urban scale air quality. *Atmos. Environ.*, 38, 2495-2509, doi.org/10.1016/j.atmosenv.2004.01.039, 2004.
- Fougnie, B., Marbach, T., Lacan, A., Lang, R., Schlüssel, P., Poli, G., Munro, R. and Couto, A.B.: The multi-viewing multi-channel multi-polarisation imager—Overview of the 3MI polarimetric mission for aerosol and cloud characterization. *J. Quant. Spec. Rad. Trans.*, 219, 23-32, doi.org/10.1016/j.jqsrt.2018.07.008, 2018.
- 515 Frost, G.J. and co-authors: A Value Assessment of an Atmospheric Composition Capability on the NOAA Next-Generation Geostationary and Extended Orbits (GEO-XO) Missions. NOAA technical report OAR CPO, 8. <https://doi.org/10.25923/1s4s-t405>, 2020.
- 520 Gao, M., Franz, B.A., Knobelspiesse, K., Zhai, P.W., Martins, V., Burton, S., Cairns, B., Ferrare, R., Gales, J., Hasekamp, O. and Hu, Y.: Efficient multi-angle polarimetric inversion of aerosols and ocean color powered by a deep neural network forward model. *Atmos. Meas. Tech.* 14, 4083-4110, 2021.
- Gao, Y., Kaufman, Y. J., Tanré, D., Kolber, D., Falkowski, P. G. Seasonal distributions of aeolian iron fluxes to the global ocean. *Geophys. Res. Lett.*, 28, 29–32. <https://doi.org/10.1029/2000GL011926>, 2001.
- 525 Gelaro, R. and Coauthors. The Modern-Era Retrospective Analysis for Research and Applications, Version 2 (MERRA-2). *J. Climate*, 30, 5419–5454, <https://doi-org.proxy-bc.researchport.umd.edu/10.1175/JCLI-D-16-0758.1>, 2017.
- 530 Giles, D.M., Sinyuk, A., Sorokin, M.G., Schafer, J.S., Smirnov, A., Slutsker, I., Eck, T.F., Holben, B.N., Lewis, J.R., Campbell, J.R. and Welton, E.J., Advancements in the Aerosol Robotic Network (AERONET) Version 3 database—automated near-real-time quality control algorithm with improved cloud screening for Sun photometer aerosol optical depth (AOD) measurements. *Atmos. Meas. Tech.*, 12(1), 169-209, <https://doi.org/10.5194/amt-12-169-2019>, 2019.
- 535 Groß, S., Freudenthaler, V., Wirth, M. and Weinzierl, B., Towards an aerosol classification scheme for future EarthCARE lidar observations and implications for research needs. *Atmospheric Science Letters*, 16(1), 77-82, doi.org/10.1002/asl2.524, 2015.



- 540 Gupta, P., Christopher, S. A. Particulate matter air quality assessment using integrated surface, satellite,
and meteorological products: Multiple regression approach, *J. Geophys. Res.*, 114, D14205,
doi:10.1029/2008JD011496, 2009.
- Gupta, P., Christopher, S. A. Particulate matter air quality assessment using integrated surface, satellite,
and meteorological products: 2. A neural network approach, *J. Geophys. Res.*, 114, D20205,
545 doi:10.1029/2008JD011497, 2009.
- Hsu, N.C., Tsay, S.C., King, M.D. and Herman, J.R., Deep blue retrievals of Asian aerosol properties
during ACE-Asia. *IEEE Transactions on Geoscience and Remote Sensing*, 44(11), 3180-3195,
10.1109/TGRS.2006.879540,
2006.
- 550 Hsu, N.C., Jeong, M.J., Bettenhausen, C., Sayer, A.M., Hansell, R., Seftor, C.S., Huang, J. and Tsay,
S.C., Enhanced Deep Blue aerosol retrieval algorithm: The second generation. *J. Geophys. Res.*
Atmos., 118(16), 9296-9315, doi.org/10.1002/jgrd.50712, 2013.
- Kahn, R., Banerjee, P., McDonald, D. and Diner, D.J., Sensitivity of multiangle imaging to aerosol
555 optical depth and to pure-particle size distribution and composition over ocean. *J. Geophys. Res.*
Atmos., 103(D24), 32195-32213, doi.org/10.1029/98JD01752, 1998.
- Kahn, R. A., Li, W.-H., Moroney, C., Diner, D. J., Martonchik, J. V., and Fishbein, E. Aerosol source
plume physical characteristics from space-based multiangle imaging, *J. Geophys. Res.*, 112, D11205,
560 doi:10.1029/2006JD007647, 2007.
- Kahn, R. A., Chen, Y., Nelson, D. L., Leung, F.-Y., Li, Q., Diner, D. J., and Logan, J. A., Wildfire
smoke injection heights: Two perspectives from space, *Geophys. Res. Lett.*, 35, L04809,
doi:10.1029/2007GL032165, 2008.
- Kahn, R.A. and Gaitley, B.J., An analysis of global aerosol type as retrieved by MISR. *J. Geophys. Res.*
565 *Atmos.*, 120(9), 4248-4281, doi.org/10.1002/2015JD023322, 2015.
- Kang, Y., Kim, M., Kang, E., Cho, D. and Im, J., Improved retrievals of aerosol optical depth and fine
mode fraction from GOCI geostationary satellite data using machine learning over East Asia. *ISPRS*
Journal of Photogrammetry and Remote Sensing, 183, 253-268,
570 <https://doi.org/10.1016/j.isprsjprs.2021.11.016>, 2022.
- Kaufman, Y. J., Tanré, D., Remer, L. A., Vermote, E. F., Chu, A., and Holben, B. N., Operational
remote sensing of tropospheric aerosol over land from EOS moderate resolution imaging
spectroradiometer, *J. Geophys. Res.*, 102(D14), 17051– 17067, doi:10.1029/96JD03988, 1997.
- 575 Kaufman, Y., Tanré, D. and Boucher, O. A satellite view of aerosols in the climate system. *Nature* 419,
215–223 doi.org/10.1038/nature01091, 2002.



- 580 Kaufman, Y. J., Koren, I., Remer, L. A., Rosenfeld, D., and Rudich, Y.: The effect of smoke, dust, and
pollution aerosol on shallow cloud development over the Atlantic Ocean, *P. Natl. Acad. Sci. USA*, 102,
11207–11212, <https://doi.org/10.1073/PNAS.05051911102>, 2005.
- .
- 585 Klüser, L., Martynenko, D. and Holzer-Popp, T., Thermal infrared remote sensing of mineral dust over
land and ocean: a spectral SVD based retrieval approach for IASI. *Atmos. Meas. Tech.*, 4(1), 461-509,
doi.org/10.5194/amt-4-757-2011, 2011.
- Koren, I., Kaufman, Y. J., Rosenfeld, D., Remer, L. A., and Rudich, Y. , Aerosol invigoration and
restructuring of Atlantic convective clouds, *Geophys. Res. Lett.*, 32, L14828,
590 [doi:10.1029/2005GL023187](https://doi.org/10.1029/2005GL023187), 2005.
- Koren, I., Martins, J. V., Remer, L. A., and Afargan, H.: Smoke Invigoration Versus Inhibition of
Clouds over the Amazon, *Science*, 321, 946–949, <https://doi.org/10.1126/science.1159185>, 2008.
- 595 LAADS DAAC (Level-1 and Atmosphere Archive & Distribution System Distributed Active Archive
Center) <https://ladsweb.modaps.eosdis.nasa.gov/about/>, referenced 05 February 2023.
- Lary, D.J., Remer, L.A., MacNeill, D., Roscoe, B. and Paradise, S., Machine learning and bias
correction of MODIS aerosol optical depth. *IEEE Geoscience and Remote Sensing Letters*, 6(4), 694-
698, [10.1109/LGRS.2009.2023605](https://doi.org/10.1109/LGRS.2009.2023605), 2009.
- 600 Lee, J., Shi, Y.R., Cai, C., Ciren, P., Wang, J., Gangopadhyay, A. and Zhang, Z., Machine learning
based algorithms for global dust aerosol detection from satellite images: inter-comparisons and
evaluation. *Remote Sensing*, 13(3), 456, doi.org/10.3390/rs13030456, 2021.
- 605 Levy, R.C., Remer, L.A., Mattoo, S., Vermote, E.F. and Kaufman, Y.J., Second-Generation Operational
Algorithm: Retrieval of Aerosol Properties over Land from Inversion of Moderate Resolution Imaging
Spectroradiometer Spectral Reflectance. *J. Geophys. Res. Atmos.*, 112, D13211,
<https://doi.org/10.1029/2006JD007811>, 2007.
- 610 Levy, R.C., Mattoo, S., Munchak, L.A., Remer, L.A., Sayer, A.M., Patadia, F. and Hsu, N.C., The
Collection 6 MODIS aerosol products over land and ocean. *Atmos. Meas. Tech.*, 6(11), 2989-3034,
doi.org/10.5194/amt-6-2989-2013
2013.
- Lipponen, A., Reinval, J., Väisänen, A., Taskinen, H., Lähivaara, T., Sogacheva, L., Kolmonen, P.,
615 Lehtinen, K., Arola, A. and Kolehmainen, V., Deep-learning-based post-process correction of the
aerosol parameters in the high-resolution Sentinel-3 Level-2 Synergy product. *Atmos. Meas.*
Tech., 15(4), 895-914, doi.org/10.5194/amt-15-895-2022, 2022.



- 620 Malm, W.C. and Hand, J.L., An examination of the physical and optical properties of aerosols collected
in the IMPROVE program. *Atmospheric Environment*, 41(16), 3407-3427,
doi.org/10.1016/j.atmosenv.2006.12.012, 2007.
- 625 Martonchik, J.V., Diner, D.J., Kahn, R.A., Ackerman, T.P., Verstraete, M.M., Pinty, B. and Gordon,
H.R., Techniques for the retrieval of aerosol properties over land and ocean using multiangle
imaging. *IEEE Transactions on Geoscience and Remote Sensing*, 36(4), 1212-1227,
10.1109/36.701027, 1998.
- 630 Martonchik, J.V., Diner, D.J., Crean, K.A. and Bull, M.A., Regional aerosol retrieval results from
MISR. *IEEE Transactions on geoscience and remote sensing*, 40(7), 1520-1531,
10.1109/TGRS.2002.801142, 2002.
- 635 Müller, D., Hostetler, C.A., Ferrare, R.A., Burton, S.P., Chemyakin, E., Kolgotin, A., Hair, J.W., Cook,
A.L., Harper, D.B., Rogers, R.R. and Hare, R.W., Airborne multiwavelength high spectral resolution
lidar (HSRL-2) observations during TCAP 2012: vertical profiles of optical and microphysical
properties of a smoke/urban haze plume over the northeastern coast of the US. *Atmos. Meas.*
Tech., 7(10), pp.3487-3496, doi.org/10.5194/amt-7-3487-2014, 2014.
- 640 Pappalardo, G., Amodeo, A., Apituley, A., Comeron, A., Freudenthaler, V., Linné, H., Ansmann, A.,
Bösenberg, J., D'Amico, G., Mattis, I. and Mona, L., EARLINET: towards an advanced sustainable
European aerosol lidar network. *Atmos. Meas Tech.*, 7(8), pp.2389-2409, doi.org/10.5194/amt-7-2389-
2014, 2014.
- Prucher, Jeff, 2006. *The Oxford Dictionary of Science Fiction*. Oxford University Press. ISBN-13:
9780195305678. On line: DOI:10.1093/acref/9780195305678.001.0001, accessed 29 May 2023.
- 645 Randles, C.A., Da Silva, A.M., Buchar, V., Colarco, P.R., Darmenov, A., Govindaraju, R., Smirnov,
A., Holben, B., Ferrare, R., Hair, J. and Shinozuka, Y., The MERRA-2 aerosol reanalysis, 1980 onward.
Part I: System description and data assimilation evaluation. *Journal of Climate*, 30(17), 6823-6850,
doi.org/10.1175/JCLI-D-16-0609.1, 2017.
- 650 Remer, L. A., Kaufman, Y. J., Tanré, D., Mattoo, S., Chu, D. A., Martins, J. V., Li, R. R., Ichoku, C.,
Levy, R. C., Kleidman, R. G., Eck, T. F., Vermote, E., and Holben, B. N., The MODIS aerosol
algorithm, products and validation, *J. Atmos. Sci.*, 62, 947–973, <https://doi.org/10.1175/JAS3385.1>,
2005.
- 655 Remer, L. A., Kaufman, Y. J. Aerosol direct radiative effect at the top of the atmosphere over cloud free
ocean derived from four years of MODIS data. *Atmos. Chem. Phys.*, 6, 237–253, doi.org/10.5194/acp-
6-237-2006, 2006.



- Remer, L. A., Kleidman, R.G., Levy, R.C., Kaufman, Y.J., Tanré, D., Mattoo, S., Martins, J.V., Ichoku, C., Koren, I., Yu, H., and Holben, B.N., Global aerosol climatology from the MODIS satellite sensors, *J. Geophys. Res.*, 113, D14S07, doi:10.1029/2007JD009661, 2008.
- Salomonson, V.V., Barnes, W.L., Maymon, P.W., Montgomery, H.E. and Ostrow, H., MODIS: Advanced facility instrument for studies of the Earth as a system. *IEEE Transactions on Geoscience and Remote Sensing*, 27(2), 145-153, 10.1109/36.20292, 1989.
- Schildhause, Chloe (November 25, 2014). "Back to the Future II's Art Director Tells Us How They Developed The Film's Somewhat Misguided Predictions". *Uproxx*. Retrieved May 2023.
- Snider, G., Weagle, C.L., Martin, R.V., Van Donkelaar, A., Conrad, K., Cunningham, D., Gordon, C., Zwicker, M., Akoshile, C., Artaxo, P. and Anh, N.X., SPARTAN: a global network to evaluate and enhance satellite-based estimates of ground-level particulate matter for global health applications. *Atmos. Meas. Tech.*, 8(1), 505-521, doi.org/10.5194/amt-8-505-2015, 2015.
- Song, Q., Zhang, Z., Yu, H., Kok, J. F., Di Biagio, C., Albani, S., Zheng, J., and Ding, J., Size-resolved dust direct radiative effect efficiency derived from satellite observations, *Atmos. Chem. Phys.*, 22, 13115–13135, <https://doi.org/10.5194/acp-22-13115-2022>, 2022.
- Tanré, D., M. Herman, Y. J. Kaufman, Information on the aerosol size distribution contained in the solar reflected spectral radiances, *J. Geophys. Res.*, 101, 19043– 19060, <https://doi.org/10.1029/96JD00333>, 1996.
- Tanré, D., Kaufman, Y. J., Herman, M., and Mattoo, S., Remote sensing of aerosol properties over oceans using the MODIS/EOS spectral radiances, *J. Geophys. Res.*, 102(D14), 16971– 16988, doi:10.1029/96JD03437, 1997.
- Ukkonen, P. Exploring pathways to more accurate machine learning emulation of atmospheric radiative transfer. *Journal of Advances in Modeling Earth Systems*, 14, e2021MS002875. doi.org/10.1029/2021MS002875, 2022
- Val Martin M, Kahn RA, Tosca MG. A Global Analysis of Wildfire Smoke Injection Heights Derived from Space-Based Multi-Angle Imaging. *Remote Sensing*, 10(10), 1609. doi-org.proxy-bc.researchport.umd.edu/10.3390/rs10101609, 2018.
- Van Donkelaar, A., Martin, R. V., Park, R. J. Estimating ground-level PM_{2.5} using aerosol optical depth determined from satellite remote sensing, *J. Geophys. Res.*, 111, D21201, doi:10.1029/2005JD006996, 2006.
- Van Donkelaar, A., Martin, R.V., Brauer, M., Hsu, N.C., Kahn, R.A., Levy, R.C., Lyapustin, A., Sayer, A.M., Winker, D.M. Global estimates of fine particulate matter using a combined geophysical-statistical



method with information from satellites, models, and monitors. *Environ. Sci. Tech.*, 50, 3762-3772, doi.org/10.1021/acs.est.5b05833, 2016.

Voss, K.K. and Evan, A.T., A new satellite-based global climatology of dust aerosol optical depth, *J. Appl. Meteorol. Clim.*, 59(1), 83-102, doi.org/10.1175/JAMC-D-19-0194.1, 2020.

700

Werdell, P.J., Behrenfeld, M.J., Bontempi, P.S., Boss, E., Cairns, B., Davis, G.T., Franz, B.A., Gliese, U.B., Gorman, E.T., Hasekamp, O., Knobelspiesse, K.D., Mannino, A., Martins, J.V., McClain, C.R., Meister, G., and Remer, L.A., The Plankton, Aerosol, Cloud, ocean Ecosystem mission: status, science, advances. *Bulletin of the American Meteorological Society*, 100(9), 1775-1794, doi.org/10.1175/BAMS-D-18-0056.1, 2019.

705

Yu, H., Kaufman, Y. J., Chin, M., Feingold, G., Remer, L. A., Anderson, T. L., Balkanski, Y., Bellouin, N., Boucher, O., Christopher, S., DeCola, P., Kahn, R., Koch, D., Loeb, N., Reddy, M. S., Schulz, M., Takemura, T., Zhou, M. A review of measurement-based assessment of the aerosol direct radiative effect and forcing., *Atmos. Chem. Phys.*, 6, 613-666, doi.org/10.5194/acp-6-613-2006, 2006.

710

Yu, H., Remer, L. A., Kahn, R. A., Chin, M., & Zhang, Y., Satellite perspective of aerosol intercontinental transport: From qualitative tracking to quantitative characterization, *Atmospheric Research*, 124, 73– 100. <https://doi.org/10.1016/j.atmosres.2012.12.013>, 2013.

715

Yu, H., Tan, Q., Chin, M., Remer, L. A., Kahn, R. A., Bian, H., et al., Estimates of African dust deposition along the trans-Atlantic transit using the decadelong record of aerosol measurements from CALIOP, MODIS, MISR, and IASI. *J. Geophys. Res. Atmos.*, 124, 7975– 7996, <https://doi.org/10.1029/2019JD030574>, 2019.

720

Yu, H., Yang, Y., Wang, H., Tan, Q., Chin, M., Levy, R. C., Remer, L. A., Smith, S. J., Yuan, T., and Shi, Y.: Interannual variability and trends of combustion aerosol and dust in major continental outflows revealed by MODIS retrievals and CAM5 simulations during 2003–2017, *Atmos. Chem. Phys.*, 20, 139–161, <https://doi.org/10.5194/acp-20-139-2020>, 2020.

725

Yuan, T., Remer, L. A., Pickering, K. E., and Yu, H.: Observational evidence of aerosol enhancement of lightning activity and convective invigoration, *Geophys. Res. Lett.*, 38, L04701, <https://doi.org/10.1029/2010GL046052>, 2011.

730

Yuan, T., Remer, L. A., and Yu, H.: Microphysical, macrophysical and radiative signatures of volcanic aerosols in trade wind cumulus observed by the A-Train, *Atmos. Chem. Phys.*, 11, 7119–7132, <https://doi.org/10.5194/acp-11-7119-2011>, 2011.

Zemeckis, Robert and Gale, Bob, 1989. *Back To the Future Part II*. Amblin Entertainment and Universal Pictures. Produced by Bob Gale and Neil Canton. 108 minutes.



- 735 Zhang, J., Christopher, S.A. Longwave radiative forcing of Saharan dust aerosols estimated from
MODIS, MISR, and CERES observations on Terra. *Geophys. Res. Lett.*, 30, 2188,
doi:10.1029/2003GL018479, 2003.
- Zhang, J., Reid, J.S., Westphal, D.L., Baker, N.L., Hyer, E.J. A system for operational aerosol optical
depth data assimilation over global oceans. *J. Geophys. Res. Atmos.*, 113, D10208,
740 doi:10.1029/2007JD009065, 2008.
- Zhang, J. and Reid, J. S.: A decadal regional and global trend analysis of the aerosol optical depth using
a data-assimilation grade over-water MODIS and Level 2 MISR aerosol products, *Atmos. Chem. Phys.*,
10, 10949–10963, doi.org/10.5194/acp-10-10949-2010, 2010.
- 745 Zheng, J., Zhang, Z., Garnier, A., Yu, H., Song, Q., Wang, C., Dubuisson, P., Di Biagio, C., The
thermal infrared optical depth of mineral dust retrieved from integrated CALIOP and IIR observations,
Remote Sensing of Environment, 270, (112841), 10.1016/j.rse.2021.112841, 2022.