

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Discovery of a variable energy-dependent X-ray polarization in the accreting neutron star GX 5–1

Sergio Fabiani¹, Fiamma Capitanio¹, Rosario Iaria², Juri Poutanen³, Andrea Gnarini⁴, Francesco Ursini⁴, Ruben Farinelli⁵, Anna Bobrikova³, James F. Steiner⁶, Jiri Svoboda⁷, Alessio Anitra², Maria C. Baglio^{8,9}, Francesco Carotenuto¹¹, Melania Del Santo¹², Carlo Ferrigno^{13,9}, Fraser Lewis^{14,15}, David M. Russell⁸, Thomas D. Russell¹², Jakob van den Eijnden¹⁰, Massimo Cocchi¹⁶, Alessandro Di Marco¹, Fabio La Monaca¹, Kuan Liu¹⁷, John Rankin¹, Martin C. Weisskopf¹⁸, Fei Xie^{17,1}, Stefano Bianchi⁴, Luciano Burderi^{19,20,16}, Tiziana Di Salvo^{2,20,16}, Elise Egron¹⁶, Giulia Illiano²¹, Philip Kaaret¹⁸, Giorgio Matt⁴, Romana Mikušincová⁴, Fabio Muleri¹, Alessandro Papitto²¹, Iván Agudo²², Lucio A. Antonelli^{21,23}, Matteo Bachetti¹⁶, Luca Baldini^{24,25}, Wayne H. Baumgartner¹⁸, Ronaldo Bellazzini²⁴, Stephen D. Bongiorno¹⁸, Raffaella Bonino^{26,27}, Alessandro Brez²⁴, Niccolò Bucciantini^{28,29,30}, Simone Castellano²⁴, Elisabetta Cavazzuti³¹, Chien-Ting Chen³², Stefano Ciprini^{33,23}, Enrico Costa¹, Alessandra De Rosa¹, Ettore Del Monte¹, Laura Di Gesu³¹, Niccolò Di Lalla³⁴, Immacolata Donnarumma³¹, Victor Doroshenko³⁵, Michal Dovčiak⁷, Steven R. Ehlert¹⁸, Teruaki Enoto³⁶, Yuri Evangelista¹, Riccardo Ferrazzoli¹, Javier A. Garcia³⁷, Shuichi Gunji³⁸, Kiyoshi Hayashida³⁹, Jeremy Heyl⁴⁰, Wataru Iwakiri⁴¹, Svetlana G. Jorstad^{42,43}, Vladimir Karas⁷, Fabian Kislat⁴⁴, Takao Kitaguchi³⁶, Jeffery J. Kolodziejczak¹⁸, Henric Krawczynski⁴⁵, Luca Latronico²⁶, Ioannis Liodakis⁴⁶, Simone Maldera²⁶, Alberto Manfreda⁴⁷, Frédéric Marin⁴⁸, Andrea Marinucci³¹, Alan P. Marscher⁴², Herman L. Marshall⁴⁹, Francesco Massaro^{26,27}, Ikuyuki Mitsui⁵⁰, Tsunefumi Mizuno⁵¹, Michela Negro^{52,53,54}, Chi-Yung Ng⁵⁵, Stephen L. O'Dell¹⁸, Nicola Omodei³⁴, Chiara Oppedisano²⁶, George G. Pavlov⁵⁶, Abel L. Peirson³⁴, Matteo Perri^{23,21}, Melissa Pesce-Rollins²⁴, Pierre-Olivier Petrucci⁵⁷, Maura Pilia¹⁶, Andrea Possenti¹⁶, Simonetta Puccetti²³, Brian D. Ramsey¹⁸, Ajay Ratheesh¹, Oliver J. Roberts³², Roger W. Romani³⁴, Carmelo Sgrò²⁴, Patrick Slane⁶, Paolo Soffitta¹, Gloria Spandre²⁴, Douglas A. Swartz³², Toru Tamagawa³⁶, Fabrizio Tavecchio⁹, Roberto Taverna⁵⁸, Yuzuru Tawara⁵⁰, Allyn F. Tennant¹⁸, Nicholas E. Thomas¹⁸, Francesco Tombesi^{59,33,60}, Alessio Trois¹⁶, Sergey S. Tsygankov³, Roberto Turolla^{58,61}, Jacco Vink⁶², Kinwah Wu⁶¹, and Silvia Zane⁶¹

(Affiliations can be found after the references)

Received ?; accepted ?

ABSTRACT

We report on the coordinated observations of the neutron star low-mass X-ray binary (NS-LMXB) GX 5–1 in X-rays (*IXPE*, *NICER*, *NuSTAR* and *INTEGRAL*), optical (REM and LCO), near-infrared (REM), mid-infrared (VLT VISIR), and radio (ATCA). This Z-source was observed by *IXPE* twice in March–April 2023 (Obs. 1 and 2). In the radio band, the source was detected, but only upper-limits to the linear polarization were obtained at a 3σ level of 6.1% at 5.5 GHz and 5.9% at 9 GHz in Obs. 1 and 12.5% at 5.5 GHz and 20% at 9 GHz in Obs. 2. The mid-IR, near-IR and optical observations suggest the presence of a compact jet which peaks in the mid- or far-IR. The X-ray polarization degree was found to be $3.7\% \pm 0.4\%$ (at 90% confidence level) during Obs. 1 when the source was in the horizontal branch of the Z-track and $1.8\% \pm 0.4\%$ during Obs. 2 when the source was in the normal-flaring branch. These results confirm the variation of polarization degree as a function of the position of the source in the color-color diagram as for previously observed Z-track sources (Cyg X-2 and XTE 1701–462). Evidence for a variation of the polarization angle $\sim 20^\circ$ with energy is found in both observations, likely related to the different, non-orthogonal polarization angles of the disk and Comptonization components which peak at different energies.

Key words. accretion, accretion disks – neutron stars – X-rays: general – X-rays: binaries – X-rays: individual: GX 5–1

1. Introduction

Persistent neutron star low-mass X-ray binaries (NS-LMXBs) are among the X-ray astronomical objects that the *Imaging X-ray Polarimetry Explorer* (*IXPE*, Weisskopf et al. 2023, 2022; Soffitta et al. 2021) is investigating. Four of them were already observed by *IXPE* during its first year campaign, namely the Z-source Cyg X-2, the peculiar Z-Atoll transient XTE J1701–462, and two bright soft-state Atoll-sources, GS 1826–238 and GX 9+9. The source classification (Z or Atoll) is based on

the tracks that they draw on the color-color-diagram (CCD; see, e.g., Hasinger & van der Klis 1989; van der Klis 1995). GS 1826–238 data are compatible with a null polarization with an upper limit on the polarization degree (PD) of 1.3% (Capitanio et al. 2023), while Cyg X-2 (Farinelli et al. 2023) and GX 9+9 (Chatterjee et al. 2023; Ursini et al. 2023) have shown a statistically significant linear polarization with the PD of $\sim 2\%$ and $\sim 1.5\%$, respectively. Strong variations in PD was reported for XTE J1701–462, observed twice, that showed a high PD \sim

4.5% in the first observation and a PD compatible with a null polarization in the second one (Cocchi et al. 2023).

GX 5–1 is a Galactic Z-source (Kuulkers et al. 1994; Jonker et al. 2002) located near the Galactic Center. It is a radio source with radio emission most likely originating from a compact jet (Fender & Hendry 2000). The radio counterpart allowed an accurate localization that, despite optical obscuration and crowded field near the Galactic Center, has led to the determination of a likely infra-red companion candidate (Jonker et al. 2000). Until the early 1990’s GX 5–1 X-ray data were likely contaminated by the black hole LMXB GRS 1758–258, located only 40’ away. Sunyaev et al. (1991) and Gilfanov et al. (1993) were able to resolve two sources, showing that GX 5–1 was ~ 30 –50 times brighter than GRS 1758–258 below 20 keV. GX 5–1 has not shown any X-ray pulsations or X-ray bursts (Paizis et al. 2005).

An X-ray halo due to scattering of X-ray photons is clearly revealed in the *Chandra* image (Smith et al. 2006; Clark 2018). Such a halo arises due to the presence of multiple clouds along the line of sight.

In X- and γ -rays, Paizis et al. (2005) studied one year of *INTEGRAL* data, which covered all the Z-track of the source (mainly the horizontal branch (HB) and normal branch (NB)). ISGRI and JEM-X average spectra showed a clear hard X-ray emission above 20 keV, not detected previously and compatible with thermal Comptonization of soft photons from a hot, optically thin plasma in the vicinity of the NS. However, Paizis et al. (2005) were not able to constrain the temperature of the Comptonizing plasma. They assessed the compatibility of GX 5–1 energy spectrum with the so called ‘eastern’ (Mitsuda et al. 1984) and ‘western’ (White et al. 1986) models and found the former being physically more meaningful, describing the spectral ‘flattening’ above ~ 20 keV as a Comptonized hard-tail emission. Paizis et al. (2006) described the GX 5–1 energy spectrum in the 20–100 keV energy band observed by *INTEGRAL* IBIS/ISGRI with a Comptonization component (comptt, Titarchuk (1994)) plus a power law to account for the hard X-ray emission. Such a high-energy tail was first detected by Asai et al. (1994) although a possible contamination from the nearby black hole GRS 1758–258 could not be excluded in this latter case.

Paizis et al. (2006) highlighted the presence of a correlation between the X-ray spectral states and the radio emission. Steady radio emission is associated with low/hard state (typical for Atoll sources; Fender & Hendry 2000; Migliari & Fender 2006). These sources brighten considerably during intermediate states, often showing bright, transient flares (HB of Z-sources), before quenching during the very soft states in the NB and FB branches of Z-sources (Fender & Hendry 2000; Di Salvo & Stella 2002). Moreover, Paizis et al. (2006) reported that the radio flux is positively correlated with the flux of the hard tail in the 40–100 keV energy range. They suggested that this correlation is related to the acceleration of electrons along open field lines in the NS magnetosphere at the base of the jet seen in the radio. Berendsen et al. (2000) reported upper limits to the radio linear polarization of 33% at 6.3 cm and 23% at 3.5 cm not sufficient to constrain the emission mechanism or the optical depth of the jet.

2. Observations and data reduction

2.1. IXPE

The X-ray polarimeter *IXPE* (Weisskopf et al. 2023, 2022; Soffitta et al. 2021) observed GX 5–1 twice (Obsid 02002799) from March 21 to 22 and from April 13 to 15, 2023 (see Table 1

and Fig. 1 for the light curves) with a nominal integration time of 50 ks each. *IXPE* provides timing, imaging, spectroscopic, and polarimetric data in the 2–8 keV band. Data reduction and analysis were performed by means of the *IXPEOBSSIM* software version 30.5.0 (Baldini et al. 2022) and the *HEASOFT* package version 6.31.1 (Nasa Heasarc 2014). Data were filtered by using *IXPEOBSSIM* tools *xpselect* and binned¹ with *xpbin* to produce images and *I*, *Q*, and *U* energy spectra for spectro-polarimetric analysis performed with *XSPEC* version 12.13.0c (Arnaud 1996). We used the latest version 12 of the *IXPE* response matrices available at the *IXPEOBSSIM* public repository² (also available at the HEASARC archive). A circular source extraction region of 60’’ in radius was selected from the image for each one of the three detector units (DUs). No background subtraction was applied due to the high count rate of the source (~ 20 –25 cts s⁻¹ per DU) (see Di Marco et al. 2023). Only *XSPEC* allows currently a ‘weighted analysis’ (Di Marco et al. 2022) in which a weight is assigned to each photo-electron track recorded by the DUs depending on the shape of the charge distribution.

The normalized Stokes parameters $q = Q/I$ and $u = U/I$, the PD and polarization angle (PA) with their uncertainties can be calculated by using the model-independent *pcube* binning algorithm of *IXPEOBSSIM*. On the other hand, PD and PA obtained with *XSPEC* require the definition of a spectro-polarimetric model. Because the PD and PA are not independent, the appropriate way to report results is by means of contour plots at certain confidence levels for the joint measurements of both parameters. We report *XSPEC* (PD,PA) contour plots as obtained by using the *steppar* command, while contour plots associated to the *IXPEOBSSIM* analysis are obtained from the statistics of the number of counts (Weisskopf et al. 2010; Strohmayer & Kallman 2013; Muleri 2022).

The spectro-polarimetric analysis was carried out by taking into account the current *IXPE* effective area instrument response function (*arf*) which may not be as accurate as possible at energies above 6 keV where there is a significant roll-off in the spectral response. Because the high-energy part of the *IXPE* band is of special interest for our study, we estimate the *IXPE* spectral systematic uncertainties to be as 3% in Obs. 1 and 2% in Obs. 2, based on the comparison with *NICER*, *NuSTAR* and *IBIS/ISGRI* energy spectra. The spectral models for both the *IXPE* observations are frozen based on the spectral analysis of these other observatories. We used the *XSPEC* gain fit tool for the *IXPE* data only, to shift the energies on which the response matrix is defined and to match the effective area curve during the fit procedure. The spectro-polarimetric fit of the *IXPE* data was carried out freeing the three DU’s normalization constants, the gain slope, gain offset and the polarimetric parameters.³

2.2. NuSTAR

GX 5–1 was observed by *NuSTAR* (Harrison et al. 2013) on March 21 and twice on April 13–14, 2023. All the relevant observation times, sequence IDs and exposure times are provided in Table 1 and the corresponding light curves are shown in Fig. 1. The unfiltered event files were processed with the *NuSTAR* Data Analysis Software (*NUSTARDAS* v.2.1.2) to produce the cleaned and calibrated level 2 data us-

¹ The default energy binning is 40 eV.

² <https://github.com/lucabaldini/ixpeobssim>

³ The normalization constants, gain slope and offset parameters are identical between *I*, *Q* and *U* spectra of the same DU. The PA and PD parameters of those spectra are identical also for each DU.

Table 1. List of observations.

Telescope	Obsid	Obs. Start	Obs. Stop	Net Exposure (ks)	Notes
<i>IXPE</i> Obs. 1					
<i>IXPE</i>	02002799	2023-03-21, 04:16:14	2023-03-22, 05:02:52	48.6	obs. segment 1
NICER	6010230101/2	2023-03-21, 03:41:20	2023-03-22, 04:50:20	13.1	
<i>NuSTAR</i>	90902310002	2023-03-21, 16:41:48	2023-03-22, 08:04:34	12.6	
<i>INTEGRAL</i>	2070006/0001	2023-03-21, 03:58:07	2023-03-22, 04:28:46	40.1	
REM	–	2023-03-22, 05:25:24	2023-03-22, 07:33:05	–	see Sect. 2.7 for exp. details
LCO	–	2023-03-22, 16:33:07	2023-03-22, 16:38:07	–	"
No <i>IXPE</i> Obs.					
NICER	6010230103/4	2023-03-24 18:19:20	2023-03-25 00:44:20	5.2	
ATCA	–	2023-03-24, 15:21:20	2023-03-25, 01:55:50	–	see Sect. 2.5 for exp. details
VISIR	110.2448	2023-03-28, 08:31:00	2023-03-28, 09:21:00	–	see Sect. 2.6 for exp. details
LCO	–	2023-03-29, 15:50:11	2023-03-31, 15:43:07	–	see Sect. 2.7 for exp. details
VISIR	110.2448	2023-03-31, 07:46:00	2023-03-31, 08:42:00	–	see Sect. 2.6 for exp. details
<i>IXPE</i> Obs. 2					
<i>IXPE</i>	02002799	2023-04-13, 23:43:42	2023-04-15, 00:37:32	47.1	obs. segment 2
NICER	6010230105/6	2023-04-13 16:39:04	2023-04-15, 23:58:20	13.1	
<i>NuSTAR</i>	90902310004/6	2023-04-13, 15:57:07	2023-04-14, 23:23:24	15.7	
<i>INTEGRAL</i>	multiple	2023-04-13, 03:51:17	2023-04-15 08:53:00	85.0	
REM	–	2023-04-14, 03:16:55	2023-04-14, 06:31:56	–	see Sect. 2.7 for exp. details
LCO	–	2023-04-14, 05:35:19	2023-04-14, 23:23:55	–	"
ATCA	–	2023-04-14, 11:32:00	2023-04-14, 17:57:40	–	see Sect. 2.5 for exp. details

Notes. The three table sections comprise the observations related to *IXPE* Obs. 1, Obs. 2, and observations in between performed with other observatories.




Fig. 1. Light curves of *NuSTAR*, NICER and IBIS/ISGRI during *IXPE* Obs. 1 (left panel) and Obs. 2. (right panel). On the top row of each panel the three *IXPE* DU's light curves are shown (DU1 blue, DU2 orange, and DU3 green). In the left panel the NICER soft purple and dark red points correspond to ObsID 6010230101 and 6010230102. In the right panel the *NuSTAR* red and purple points correspond to ObsID 90902310004 and 90902310006. The NICER gray points correspond to ObsID 6010230106.

ing the latest calibration files (CALDB v.20221130) and the standard filtering criteria with the `nupipeline` task, where `statusexpr="STATUS==b00000xxx00xxx000"` was set due to the source flux exceeding 100 counts s^{-1} . The spectra and light curves were extracted using the `nuproducts` task, selecting a circular region of $60''$ in radius centered on the source.

2.3. NICER

NICER (Gendreau et al. 2016) observed GX 5–1 between March 21–25 and April 13–14, 2023. The observations identi-

fied with ID 6010230101, 6010230102, and 6010230106 were included in the spectro-polarimetric analysis, because they are simultaneous with the *IXPE* observations. The calibrated and cleaned files were extracted by using the standard `nicer12` command of the NICER Data Analysis Software (NICERDAS v.10) together with the latest calibration files (CALDB v.20221001). The spectra and the light curves were then obtained with the `nicer13-spect` and `nicer13-lc` tasks, while the background was computed using the SCORPEON⁴ model. Light curves were

⁴ https://heasarc.gsfc.nasa.gov/docs/nicer/analysis_threads/scorpeon/

obtained including 50 NICER FPMs available during the ObsIDs included in the analysis (out of 52 available, excluding noisy detectors ID. 14 and ID. 34). Thus, no discontinuities in the NICER light curves are present (see Fig. 1).

During the contemporary observation with *IXPE* there were two significant increases of count rate in the NICER data, up to the double of the typical value. These events were coincident with two solar flares. The first one was a C class peaking at about 60048.5389 MJD and the second one was an M class peaking at about MJD 60048.6806. We removed them manually from the GTIs (the C class flare from MJD 60048.53535 to MJD 60048.60033 and the M class flare from MJD 60048.67506 to MJD 60048.73744).

We noticed some relevant features in the energy spectrum below 4 keV, most likely due to spectral features unaccounted for in the NICER ARF. Because of the source's high count rate, such features become apparent in the spectral modeling. We therefore accounted for calibration artifacts owing to imperfections in modeling the dead layer of the silicon detector at the Si-K edge, and of the concentrator mirror surface roughness at the Au-M edges, which affects the ≈ 2.2 – 3.5 keV range.⁵ We froze the energy of those edges at their best-fit value as reported in Table 4.

2.4. INTEGRAL IBIS/ISGRI

INTEGRAL IBIS/ISGRI (Winkler et al. 2003) observed the region of GX 5–1 from March 21 to 22 and from April 13 to 15, 2023, responding to a request by the *IXPE* team. The data for this source are publicly available and were reduced for the imager IBIS (Ubertini et al. 2003) and the ISGRI detector (Lebrun et al. 2003). We used the MMODA⁶ platform that empowers the Off-Line Science Analysis (OSA) version 11.2 distributed by the ISDC (Courvoisier et al. 2003) with the most recent calibration files that are continuously ingested in the Instrument Characteristic repository. We first built a mosaicked image of all individual pointings that constitute the standard dithering strategy of observation for IBIS/ISGRI in the 28–40 keV energy range. These images were used to make the catalog of detected sources with a signal-to-noise ratio larger than 7. Using these catalog, we extracted light curves with 1000 s time bins and spectra in 256 standard channels for IBIS/ISGRI, these were grouped in 10 equally spaced logarithmic channels between 28 and 150 keV. We also accounted for a systematic uncertainty of the spectra at the 1.5% level. The equivalent on-axis exposures of the IBIS/ISGRI spectra are 40 and 85 ks, respectively, after correction for dead time and vignetting. Products are available at the *INTEGRAL* Product gallery.⁷

2.5. ATCA

The Australia Telescope Compact Array (ATCA) observed GX 5–1 on March 24 and April 14, 2023. On March 24, the telescope observed with the array in its 750C configuration.⁸ Observations taken on April 14 were carried out with the array in a relatively compact H214 configuration, in combination with an isolated antenna located 6 km from the array core, which was also included for our analysis. For both observations, data were recorded simultaneously at central frequencies of 5.5 GHz and 9.0 GHz, with 2 GHz of bandwidth at each frequency.

We used PKS 1934–638 for bandpass and flux density calibration. PKS 1934–638 was also used to solve for the antenna leakages (D-terms) for the polarization calibration. The nearby source B1817–254 was used for gain calibration and to calibrate the PA using the Common Astronomy Software Applications for radio astronomy (CASA, version 5.1.2; CASA Team et al. 2022) `atcapolhelpers.py` task `qufromgain`.⁹ Calibration and imaging followed standard procedures within CASA. When imaging, we used a Briggs robust parameter of 0 to balance sensitivity and resolution (Briggs 1995), as well as suppress the effects from some bright, diffuse emission within the field.

For our March 24 observations, fitting for a point source in the image plane, we detect GX 5–1 at a flux density of $960 \pm 19 \mu\text{Jy}$ at 5.5 GHz and $810 \pm 11 \mu\text{Jy}$ at 9 GHz coincident with the previously reported radio the X-ray position (e.g., Berendsen et al. 2000; Liu et al. 2007). These detections correspond to a radio energy spectral index of -0.37 ± 0.09 . The Stokes Q and U values were measured at the position of the peak source flux density (Stokes I). No significant linearly polarized (LP) emission was detected at either frequency. Measuring the root mean square of the image noise in a $50'' \times 50''$ region over the source position (taken as 1σ), provides 3σ upper limits on the polarized intensity $\sqrt{Q^2 + U^2}$ of $58 \mu\text{Jy beam}^{-1}$ at 5.5 GHz and $48 \mu\text{Jy beam}^{-1}$ at 9 GHz. These corresponds to a 3σ upper limit on the PD of 6.1% at 5.5 GHz and 5.9% at 9 GHz. Stacking the two frequencies to maximize the sensitivity also yields a non-detection of linearly polarized emission, with a 3σ upper-limit of 4.2% (centered at 7.25 GHz).

On April 14, following the same calibration and imaging procedure, we measured the flux density of GX 5–1 to be $750 \pm 50 \mu\text{Jy}$ and $620 \pm 40 \mu\text{Jy}$ at 5.5 and 9 GHz, respectively. These detections correspond to a radio spectral index of -0.4 ± 0.1 . We note that due to the more compact array configuration for this epoch, and the presence of diffuse emission in the field, we imaged with a strictly uniform weighting scheme (setting the Briggs robust parameter to -2), reducing the impact of diffuse emission in the field on our resultant images. The compact configuration coupled with a shorter exposure time did result in a higher noise level in the images. At 5.5 GHz, we do not detect any linearly polarized emission, with a 3σ upper-limit on the PD of 12.5%. Similarly, at 9 GHz we measure a 3σ upper-limit on the PD of 20%. Stacking the two frequencies places a 3σ upper-limit on the PD of 8% at 7.25 GHz. We do note that during the final ~ 30 min of the observation, some linear polarization was detected from close to the source position, but only at 9 GHz. Due to the non-detection at 5.5 GHz and the short and sudden nature of this emission, we attribute it to radio frequency interference and not an astrophysical event.

2.6. VLT VISIR

Mid-IR observations of the field of GX 5–1 were made with the European Southern Observatory's Very Large Telescope (VLT) on March 28 and 31 2023 under the program 110.2448 (PI: D. Russell). The VLT Imager and Spectrometer for the mid-Infrared (VISIR; Lagage et al. 2004) instrument on the VLT was used in small-field imaging mode. Four filters (M -band, $J8.9$, $B10.7$ and $B11.7$) were used, with central wavelengths of 4.67, 8.70, 10.64, and $11.51 \mu\text{m}$, respectively. For each observation, the integration time on source was composed of a number of nodding cycles, alternating between source and sky. The total observing time was usually almost twice the integration time.

Figure 1: *INTEGRAL* IBIS/ISGRI and *IXPE* observation times. The total observing time was usually almost twice the integration time.

⁵ https://heasarc.gsfc.nasa.gov/docs/nicer/analysis_threads/archiving

⁶ <https://www.astro.unige.ch/mmoda/>

⁷ <https://www.astro.unige.ch/mmoda/gallery/astrophysical-entity/gx-5-1>

⁸ https://www.narrabri.atnf.csiro.au/operations/array_configuration/arrays/atca.html




Fig. 2. Polarization of GX 5–1 measured by *IXPE* in the 2–8 keV energy range during Obs. 1 and Obs. 2 obtained with *xspec* and *IXPEOBSSIM*. Contours are computed for two parameters of interest at 50%, 90%, and 99.9% confidence levels.

Observations of standard stars were made on the same nights as the target, in the same filters (photometric standards HD137744, HD169916, HD130157 and HD145897 were observed, all at airmass 1.0–1.1). Conditions were clear on both nights. All data (target and standard stars) were reduced using the VISIR pipeline in the *GASGANO* environment.¹⁰ Raw images from the chop/nod cycle were recombined. Photometry was performed on the combined images using PHOT in IRAF.¹¹ For the B10.7 filter, two standard stars were observed on each night, just before and after each observation of GX 5–1, to check for stability. We found that the counts/flux ratio values from these standards agree to a level of 6.4%, which we adopt as the systematic error of all flux measurements. The estimated counts/flux ratio in each filter was used to convert count rates (or upper limits) of GX 5–1 to flux densities.

On 2023-03-28 (MJD 60031.37), the B10.7 filter was used only, and we derive a 3σ flux density upper limit of 3.27 mJy at $10.64 \mu\text{m}$ (the airmass was 1.05–1.10). On 2023-03-31 (MJD 60034.34), GX 5–1 was detected in *M*-band and *J*8.9, with flux densities of 4.2 ± 1.4 mJy at $4.67 \mu\text{m}$ and 10.0 ± 2.3 mJy at $8.70 \mu\text{m}$, respectively (at airmass 1.07–1.17). The significance of the detection was 5.9σ in both filters. The errors on the fluxes incorporate the statistical error on each detection, and the systematic error from the standard stars, in quadrature. The source was not detected in the B10.7 and B11.7 filters on 2023-03-31, with flux upper limits that were less constraining than on 2023-03-28.

GX 5–1 lies in a crowded region of the Galactic plane, with several stars detected within $5''$ of the source. The near-IR counterpart was confirmed through photometry and spectroscopy (named as star 513; Jonker et al. 2000; Bandyopadhyay et al. 2003), and its coordinates agree with the radio and X-ray position. We are confident that the source we detect with VISIR is indeed GX 5–1, because star 503 from Jonker et al. (2000) and Bandyopadhyay et al. (2003) is also detected (at a low significance of 3.1 – 3.3σ) at the correct coordinates. This star is estimated from Fig. 1 of Jonker et al. (2000) to lie $4''.3$ to the south-east of GX 5–1. In the *J*8.9 VISIR image the detected source is measured to be $4''.28$ to the south-east of the position of the detected GX 5–1, as expected. The flux density of star 503 is 1.4 ± 0.5 mJy in *M*-band and 4.5 ± 2.2 mJy in *J*8.9.

2.7. REM and LCO

Optical (SDSS *griz* filters) and near infrared (NIR; 2MASS *H*-band) observations of GX 5–1 were acquired with the robotic 60 cm Rapid Eye Mount (REM; Zerbi et al. 2001; Covino et al. 2004) telescope on March 22 and on April 14, 2023 (see Table 1). Strictly simultaneous observations were obtained in all bands; on March 22, a series of 150 30-s exposures were acquired in *H*-band (dithering was applied), and 20 300-s exposures in each of the optical bands; on April 14, a series of 225 30-s exposures were acquired in *H*-band (dithering was applied), and 30 300-s exposures in each of the optical bands.

Optical observations were also acquired with the 1 m and 2 m telescopes of the Las Cumbres Observatory (LCO) network, using the *i'* and *Y* filters. Observations with the 2 m telescope (Siding Spring - Faulkes Telescope South) were performed on 2023-03-22T16:33:07, 2023-03-29T15:50:11, 2023-03-30T15:46:39 and 2023-03-31T15:43:07 (300s integration in all epochs and bands), and with the 1 m telescopes (at the locations of Cerro Tololo, Siding Spring and Sutherland) on 2023-03-21T07:09:39, 2023-03-22T01:01:36, 2023-03-22T08:11:35, 2023-04-14T05:35:19, T08:11:37, T15:11:30, T23:23:55 (300s integration in all epochs and bands).

The optical images were bias- and flat-field corrected using standard procedures; the contribution of the sky in the NIR images was evaluated by performing a median of the dithered images five-by-five, and was then subtracted from each image. In all bands and epochs, all reduced images were then aligned and averaged in order to increase the signal-to-noise. Aperture photometry was performed using PHOT in IRAF. Flux-calibration was performed against a group of 6 stars with magnitudes tabulated in the 2MASS catalog¹² and 6 stars from the PanSTARRS catalog.¹³

Due to the very high extinction of the source ($N_H = 4.93 \times 10^{22} \text{ cm}^{-2}$ and $4.88 \times 10^{22} \text{ cm}^{-2}$, which translates¹⁴ into $A_V = 17.2$ mag and 17 mag in Obs. 1 and Obs. 2, respectively) and to the combination of the low spatial resolution of our images and the crowded field of the source, GX 5–1 is not detected in any of the optical and NIR images acquired. A blend of our target with at least one of the nearby stars can be detected at very low significance in the averaged *H*-band image, at a position consistent with the proposed optical and NIR counterpart of the source (Jonker et al. 2000, star 513). However, this detection is not significant enough to extract a flux for the blend.

We estimate the following 3σ upper limits (only the most constraining ones per epoch are quoted): $H = 13.84$, $Y = 19.08$, $z = 18.28$, $i' = 21.79$ (LCO), $r' = 20.22$, $g' = 20.45$ for Obs 1, $H = 14.00$, $Y = 18.86$, $z = 18.35$, $i' = 21.22$, $r = 20.46$, $g = 20.50$ for Obs 2. These upper limits are consistent with the *H*-band magnitude of the proposed NIR counterpart reported by Jonker et al. (2000) (Star 513; $H = 14.1 \pm 0.2$).

3. Results

3.1. Polarimetric model-independent analysis

The model-independent analysis of the X-ray polarization with *ixpeobssim* (see Table 2) in the 2–8 keV energy band gives the PD of $4.3\% \pm 0.3\%$ and $2.0\% \pm 0.3\%$ in Obs. 1 and Obs. 2, respectively (with 1σ errors). The corresponding PAs are -9.7 ± 2.0

¹⁰ <https://www.eso.org/sci/software/gasgano.html>

¹¹ IRAF is distributed by the National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation

¹² <https://irsa.ipac.caltech.edu/Missions/2mass.html>

¹³ <https://catalogs.mast.stsci.edu/panstarrs/>

¹⁴ See Sect. 3.5 and Foight et al. (2016a).




Fig. 3. Stokes parameters as a function of energy for *IXPE* Obs. 1 and Obs. 2 (left panels) and for the combined data set (right panels) obtained with *IXPEOBSSIM*. The normalized Stokes q parameter is compatible with a constant value in each one of the *IXPE* observations, whereas the Stokes u parameter is not (see Table 3). This behavior of the Stokes parameters is consistent with a variation of the PA (see Fig. 4).

Table 2. Polarization in the 2–8 keV band (see Fig. 2) and in 4 energy bins estimated with *IXPEOBSSIM* for Obs. 1 and Obs. 2.

Energy (keV)	Parameters	Obs. 1	Obs. 2
2–8	PD (%)	4.3 ± 0.3	2.0 ± 0.3
	PA (deg)	-9.7 ± 2.0	-9.2 ± 4.0
2–3	PD (%)	4.0 ± 0.5	2.0 ± 0.4
	PA (deg)	0.2 ± 3.4	6.0 ± 6.2
3–4	PD (%)	3.8 ± 0.4	1.7 ± 0.3
	PA (deg)	-14.4 ± 3.8	-3.5 ± 6.5
4–5	PD (%)	4.4 ± 0.5	2.5 ± 0.5
	PA (deg)	-9.8 ± 3.5	-19.2 ± 6.0
5–8	PD (%)	5.4 ± 0.7	2.6 ± 0.7
	PA (deg)	-14.0 ± 3.7	-21.0 ± 7.9

Notes. The errors are at 68% confidence level.

and -9.2 ± 4.0 . We see a significantly larger PD in Obs. 1 compared to Obs. 2. However, the PA is compatible with being constant during the two observations. Contour plots of the *IXPEOBSSIM* analysis in the 2–8 keV energy range are reported in Fig. 2 (dashed lines).

The behavior of polarization as a function of energy is reported in Table 2. The energy dependence of the PD is essentially the same in the two observation, whereas the PA varies from positive to negative values. A proper assessment of this behavior requires the use of Stokes parameters. The left panels of Fig. 3 show the Stokes parameters as a function of energy of both the observations separately. For each observation, the Stokes q parameters are consistent with being constant in energy, albeit at different values between the first and second observations. On

Table 3. Results of the Stokes parameters fit with a constant value for Obs. 1, Obs. 2, and the combined data set.

Stokes parameter	Best-fit parameters	Obs. 1	Obs. 2	Obs. 1 + Obs. 2
q	constant (%)	3.86 ± 0.25	1.86 ± 0.24	2.83 ± 0.17
	χ^2_ν	1.31	0.15	1.26
	α (%)	26.8	93.0	28.7
	constant (%)	-1.32 ± 0.25	-0.74 ± 0.24	-0.89 ± 0.17
u	χ^2_ν	4.26	4.00	7.16
	α (%)	0.52	0.74	0.01

Notes. The reduced χ^2 of the fit is χ^2_ν . The significance level of the χ^2 value is α . The number of degrees of freedom is 3.

the other hand, the Stokes u parameters are not compatible with a constant (see Obs. 1 and Obs. 2 columns of Table 3). This requires the variation of the PA.

On the basis of the assumption that the geometry and the physical process producing polarization are the same in the two observations, we also calculated the Stokes parameters of the *IXPE* observations combined (Obs. 1 and Obs. 2) to improve the statistics. While the resulting normalized Stokes parameter q still remains compatible with a constant value, the Stokes u is even more far from being a constant with the reduced χ^2 values for $\nu = 3$ degrees of freedom being $\chi^2_\nu = 1.26$ and $\chi^2_\nu = 7.16$ for q and u , respectively. As anticipated by the separate analysis of the two observations, this behavior of q and u implies a variation of the PA. Such a variation by about 20° is highlighted in Fig. 4. In the top panel the contour plots on the PD–PA plane for the *IXPE* whole observation are shown. Polarization in the 2–3 keV energy bin is not compatible with that in the 5–8 keV bin with a probability of $\sim 98.7\%$. In the bottom panel the variation of the




Fig. 4. Polarization contour plot (top panel) and PA (bottom panel) as a function of energy of the combined *IXPE* data set (Obs. 1 and Obs. 2).

PA with energy is shown together with the fit by a constant giving a value of -9.1 ± 1.6 with $\chi^2_\nu = 6.44$ for $\nu = 3$ corresponding to a significance level $\alpha = 0.02\%$.

3.2. X-ray spectral analysis

The NICER, *NuSTAR* and IBIS/ISGRI light curves contemporary to *IXPE* observations are shown in Fig. 1. The time-resolved CCD and hardness-intensity diagram (HID) for all the observations with *IXPE*, NICER and *NuSTAR* show GX 5–1 moving along the complete Z-track from March 21 to April 14. *NuSTAR* CCD and HID, obtained from the GTIs contemporary to the *IXPE* observations (see Fig. 5), highlight clearly the Z shape, disentangling also the NB with respect to the FB, due to the wide energy band from 3 to 20 keV used to construct the CCD colors. During *IXPE* Obs. 1, the source was in the HB, whereas it was in the NB-FB during Obs. 2 which we have checked also by constructing CCD and HID from the *IXPE* data of both observations. From March 24 to 25 (about 3 days after the end of Obs. 1), when ATCA was observing, NICER detected GX 5–1 moving from the HB-NB corner towards the NB.

We fit the NICER, *NuSTAR* and IBIS/ISGRI data of Obs. 1 and Obs. 2 presenting results in Fig. 6 and Table 4. The better

Table 4. Best-fit parameters of GX 5–1 spectral model from NICER, *NuSTAR* and IBIS/ISGRI data simultaneous to *IXPE* observation.

Components	Parameters	Obs. 1	Obs. 2
edge	E (keV)	1.82148 (frozen)	
	τ	$0.172^{+0.028}_{-0.016}$	0.152 ± 0.017
edge	E (keV)	1.95197 (frozen)	
	τ	0.051 ± 0.016	0.484 ± 0.016
edge	E (keV)	2.28003 (frozen)	
	τ	0.037 ± 0.014	0.025 ± 0.014
edge	E (keV)	2.44444 (frozen)	
	τ	0.050 ± 0.013	0.038 ± 0.013
edge	E (keV)	3.16139 (frozen)	
	τ	$0.020^{+0.007}_{-0.008}$	0.012 ± 0.007
Continuum parameters			
tbabs	N_H (10^{22} cm^{-2})	$4.93^{+0.12}_{-0.06}$	4.88 ± 0.04
diskbb	kT (keV)	0.95 ± 0.07	1.20 ± 0.03
	$R_{\text{in}} \sqrt{\cos \theta}$ (km) ^a	25^{+2}_{-3}	$19.6^{+0.9}_{-0.8}$
thcomp	Γ	$2.35^{+0.13}_{-0.49}$	< 2.1
	kT_e (keV)	$2.99^{+0.02}_{-0.10}$	$3.08^{+0.37}_{-0.11}$
	f	$0.99^{+0.01}_{-0.36}$	$0.032^{+0.08}_{-0.005}$
bbodyrad	kT_{bb} (keV)	$1.27^{+0.26}_{-0.11}$	$1.68^{+0.03}_{-0.04}$
	R_{bb} (km) ^a	19^{+1}_{-6}	$9.3^{+0.6}_{-0.5}$
expabs	E_{cut} (keV)	$[= kT_{\text{bb}}]$	–
powerlaw	Γ	$2.67^{+0.49}_{-0.92}$	–
	norm	$0.45^{+0.23}_{-0.43}$	–
Cross-calibration constants			
const	C_{NICER}	1.107 ± 0.003	1.034 ± 0.004
	$C_{\text{NuSTAR,FPMA}}$	1 (frozen)	
	$C_{\text{NuSTAR,FPMB}}$	1.0136 ± 0.0013	0.9908 ± 0.0014
	$C_{\text{IBIS/ISGRI}}$	0.66 ± 0.08	1.29 ± 0.03
	$\chi^2/\text{d.o.f.}$	421/451	338/365
	τ_{thcomp}^b	$8.7^{+2.5}_{-0.7}$	> 10.0
	$f_{2-8 \text{ keV}}^c$	2.3	2.4
	$f_{2-5 \text{ keV}}^c$	1.5	1.7
	$f_{5-8 \text{ keV}}^c$	0.8	0.7
	$f_{\text{thcomp+bodyrad}} / f_{2-8 \text{ keV}}$	61%	33%

Notes. The errors are at 90% confidence level. The edges reported on the top section of the table are referred only to the NICER energy spectrum. ^(a) Radii are estimated assuming the distance to the source of 7.6 kpc (see Sect. 3.5). ^(b) The optical depth τ_{thcomp} comes from Eq. (14) in Zdziarski et al. (2020). ^(c) The unabsorbed flux is measured in units $10^{-8} \text{ erg cm}^{-2} \text{ s}^{-1}$.

fits we obtained are based on the following spectral models for Obs. 1 and Obs. 2:

$$\text{tbabs} * (\text{diskbb} + \text{expabs} * \text{powerlaw} + \text{thcomp} * \text{bodyrad}),$$

$$\text{tbabs} * (\text{diskbb} + \text{thcomp} * \text{bodyrad}),$$

respectively.

We included the normalizing cross-calibration multiplicative factors for the NICER, IBIS/ISGRI and *NuSTAR* FPMA (frozen at unity) and FPMB telescopes. A *tbabs* (Wilms et al. 2000) multiplicative model component was used to take into account the low-energy absorption due to the interstellar medium. We used the abundances and cross-section tables according to Wilms et al. (2000) and Verner et al. (1996) (*wilm*, *vern* in *XSPEC*). We modeled the GX 5–1 energy spectra of both the observations with a multi-color disk and a harder boundary/spreading layer (BL, or SL) emission (e.g., Popham & Sunyaev 2001; Revnivtsev et al. 2013). In both observations, the convolution model component *thcomp* (Zdziarski et al. 2020) is used to represent Comptonized emission from the BL/SL (e.g. Di Salvo et al. 2002; Farinelli et al. 2009) modeled with the *bodyrad* component. The f parameter of *thcomp* represents the fraction of Comptonized seed photons. In Obs. 1, this parameter is > 0.63 with the best-fit value equal to 0.99. Effectively, all photons from the BL/SL *bodyrad* are Comptonized. In Obs. 2, only a fraction between 2.7% and 11.2% (best-fit 3.2%) of seed photons are Comptonized. It is worth noting that a higher Comptonization fraction corresponds




Fig. 5. *NuSTAR* CCD and HID of GX 5–1 contemporary to *IXPE* observation. During *IXPE* Obs. 1 the source was in the HB, while during Obs. 2 it was in the NB-FB. Black, red and purple colors refer to the observing days March 21, April 13 and 14, 2023, respectively.




Fig. 6. Deconvolved *NICER* (1.5–10 keV), *NuSTAR* (3.5–45 keV for Obs. 1 and 3.5–35 keV for Obs. 2) and *IBIS/ISGRI* (28–50 keV) spectra simultaneous with both *IXPE* observations as obtained from the best-fit model reported in Table 4 and residuals between data and model in units of σ .

to a higher PD observed from the source. During Obs. 1, when the source is on the HB, the energy spectrum is harder, and a high-energy excess was seen (it was absent in Obs. 2). This excess is modeled with an additional hard tail represented by a power-law component with a low-energy exponential roll-off. The e-folding energy for the absorption of the exponential roll-off is set equal to the $\text{bbodyrad } kT_{\text{bb}}$ energy of the SL/BL. This parameter link comes from the assumption that the power-law emission originates from the seed distribution of the SL/BL blackbody. The BL and the inner disk are hotter in Obs. 2 than in Obs. 1, while the BL sphere-equivalent radius and inner disk radius are larger in Obs. 1. In contrast to Obs. 1, the additional power-law component is not needed to model the spectrum of Obs. 2.

The softer disk component dominates up to ~ 3.5 keV in Obs. 1 and up to ~ 5.5 keV in Obs. 2 (see Fig. 6). In the 2–8 keV

band, the energy flux in the Comptonized component (including the contribution of the exponentially absorbed power-law component) accounts for $\sim 61\%$ of the total in Obs. 1. In Obs. 2, the energy flux of the Comptonized component drops to $\sim 33\%$ of the total in the same energy band.

3.3. X-ray spectro-polarimetric analysis

The results of the spectro-polarimetric analysis, including *IXPE*, once the spectral model used to fit the data from the other observatories is frozen, is reported in Table 5. The PD obtained with a *polconst* multiplicative model component applied to the whole spectral model of GX 5–1 in Obs. 1 and Obs. 2 is $3.7\% \pm 0.4\%$ and $1.8\% \pm 0.4\%$, respectively. The PA is around -9° , in both the observations. The polarization contour plot in the 2–8 keV energy band obtained with *xSPEC* (*polconst* model applied to

Table 5. Best-fit parameters of polarization analysis with *xspec*.

DU	Parameters	Obs. 1	Obs. 2
DU1	N_{DU1}	0.796 ± 0.006	0.798 ± 0.004
	gain slope	$0.953^{+0.003}_{-0.003}$	0.961 ± 0.003
	gain offset (keV)	0.13 ± 0.02	0.102 ± 0.015
DU2	N_{DU2}	0.772 ± 0.006	0.769 ± 0.004
	gain slope	0.952 ± 0.003	0.960 ± 0.003
	gain offset (keV)	0.15 ± 0.02	0.131 ± 0.014
DU3	N_{DU3}	0.734 ± 0.005	0.736 ± 0.004
	gain slope	0.976 ± 0.004	0.966 ± 0.003
	gain offset (keV)	0.10 ± 0.02	0.114 ± 0.015
Components	Parameters	Obs. 1	Obs. 2
$\text{polconst}^*(\text{diskbb} + [\text{expabs}^*\text{powerlaw}]^a + \text{thcomp}^*\text{bbodyrad})$			
	PD (%)	3.7 ± 0.4	1.8 ± 0.4
	PA (deg)	-9 ± 3	-9 ± 6
	$\chi^2/\text{d.o.f.}$	1806/1799	1733/1711
$\text{polconst}_1^*\text{diskbb} + \text{polconst}_2^*([\text{expabs}^*\text{powerlaw}]^a + \text{thcomp}^*\text{bbodyrad})$			
polconst_1	PD (%)	2.3 ± 0.9	1.8 ± 0.9
	PA (deg)	21 ± 11	14 ± 15
polconst_2	PD (%)	5.7 ± 1.4	4.3 ± 2.0
	PA (deg)	-16 ± 20	-32 ± 14
	$\chi^2/\text{d.o.f.}$	1792/1806	1726/1718

Notes. The errors are at 90% confidence level. See Table 4 for the corresponding spectral analysis. Polarization is computed in the 2–8 keV energy range. ^(a) $\text{expabs}^*\text{powerlaw}$ component was included only in Obs. 1.

the spectral model) is shown in Fig. 2. These contours are nearly identical to those obtained using *IXPEOBSSIM*.

The spectro-polarimetric analysis with *xspec* allows us to assign a polarization to the different spectral components. The result of such an analysis is reported in Table 5 and Fig. 7. For Obs. 1, we assigned the same polarization for the $\text{expabs}^*\text{powerlaw}$ and $\text{thcomp}^*\text{bbodyrad}$ components, assuming that the power law is just a continuation of the BL/SL component, with the power-law low energy rollover E_{cut} being equal to the temperature of the BL/SL bbodyrad kT_{bb} .

The Comptonization component is well constrained at a 99.9% confidence level only in Obs. 1, while the disk component is not constrained at 99% in either observation (see Fig. 7). Moreover, in both the observations the polarization of each component has similar PAs, suggesting that the geometry responsible for the polarization is similar. Figure 8 shows the fit of the Q and U Stokes parameters as a function of energy with the two polconst components for both the observations. The contribution to the total flux of the polarized component is different (higher in Obs. 1) probably due to a dilution effect by unpolarized radiation connected with a low covering fraction of the Comptonization component (see Table 4). Unfortunately, even if the source was active in the radio band during the observation campaign, it is impossible to compare the direction of the PA with the direction of the jet, because the radio observations reported in this paper do not have the spatial resolution to resolve it. Moreover, literature does not report any information about jet direction.

The variation of the PA of the total emission, reported in Sect. 3.1, obtained with *IXPEOBSSIM* model independent analysis

can be explained by the different PAs of the disk and Comptonized spectral components which are not aligned, nor being 90° apart. Indeed, the PA of the total emission varies from being positive at lower energies (dominated by the diskbb component) to negative values at higher energies dominated by the Comptonization component with a positive PA.

When assessing polarization as a function of time, no significant variations are seen. This implies that polarization is not sensitive to the flux variation present in the light curves (see Fig. 1).




Fig. 7. Polarization contour plots from the spectro-polarimetric analysis in the 2–8 keV energy range of Obs. 1 (top panel) and Obs. 2 (bottom panel). The polconst polarimetric model is applied separately to the diskbb component of both *IXPE* observations and to the $(\text{expabs}^*\text{powerlaw} + \text{thcomp}^*\text{bbodyrad})$ components as a whole. The $\text{expabs}^*\text{powerlaw}$ component (labeled as pl. in the plot legend) is included only in Obs. 1. Contour plots are computed for 4 parameters of interest, taking into account the fact that the polarization of the disk and Comptonization components are correlated in the simultaneous fit.

3.4. Spectral energy distribution

Thanks to the multi energy observation campaign it was possible to produce a broadband (radio–X-ray) spectral energy distribution (SED) of GX 5–1 for both the observations (see Fig. 9). Optical and infrared fluxes have been de-reddened using the hydrogen column density reported in this work ($N_{\text{H}} = 4.93 \times 10^{22} \text{ cm}^{-2}$ and $4.88 \times 10^{22} \text{ cm}^{-2}$ in Obs. 1 and 2, respectively), which was converted into an estimate of the V-band extinction A_V using the relation reported in Foight et al. (2016b), resulting in $A_V = 17.18 \pm 0.77 \text{ mag}$ and $17.00 \pm 0.72 \text{ mag}$ for the two observations, respectively. The different absorption coefficients at optical and near-IR wavelengths have then been evaluated using the relations reported in Cardelli et al. (1989) and Nishiyama et al. (2008), respectively. For the mid-IR, the coefficients reported in Weingartner & Draine (2001) have instead been used.

GX 5–1 is detected at radio and mid-IR wavelengths, with upper limits in the near-IR and optical. As mentioned in Sect. 2.7, this is due to the large value of dust extinction of $A_V \approx 17$. This implies an extinction of $\sim 7 \text{ mag}$ at $1 \mu\text{m}$ (Y -band) and $\sim 3 \text{ mag}$ at $1.6 \mu\text{m}$ (H -band). The radio spectral index of -0.4 ± 0.1 is too steep for the radio emission to arise from a steady, compact jet (for which we expect a spectral index from ~ 0 to $+0.5$). As such, it is likely due to optically thin jet ejections, or a combination of compact jet and optically thin ejections. In the mid-IR, we report the first detection of this source. The de-reddened flux densities are comparable to the near-IR values reported in the literature (Naylor et al. 1991; Jonker et al. 2000; Bandyopadhyay et al. 2003); however, the de-reddened near-IR fluxes depend sensitively on the value of the interstellar extinction. We note that different values of the neutral hydrogen col-




Fig. 8. Fit of the *IXPE* Stokes parameters Q and U as a function of energy with the model comprising two `polconst` components applied to the disk (dotted lines) and to the Comptonization (dashed lines) for Obs. 1 (left column) and Obs. 2 (right column).




Fig. 9. Broad-band radio to X-ray SED of GX 5–1 during the two observations. For Obs. 2, no mid-IR flux is reported as observations were not performed with VISIR at that time (see Sect. 2.6 for details). The flux densities at mid-IR, near-IR and optical frequencies have been dereddened as described in Sect. 3.4. The errors are at 68% confidence level.

umn densities are reported in the literature, ranging from $N_{\text{H}} = 2.54 \times 10^{22} \text{ cm}^{-2}$ to $6.20 \times 10^{22} \text{ cm}^{-2}$ (Christian & Swank 1997; Zeegers et al. 2017; Homan et al. 2018; Clark 2018; Bhulla et al. 2019). Yang et al. (2022) has derived $N_{\text{H}} = (4.52 \pm 0.01) \times 10^{22} \text{ cm}^{-2}$ by measuring the Si K edge due to scattering by dust us-

ing the *Chandra* gratings. This value is in agreement with our measurements because N_{H} is slightly overestimated when fitting the continuum with the absorption model `tbbabs` (Corrales et al. 2016). In this work, we observed GX 5–1 over a significantly larger energy range, and we are confident to have properly constrained the absorption in the interstellar medium. However, if there is a component of the neutral hydrogen column which is intrinsic to the LMXB, this could cause varying measurements and introduce uncertainty into the relation between N_{H} and the extinction A_{V} . Moreover, older works using Anders & Grevesse (1989) solar abundances rather than Wilms et al. (2000) interstellar abundances, could be affected by model systematic.

The mid-IR flux measured with the VLT VISIR is higher than the extrapolation of the ATCA spectral index from radio to mid-IR. The mid-IR emission is therefore unlikely to be due to optically thin synchrotron from discrete jet ejections that were seen in the radio, but it could be optically thin synchrotron emission from the compact jet (from above the jet spectral break; Russell et al. 2013). We find evidence of strong mid-IR variability between the two epochs, with the $9\text{--}11\mu\text{m}$ flux density changing from $< 3.27 \text{ mJy}$ on 2023-03-28, to $9.95 \pm 2.31 \text{ mJy}$ on 2023-03-31. While this variation by a factor of ≥ 3 (≥ 2.5 magnitudes) is quite high for a LMXB accretion disk on these timescales, high amplitude (spanning several magnitudes) infrared variability has been reported from a number of bright, persistent NS-LMXBs, including GX 17+2, Cir X-1, 4U 1705–440 and GX 13+1 (e.g. Glass 1994; Callanan et al. 2002; Bandyopadhyay et al. 2002; Homan et al. 2009; Corbet et al. 2010; Harrison et al. 2011).




Fig. 10. Fit of the infrared extinction A_{K_s} between 5 and 15 kpc from Marshall et al. (2006) with a linear function (the best-fit line and the lines taking into account the associated errors are in blue). The red horizontal lines represent the estimate with the corresponding upper and lower limits of A_{K_s} from Eq. (3). The distance to the source is thus $d = 7.6 \pm 1.1$ kpc.

This variability has been generally interpreted as indicative of highly variable synchrotron emission from a compact jet. Variable near-IR polarization has also been reported from the bright, persistent NS-LMXBs Sco X-1 and Cyg X-2, as well as the NS-LMXB transient SAX J1808.4–3658 (Shahbaz et al. 2008; Russell & Fender 2008; Baglio et al. 2020). The de-reddened mid-IR 4.7–8.7 μm spectral index on 2023-03-31 is -2.3 to -0.4 (adopting $A_V = 17.18 \pm 0.77$), which is consistent with optically thin synchrotron emission from relativistic particles, or a steeper particle distribution with a thermal (Maxwellian) component, as has been seen at infrared wavelengths from jets in some black hole LMXBs (e.g., Russell et al. 2010; Shahbaz et al. 2013). Thus, this emission could originate from a compact jet which peaks in the mid- or far-IR, although follow-up observations characterizing the IR spectrum and variability would be beneficial to confirm the nature of the IR emission. If the compact jet is present, its spectrum from radio to mid-IR must be inverted, with index > 0.33 (and fainter than the observed radio emission). The radio to IR spectrum is similar in some ways to the NS-LMXBs 4U 1728–34 and 4U 0614+091 (Migliari et al. 2010; Díaz Trigo et al. 2017).

3.5. Measurement of the distance to the source

In order to derive some spectral model parameters (namely $R_{\text{in}} \sqrt{\cos \theta}$ of `diskbb` and R_{bb} of `bbodyrad`), an estimate of the distance to the source is needed. Such an estimate can be obtained from the equivalent hydrogen column density N_{H} which we get from our analysis. We find that $N_{\text{H}} = (4.93^{+0.12}_{-0.06}) \times 10^{22} \text{ cm}^{-2}$ and $(4.88^{+0.03}_{-0.04}) \times 10^{22} \text{ cm}^{-2}$ for the first and the second observations, respectively. Because the two values are the same at the 90% confidence levels, we use the average value and its largest uncertainty range, that is $(4.91^{+0.14}_{-0.07}) \times 10^{22} \text{ cm}^{-2}$, in the following discussion. To estimate the distance to the source we adopt the approach proposed by Gambino et al. (2016). We use the model of the infrared Galactic interstellar extinction discussed by Marshall et al. (2006). Because the Galactic coordinates of GX 5–1 are $l = 5^\circ 08'$ and $b = -1^\circ 02'$ we adopt the map

that relates the infrared extinction A_{K_s} with the source distance d valid for $l = 5^\circ$ and $b = -1^\circ$ (see black dots with the error bars in Fig. 10).

Because the visual extinction A_V is related to N_{H} as (Foight et al. 2016a)

$$N_{\text{H}} = (2.87 \pm 0.12) \times 10^{21} A_V, \quad (1)$$

and the relation between A_V and the extinction in the K_s band is (Nishiyama et al. 2008)

$$A_{K_s} = (0.062 \pm 0.005) A_V, \quad (2)$$

we obtain

$$A_{K_s} = \frac{(0.062 \pm 0.005)}{(2.87 \pm 0.12) \times 10^{21}} N_{\text{H}} \text{ mag} = 1.06 \pm 0.10 \text{ mag}. \quad (3)$$

We fit the A_{K_s} values between 5 and 15 kpc with a linear function (the best-fit line and the lines taking into account the associated errors are in blue color in Fig. 10). We infer that the distance to the source is $d = 7.6 \pm 1.1$ kpc. This distance is in agreement with the previously reported values (Penninx 1989; Smith et al. 2006).

4. Discussion

As shown in Sect. 3, the two multiwavelength observations of GX 5–1 allowed us to catch the source when it was covering the complete Z-track on its CCD/HID (see Fig. 5). In particular, during the first observation, GX 5–1 was on the HB of the track, while in the second it moved across to the NB and FB. Very interestingly, we found the same behavior in the peculiar transient XTE J1701–462 (Cocchi et al. 2023), in which the PD was correlated with the source position on the Z-track, being higher in the HB and decreasing by a factor about two in the NB. Long et al. (2022) hypothesized the same behavior also for Sco X-1. There are at least three regions which may potentially contribute to the polarization: the BL/SL, the accretion disk, and the reflection of the BL/SL photons from the disk atmosphere or a wind.

The spectro-polarimetric analysis of the *IXPE* data (Table 5) shows that the disk polarization is about 2% in both observations, which is compatible with the classical results of a high optical depth scattering atmosphere at an inclination of 60° (Chandrasekhar 1960). The higher PD value of the hard component on the other hand, cannot be explained by repeated Compton scattering in high optical depth environment, neither for a boundary layer coplanar with the disk (which otherwise would resemble a Chandrasekhar-like slab) nor for a spreading layer around the neutron star, for which a maximum PD of $\lesssim 2\%$ is expected.

Disk reflection is probably the most natural way to explain a PD of order of 4%–5%, as shown by Lapidus & Sunyaev (1985). It is important to note that we do not find a strong reflection signature in the spectral analysis, and it deserves mentioning that the reflection contribution to the spectrum may be low and sometimes may be embedded in the continuum but, nevertheless, make a large contribution to the net polarization signal (Schnittman & Krolik 2009). This may be particularly true if the primary spectrum is not a hard power law, but a blackbody-like spectrum with a rollover below 30 keV. Note that a similar argument has been used also for Cyg X-2 (Farinelli et al. 2023) and GX 9+9 (Ursini et al. 2023) to explain the high PD attributed to the Comptonization spectrum in a two-fold spectro-polarimetric approach.

Table 6. Best-fit parameters with reflection included in the energy spectrum model of GX 5–1 from NICER, *NuSTAR* and *INTEGRAL* data simultaneous to *IXPE* observation.

Components	Parameters	Obs1	Obs2
edge	E (keV)	1.820 ± 0.013	1.818 ± 0.013
	τ	0.15 ± 0.02	0.15 ± 0.02
edge	E (keV)	1.95197 (fixed)	
	τ	0.05 ± 0.02	0.05 ± 0.02
edge	E (keV)	2.28003 (fixed)	
	τ	0.034 ± 0.014	0.027 ± 0.013
edge	E (keV)	2.44444 (fixed)	
	τ	0.049 ± 0.013	0.038 ± 0.013
edge	E (keV)	3.16139 (fixed)	
	τ	0.021 ± 0.007	0.011 ± 0.007
tbabs	N_{H} (10^{22} cm^{-2})	4.80 ± 0.05	4.87 ± 0.05
diskbb	kT (keV)	1.21 ± 0.05	1.20 ± 0.03
	$R_{\text{in}} \sqrt{\cos \theta}$ (km) ^a	16 ± 1	$19^{+1}_{-0.9}$
comptb	kT_0 (keV)	1.93 ± 0.13	1.71 ± 0.04
	α	$2.9^{+1.7}_{-0.8}$	< 1.16
	kT_e (keV)	$3.6^{+1.9}_{-0.4}$	$3.24^{+0.38}_{-0.10}$
	$\log A$	8 (fixed)	$-1.54^{+0.54}_{-0.06}$
	N_{comptb}	0.148 ± 0.003	0.123 ± 0.006
expabs	E_{cut} (keV)	[= kT_0]	–
powerlaw	Γ	2.4 (fixed)	–
	norm	0.17 ± 0.05	–
rdblur	betor	–2.5 (fixed)	–
	R_{in} (R_g)	7 (fixed)	–
	R_{out} (R_g)	1000 (fixed)	–
	incl. (deg)	60 (fixed)	–
rfxconv	rel-refl	0.3 (fixed)	–
	$\log \xi$	3.8 (fixed)	–
	$\chi^2/\text{d.o.f.}$	268/295	247/260

Notes. The errors are at 90% confidence level. ^(a) Radii are estimated assuming the distance to the source of 7.6 kpc (see Sect. 3.5).

The PD of the reflected radiation is not easy to predict because it depends on geometrical and physical factors (e.g., the disk ionization parameter) however, it is not likely to exceed ~20% (Matt 1993; Poutanen et al. 1996; Schnittman & Krolik 2009). We tested the hypothesis as to whether there is a reflection component in the GX 5–1 energy spectrum by making some assumptions: (1) highly ionized disk due to the absence of a broad emission line in the Fe-K region of the spectrum, (2) inclination $i = 60^\circ$ since GX 5–1 is a Cyg-like Z-source (Homan et al. 2018), (3) keep the reflection amplitude $f = \Omega/2\pi$ at a typical value for the NS-LMXBs (namely 30%, see, e.g., Di Salvo et al. 2015; Matranga et al. 2017). We applied the following spectral models:

Obs. 1: `constant*tbabs(diskbb + expabs*powerlaw + rdblur*rfxconv*comptb + comptb)`,

Obs. 2: `constant*tbabs(diskbb + rdblur*rfxconv*comptb + comptb)`.

The `comptb` (Farinelli et al. 2008) model component was included in place of the `thcomp` to prevent a double convolution of the BL/SL blackbody. Details on `rfxconv` and `rdblur` model components can be found in Kolehmainen et al. (2011) and Fabian et al. (1989), respectively. The sum `rdblur*rfxconv*comptb + comptb` accounts for the reflected radiation (the first term) plus the incident radiation (the second term).

This modeling of the reflection component contribute ~22% of the flux in Obs. 1 and ~12% in Obs. 2. Such a contribution to the total emission can easily account for the polarization detected (see Table 6 for details of the fit parameters of the energy spectrum). The fraction of Comptonized photons in Obs. 2 is significantly smaller with respect to Obs. 1, as obtained also in Sect. 3 by applying only the Comptonization model `thcomp`. The parameter $\log A = -1.54^{+0.54}_{-0.06}$ in `comptb` corresponds to a $2.8^{+6.3}_{-0.4}\%$

of photons upscattered in energy (consistent with $3.2^{+8.0}_{-0.5}\%$ obtained when reflection is not taken into account as in Sect. 3). This confirms the presence of a blackbody component observed through an almost vanished Comptonization medium in Obs. 2, even if reflection is included. However, the fraction of Comptonized photons, albeit small (~3%), is still sufficient to manifest its presence at high energy. Indeed, if we would neglect this small fraction of Comptonized photons by substituting `comptb` with a blackbody component, such as `bbodyrad`, we obtain an unacceptable fit result ($\chi^2_\nu = 3.6$, with a significant excess at high energy due to this small fraction of Comptonized photons).

Another possible mechanism for producing polarization is related to scattering in the wind above the accretion disk: as was shown in Sunyaev & Titarchuk (1985), the emission scattered once in a plane (e.g. equatorial wind) can be polarized up to 27% for an inclination $i = 60^\circ$. This polarization degree is only weakly dependent on the opening angle of the wind. Assuming that 20% of the source emission is scattered, we can obtain the observed PD. Recently, a similar model was shown to explain well the presence of a constant polarized component in the X-ray pulsars RX J0440.9+4431 / LS V +44 17 (Doroshenko et al. 2023). Although it is well known that strong winds can be indeed present in the soft states of X-ray binaries (e.g., Neilsen & Lee 2009; Ponti et al. 2012, 2014), it is worth noting that we do not see wind absorption features in the energy spectrum of GX 5–1, implying that, if present, the wind should be completely ionized.

Both *IXPE* observations show similar behavior of the PD values: they are smaller in the 2–4 keV band and increase slightly with energy. The reduction of the PD at lower energies can be explained by the energy dependence of the disk emission: in the low energy band we have a marginally polarized disk emission dominating the spectrum, while at higher energies, the emission of the SL and/or the scattered/reflected component is more visible. Another feature of the observed emission that needs to be addressed is the variation of the PA with energy. In the spectropolarimetric analysis the disk and the Comptonization components have non-orthogonal PAs, thus the variation of the polarization plane of the total emission with energy can be interpreted as the energy-dependent contribution of these two components to the total emission. The same applies if, for instance, the SL emission is scattered in the wind and the disk emission is polarized with a different and non-orthogonal angle. The PA of the combined emission will be energy dependent. It is well known that the disk emission exhibits the rotation of the polarization plane with energy (Connors et al. 1980; Dovčiak et al. 2008; Loktev et al. 2022). Simulations of the SL emission also show a change of the PA with energy but predict a PD at most 1.5% (Bobrikova A., in prep.). Thus, a single component explanation of the PA variation cannot be considered compatible with the high measured PD. Further modeling will be needed to satisfactory address the explanations presented above.

5. Summary

IXPE observed the NS-LMXB GX 5–1 twice in the period March–April 2023. Contemporary observations in the X-ray energy band were put in place with NICER, *NuSTAR*, and *INTEGRAL*. Multi-wavelength coverage was ensured by ATCA in the radio, VLT VISIR in mid-IR, REM in optical, and NIR and LCO in optical.

During the observations GX 5–1 moved across the entire Z-track. *NuSTAR* disentangled clearly the NB with respect to the FB thanks to its extended energy band. The presence of a hard tail, reported in previous analyses, was clearly detected

in Obs. 1, but not in Obs. 2, when the source had a softer energy spectrum. The X-ray PD was $\sim 4\%$ during Obs. 1 when the source was in the HB and $\sim 2\%$ during Obs. 2 with the source in the NB-FB. This result is in agreement with findings from the other Z-sources observed by *IXPE* (namely Cyg X-2 and XTE J1701–462).

The source manifested an unexpected variation of the PA as a function of energy by $\sim 20^\circ$. The magnitude of the variation combined with the magnitude of the PD require further modeling. However, it is likely related to the different PAs of the disk and Comptonization components which are non-orthogonal and, moreover, have the emission peaks at different energies.

In the radio band, the source was detected, but only upper-limits to the polarization were obtained ($\sim 6\%$ at 5.5 GHz and 9 GHz in Obs. 1 and 12.5% at 5.5 GHz and 20% at 9 GHz in Obs. 2). The mid-IR counterpart was detected in M and the J8.9 bands. This emission could originate from a compact jet which peaks in the mid- or far-IR. Follow-up observations characterizing the IR spectrum and its variability would be beneficial. Due to the very high extinction toward the source and to the crowded field, the source was not detected in any of the optical and NIR images acquired.

Acknowledgements. The Imaging X-ray Polarimetry Explorer (*IXPE*) is a joint US and Italian mission. The US contribution is supported by the National Aeronautics and Space Administration (NASA) and led and managed by its Marshall Space Flight Center (MSFC), with industry partner Ball Aerospace (contract NNM15AA18C). The Italian contribution is supported by the Italian Space Agency (Agenzia Spaziale Italiana, ASI) through contract ASI-OHBI-2017-12-I.O, agreements ASI-INAF-2017-12-H0 and ASI-INFN-2017.13-H0, and its Space Science Data Center (SSDC) with agreements ASI-INAF-2022-14-HH.0 and ASI-INFN 2021-43-HH.0, and by the Istituto Nazionale di Astrofisica (INAF) and the Istituto Nazionale di Fisica Nucleare (INFN) in Italy. This research used data products provided by the *IXPE* Team (MSFC, SSDC, INAF, and INFN) and distributed with additional software tools by the High-Energy Astrophysics Science Archive Research Center (HEASARC), at NASA Goddard Space Flight Center (GSFC). ATCA is part of the Australia Telescope National Facility (<https://ror.org/05qajvd42>) which is funded by the Australian Government for operation as a National Facility managed by CSIRO. We acknowledge the Gomeri people as the Traditional Owners of the ATCA Observatory site. Partly based on observations with INTEGRAL, an ESA project with instruments and science data centre funded by ESA member states (especially the PI countries: Denmark, France, Germany, Italy, Switzerland, Spain), and with the participation of the Russian Federation and the USA. Data have been reduced using the Multi-Messenger Online Data Analysis platform provided by the ISDC (University of Geneva), EPFL, APC, and KAU. J.P. acknowledges support from the Academy of Finland grant 333112. A.B. is supported by the Finnish Cultural Foundation grant 00220175. T.D.R. thanks Jamie Stevens and James Miller-Jones for the helpful discussions regarding the radio observations. J.S. and M.D. acknowledge the support from the GACR project 21-06825X.

References

- Anders, E. & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197
- Arnaud, K. A. 1996, in *ASP Conf. Ser.*, Vol. 101, *Astronomical Data Analysis Software and Systems V*, ed. G. H. Jacoby & J. Barnes (San Francisco: ASP), 17–20
- Asai, K., Dotani, T., Mitsuda, K., et al. 1994, *PASJ*, 46, 479
- Baglio, M. C., Russell, D. M., Crespi, S., et al. 2020, *ApJ*, 905, 87
- Baldini, L., Bucciantini, N., Lalla, N. D., et al. 2022, *SoftwareX*, 19, 101194
- Bandyopadhyay, R. M., Charles, P. A., Shahbaz, T., & Wagner, R. M. 2002, *ApJ*, 570, 793
- Bandyopadhyay, R. M., Shahbaz, T., & Charles, P. A. 2003, *MNRAS*, 340, L13
- Berendsen, S. G. H., Fender, R., Kuulkers, E., Heise, J., & van der Klis, M. 2000, *MNRAS*, 318, 599
- Bhulla, Y., Misra, R., Yadav, J. S., & Jaaffrey, S. N. A. 2019, *Research in Astronomy and Astrophysics*, 19, 114
- Briggs, D. S. 1995, PhD thesis, New Mexico Institute of Mining and Technology
- Callanan, P. J., Curran, P., Filippenko, A. V., et al. 2002, *ApJ*, 574, L143
- Capitanio, F., Fabiani, S., Gnarini, A., et al. 2023, *ApJ*, 943, 129
- Cardelli, J. A., Clayton, G. C., & Mathis, J. S. 1989, *ApJ*, 345, 245
- CASA Team, Bean, B., Bhatnagar, S., et al. 2022, *PASP*, 134, 114501
- Chandrasekhar, S. 1960, *Radiative Transfer* (New York: Dover)
- Chatterjee, R., Agrawal, V. K., Jayasurya, K. M., & Katoch, T. 2023, *MNRAS*, 521, L74
- Christian, D. J. & Swank, J. H. 1997, *ApJS*, 109, 177
- Clark, G. W. 2018, *ApJ*, 852, 121
- Cocchi, M., Gnarini, A., Fabiani, S., et al. 2023, *A&A*, 674, L10
- Connors, P. A., Piran, T., & Stark, R. F. 1980, *ApJ*, 235, 224
- Corbet, R. H. D., Pearlman, A. B., Buxton, M., & Levine, A. M. 2010, *ApJ*, 719, 979
- Corrales, L. R., García, J., Wilms, J., & Baganoff, F. 2016, *MNRAS*, 458, 1345
- Courvoisier, T. J. L., Walter, R., Beckmann, V., et al. 2003, *A&A*, 411, L53
- Covino, S., Stefanon, M., Sciuto, G., et al. 2004, in *Proc. SPIE*, Vol. 5492, *Ground-based Instrumentation for Astronomy*, ed. A. F. M. Moorwood & M. Iye, 1613–1622
- Di Marco, A., Costa, E., Muleri, F., et al. 2022, *AJ*, 163, 170
- Di Marco, A., Soffitta, P., Costa, E., et al. 2023, *AJ*, 165, 143
- Di Salvo, T., Farinelli, R., Burderi, L., et al. 2002, *A&A*, 386, 535
- Di Salvo, T., Iaria, R., Matranga, M., et al. 2015, *MNRAS*, 449, 2794
- Di Salvo, T. & Stella, L. 2002, *arXiv e-prints*, astro
- Díaz Trigo, M., Migliari, S., Miller-Jones, J. C. A., et al. 2017, *A&A*, 600, A8
- Doroshenko, V., Poutanen, J., Heyl, J., et al. 2023, *A&A*, 677, A57
- Dovčiak, M., Muleri, F., Goosmann, R. W., Karas, V., & Matt, G. 2008, *MNRAS*, 391, 32
- Fabian, A. C., Rees, M. J., Stella, L., & White, N. E. 1989, *MNRAS*, 238, 729
- Farinelli, R., Fabiani, S., Poutanen, J., et al. 2023, *MNRAS*, 519, 3681
- Farinelli, R., Paizis, A., Landi, R., & Titarchuk, L. 2009, *A&A*, 498, 509
- Farinelli, R., Titarchuk, L., Paizis, A., & Frontera, F. 2008, *ApJ*, 680, 602
- Fender, R. P. & Hendry, M. A. 2000, *MNRAS*, 317, 1
- Foight, D. R., Güver, T., Özel, F., & Slane, P. O. 2016a, *ApJ*, 826, 66
- Foight, D. R., Güver, T., Özel, F., & Slane, P. O. 2016b, *ApJ*, 826, 66
- Gambino, A. F., Iaria, R., Di Salvo, T., et al. 2016, *A&A*, 589, A34
- Gendreau, K. C., Arzoumanian, Z., Adkins, P. W., et al. 2016, in *Proc. SPIE*, Vol. 9905, *Space Telescopes and Instrumentation 2016: Ultraviolet to Gamma Ray*, ed. J.-W. A. den Herder, T. Takahashi, & M. Bautz, 99051H
- Gilfanov, M., Churazov, E., Sunyaev, R., et al. 1993, *ApJ*, 418, 844
- Glass, I. S. 1994, *MNRAS*, 268, 742
- Harrison, F. A., Craig, W. W., Christensen, F. E., et al. 2013, *ApJ*, 770, 103
- Harrison, T. E., McNamara, B. J., Bornak, J., et al. 2011, *ApJ*, 736, 54
- Hasinger, G. & van der Klis, M. 1989, *A&A*, 225, 79
- Homan, J., Kaplan, D. L., van den Berg, M., & Young, A. J. 2009, *ApJ*, 692, 73
- Homan, J., Steiner, J. F., Lin, D., et al. 2018, *ApJ*, 853, 157
- Jonker, P. G., Fender, R. P., Hambly, N. C., & van der Klis, M. 2000, *MNRAS*, 315, L57
- Jonker, P. G., van der Klis, M., Homan, J., et al. 2002, *MNRAS*, 333, 665
- Kolehmainen, M., Done, C., & Díaz Trigo, M. 2011, *MNRAS*, 416, 311
- Kuulkers, E., van der Klis, M., Oosterbroek, T., et al. 1994, *A&A*, 289, 795
- Lagage, P. O., Pel, J. W., Authier, M., et al. 2004, *The Messenger*, 117, 12
- Lapidus, I. I. & Sunyaev, R. A. 1985, *MNRAS*, 217, 291
- Lebrun, F., Leray, J. P., Lavocat, P., et al. 2003, *A&A*, 411, L141
- Liu, Q. Z., van Paradijs, J., & van den Heuvel, E. P. J. 2007, *A&A*, 469, 807
- Loktev, V., Veledina, A., & Poutanen, J. 2022, *A&A*, 660, A25
- Long, X., Feng, H., Li, H., et al. 2022, *ApJ*, 924, L13
- Marshall, D. J., Robin, A. C., Reylé, C., Schultheis, M., & Picaud, S. 2006, *A&A*, 453, 635
- Matranga, M., Papitto, A., Di Salvo, T., et al. 2017, *A&A*, 603, A39
- Matt, G. 1993, *MNRAS*, 260, 663
- Migliari, S. & Fender, R. P. 2006, *MNRAS*, 366, 79
- Migliari, S., Tomsick, J. A., Miller-Jones, J. C. A., et al. 2010, *ApJ*, 710, 117
- Mitsuda, K., Inoue, H., Koyama, K., et al. 1984, *PASJ*, 36, 741
- Muleri, F. 2022, in *Handbook of X-ray and Gamma-ray Astrophysics*, ed. C. Bambi & A. Santangelo (Singapore: Springer)
- Nasa Heasarc. 2014, *HEASoft: Unified Release of FTOOLS and XANADU*, Astrophysics Source Code Library, record ascl:1408.004
- Naylor, T., Charles, P. A., & Longmore, A. J. 1991, *MNRAS*, 252, 203
- Neilsen, J. & Lee, J. C. 2009, *Nature*, 458, 481
- Nishiyama, S., Nagata, T., Tamura, M., et al. 2008, *ApJ*, 680, 1174
- Paizis, A., Ebisawa, K., Tikkanen, T., et al. 2005, *A&A*, 443, 599
- Paizis, A., Farinelli, R., Titarchuk, L., et al. 2006, *A&A*, 459, 187
- Penninx, W. 1989, in *ESA Special Publication*, Vol. 1, *Two Topics in X-Ray Astronomy*, Volume 1: X Ray Binaries. Volume 2: AGN and the X Ray Background, ed. J. Hunt & B. Battrick, 185
- Ponti, G., Fender, R. P., Begelman, M. C., et al. 2012, *MNRAS*, 422, L11
- Ponti, G., Muñoz-Darias, T., & Fender, R. P. 2014, *MNRAS*, 444, 1829
- Popham, R. & Sunyaev, R. 2001, *ApJ*, 547, 355
- Poutanen, J., Nagendra, K. N., & Svensson, R. 1996, *MNRAS*, 283, 892
- Revnivtsev, M. G., Suleimanov, V. F., & Poutanen, J. 2013, *MNRAS*, 434, 2355
- Russell, D. M. & Fender, R. P. 2008, *MNRAS*, 387, 713
- Russell, D. M., Maitra, D., Dunn, R. J. H., & Markoff, S. 2010, *MNRAS*, 405, 1759
- Russell, D. M., Markoff, S., Casella, P., et al. 2013, *MNRAS*, 429, 815

- Schnittman, J. D. & Krolik, J. H. 2009, *ApJ*, 701, 1175
- Shahbaz, T., Fender, R. P., Watson, C. A., & O'Brien, K. 2008, *ApJ*, 672, 510
- Shahbaz, T., Russell, D. M., Zurita, C., et al. 2013, *MNRAS*, 434, 2696
- Smith, R. K., Dame, T. M., Costantini, E., & Predehl, P. 2006, *ApJ*, 648, 452
- Soffitta, P., Baldini, L., Bellazzini, R., et al. 2021, *AJ*, 162, 208
- Strohmayer, T. E. & Kallman, T. R. 2013, *ApJ*, 773, 103
- Sunyaev, R., Churazov, E., Gilfanov, M., et al. 1991, *A&A*, 247, L29
- Sunyaev, R. A. & Titarchuk, L. G. 1985, *A&A*, 143, 374
- Titarchuk, L. 1994, *ApJ*, 434, 570
- Ubertini, P., Lebrun, F., Di Cocco, G., et al. 2003, *A&A*, 411, L131
- Ursini, F., Farinelli, R., Gnarini, A., et al. 2023, *A&A*, 676, A20
- van der Klis, M. 1995, in *X-ray Binaries*, ed. W. H. G. Lewin, v. E. P. J., & J. van Paradijs, Cambridge Astrophysics Series, No. 26 (Cambridge: Cambridge University Press), 252–307
- Verner, D. A., Ferland, G. J., Korista, K. T., & Yakovlev, D. G. 1996, *ApJ*, 465, 487
- Weingartner, J. C. & Draine, B. T. 2001, *ApJ*, 563, 842
- Weisskopf, M. C., Elsner, R. F., & O'Dell, S. L. 2010, in *Proc. SPIE*, Vol. 7732, *Space Telescopes and Instrumentation 2010: Ultraviolet to Gamma Ray*, ed. M. Arnaud, S. S. Murray, & T. Takahashi, 77320E
- Weisskopf, M. C., Soffitta, P., Baldini, L., et al. 2022, *JATIS*, 8, 026002
- Weisskopf, M. C., Soffitta, P., Ramsey, B. D., et al. 2023, *Nature Astronomy*, 7, 635
- White, N. E., Peacock, A., Hasinger, G., et al. 1986, *MNRAS*, 218, 129
- Wilms, J., Allen, A., & McCray, R. 2000, *ApJ*, 542, 914
- Winkler, C., Courvoisier, T. J. L., Di Cocco, G., et al. 2003, *A&A*, 411, L1
- Yang, J., Schulz, N. S., Rogantini, D., et al. 2022, *AJ*, 164, 182
- Zdziarski, A. A., Szanecki, M., Poutanen, J., Gierliński, M., & Biernacki, P. 2020, *MNRAS*, 492, 5234
- Zeegers, S. T., Costantini, E., de Vries, C. P., et al. 2017, *A&A*, 599, A117
- Zerbi, R. M., Chincarini, G., Ghisellini, G., et al. 2001, *Astronomische Nachrichten*, 322, 275
- ¹ INAF Istituto di Astrofisica e Planetologia Spaziali, Via del Fosso del Cavaliere 100, 00133 Roma, Italy e-mail: sergio.fabiani@inaf.it
- ² Università degli Studi di Palermo, Dipartimento di Fisica e Chimica, Emilio Segrè, via Archirafi 36 I-90123 Palermo, Italy
- ³ Department of Physics and Astronomy, FI-20014 University of Turku, Finland
- ⁴ Dipartimento di Matematica e Fisica, Università degli Studi Roma Tre, via della Vasca Navale 84, 00146 Roma, Italy
- ⁵ INAF Osservatorio di Astrofisica e Scienza dello Spazio di Bologna, Via P. Gobetti 101, I-40129 Bologna, Italy
- ⁶ Center for Astrophysics, Harvard & Smithsonian, 60 Garden St, Cambridge, MA 02138, USA
- ⁷ Astronomical Institute of the Czech Academy of Sciences, Boční II 1401/1, 14100 Praha 4, Czech Republic
- ⁸ Center for Astro, Particle and Planetary Physics, New York University Abu Dhabi, PO Box 129188, Abu Dhabi, UAE
- ⁹ INAF Osservatorio Astronomico di Brera, via E. Bianchi 46, 23807 Merate (LC), Italy
- ¹⁰ Department of Physics, University of Warwick, Coventry CV4 7AL, UK
- ¹¹ Astrophysics, Department of Physics, University of Oxford, Keble Road, Oxford OX1 3RH
- ¹² INAF Istituto di Astrofisica Spaziale e Fisica Cosmica, Via U. La Malfa 153, I-90146 Palermo, Italy
- ¹³ Department of Astronomy, University of Geneva, Ch. d'Ecogia 16, CH-1290 Versoix, Geneva, Switzerland
- ¹⁴ Faulkes Telescope Project, School of Physics and Astronomy, Cardiff University, The Parade, Cardiff CF24 3AA, Wales, UK
- ¹⁵ Astrophysics Research Institute, Liverpool John Moores University, 146 Brownlow Hill, Liverpool L3 5RF, UK
- ¹⁶ INAF Osservatorio Astronomico di Cagliari, Via della Scienza 5, 09047 Selargius (CA), Italy
- ¹⁷ Guangxi Key Laboratory for Relativistic Astrophysics, School of Physical Science and Technology, Guangxi University, Nanning 530004, China
- ¹⁸ NASA Marshall Space Flight Center, Huntsville, AL 35812, USA
- ¹⁹ Dipartimento di Fisica, Università degli Studi di Cagliari, SP Monserrato-Sestu, KM 0.7, Monserrato, I-09042 Italy
- ²⁰ INFN, Sezione di Cagliari, Cittadella Universitaria, I-09042 Monserrato, CA, Italy
- ²¹ INAF Osservatorio Astronomico di Roma, Via Frascati 33, 00040 Monte Porzio Catone (RM), Italy
- ²² Instituto de Astrofísica de Andalucía – CSIC, Glorieta de la Astronomía s/n, 18008 Granada, Spain
- ²³ Space Science Data Center, Agenzia Spaziale Italiana, Via del Politecnico snc, 00133 Roma, Italy
- ²⁴ Istituto Nazionale di Fisica Nucleare, Sezione di Pisa, Largo B. Pontecorvo 3, 56127 Pisa, Italy
- ²⁵ Dipartimento di Fisica, Università di Pisa, Largo B. Pontecorvo 3, 56127 Pisa, Italy
- ²⁶ Istituto Nazionale di Fisica Nucleare, Sezione di Torino, Via Pietro Giuria 1, 10125 Torino, Italy
- ²⁷ Dipartimento di Fisica, Università degli Studi di Torino, Via Pietro Giuria 1, 10125 Torino, Italy
- ²⁸ INAF Osservatorio Astrofisico di Arcetri, Largo Enrico Fermi 5, 50125 Firenze, Italy
- ²⁹ Dipartimento di Fisica e Astronomia, Università degli Studi di Firenze, Via Sansone 1, 50019 Sesto Fiorentino (FI), Italy
- ³⁰ Istituto Nazionale di Fisica Nucleare, Sezione di Firenze, Via Sansone 1, 50019 Sesto Fiorentino (FI), Italy
- ³¹ Agenzia Spaziale Italiana, Via del Politecnico snc, 00133 Roma, Italy
- ³² Science and Technology Institute, Universities Space Research Association, Huntsville, AL 35805, USA
- ³³ Istituto Nazionale di Fisica Nucleare, Sezione di Roma “Tor Vergata”, Via della Ricerca Scientifica 1, 00133 Roma, Italy
- ³⁴ Department of Physics and Kavli Institute for Particle Astrophysics and Cosmology, Stanford University, Stanford, California 94305, USA
- ³⁵ Institut für Astronomie und Astrophysik, Universität Tübingen, Sand 1, D-72076 Tübingen, Germany
- ³⁶ RIKEN Cluster for Pioneering Research, 2-1 Hirosawa, Wako, Saitama 351-0198, Japan
- ³⁷ California Institute of Technology, Pasadena, CA 91125, USA
- ³⁸ Yamagata University, 1-4-12 Kojirakawa-machi, Yamagata-shi 990-8560, Japan
- ³⁹ Osaka University, 1-1 Yamadaoka, Suita, Osaka 565-0871, Japan
- ⁴⁰ University of British Columbia, Vancouver, BC V6T 1Z4, Canada
- ⁴¹ International Center for Hadron Astrophysics, Chiba University, Chiba 263-8522, Japan
- ⁴² Institute for Astrophysical Research, Boston University, 725 Commonwealth Avenue, Boston, MA 02215, USA
- ⁴³ Department of Astrophysics, St. Petersburg State University, Universitetskij pr. 28, Petrodvoretz, 198504 St. Petersburg, Russia
- ⁴⁴ Department of Physics and Astronomy and Space Science Center, University of New Hampshire, Durham, NH 03824, USA
- ⁴⁵ Physics Department and McDonnell Center for the Space Sciences, Washington University in St. Louis, St. Louis, MO 63130, USA
- ⁴⁶ Finnish Centre for Astronomy with ESO, 20014 University of Turku, Finland
- ⁴⁷ Istituto Nazionale di Fisica Nucleare, Sezione di Napoli, Strada Comunale Cinthia, 80126 Napoli, Italy
- ⁴⁸ Université de Strasbourg, CNRS, Observatoire Astronomique de Strasbourg, UMR 7550, 67000 Strasbourg, France
- ⁴⁹ MIT Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139, USA
- ⁵⁰ Graduate School of Science, Division of Particle and Astrophysical Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan
- ⁵¹ Hiroshima Astrophysical Science Center, Hiroshima University, 1-3-1 Kagamiyama, Higashi-Hiroshima, Hiroshima 739-8526, Japan
- ⁵² University of Maryland, Baltimore County, Baltimore, MD 21250, USA
- ⁵³ NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA
- ⁵⁴ Center for Research and Exploration in Space Science and Technology, NASA/GSFC, Greenbelt, MD 20771, USA

- ⁵⁵ Department of Physics, University of Hong Kong, Pokfulam, Hong Kong
- ⁵⁶ Department of Astronomy and Astrophysics, Pennsylvania State University, University Park, PA 16801, USA
- ⁵⁷ Université Grenoble Alpes, CNRS, IPAG, 38000 Grenoble, France
- ⁵⁸ Dipartimento di Fisica e Astronomia, Università degli Studi di Padova, Via Marzolo 8, 35131 Padova, Italy
- ⁵⁹ Dipartimento di Fisica, Università degli Studi di Roma “Tor Vergata”, Via della Ricerca Scientifica 1, 00133 Roma, Italy
- ⁶⁰ Department of Astronomy, University of Maryland, College Park, Maryland 20742, USA
- ⁶¹ Mullard Space Science Laboratory, University College London, Holmbury St Mary, Dorking, Surrey RH5 6NT, UK
- ⁶² Anton Pannekoek Institute for Astronomy & GRAPPA, University of Amsterdam, Science Park 904, 1098 XH Amsterdam, The Netherlands