

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

**Please provide feedback**

Please support the ScholarWorks@UMBC repository by emailing [scholarworks-group@umbc.edu](mailto:scholarworks-group@umbc.edu) and telling us what having access to this work means to you and why it's important to you. Thank you.

EGU22-10688

<https://doi.org/10.5194/egusphere-egu22-10688>

EGU General Assembly 2022

© Author(s) 2022. This work is distributed under the Creative Commons Attribution 4.0 License.



## Multi-Spacecraft Observations of Interplanetary Shocks

**Oksana Kruparova**<sup>1,2</sup>, Vratislav Krupar<sup>1,2</sup>, and Adam Szabo<sup>2</sup>

<sup>1</sup>Goddard Planetary Heliophysics Institute, University of Maryland, Baltimore County, Baltimore, MD 21250, USA

<sup>2</sup>Heliospheric Physics Laboratory, Heliophysics Division, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

Interplanetary (IP) shocks provide us with a unique opportunity to extensively investigate properties of collisionless shocks using in situ measurements under a wide range of upstream conditions. Here we report a case study of several IP shock crossings observed by the Wind, Solar and Heliospheric Observatory (SOHO), Advanced Composition Explorer (ACE), and Deep Space Climate Observatory (DSCOVR) spacecraft. By applying a simple timing method to multipoint measurements, we are able to investigate their characteristic spatiotemporal features. We assume that an IP shock can be represented by a moving plane with a constant velocity, when observed at closely separated points in space and time. We compared IP shock parameters obtained with the timing method with those obtained using the magnetic coplanarity, the mixed mode methods, and Rankine-Hugoniot jump relations.