

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Sub-Alfvenic/Super-Sonic Impulsive Structures in the Magnetosphere:

*First Results from Hybrid Fluid-Kinetic Modeling and
Comparison with MMS Observations*

Alexander S. Lipatov^{a,b}, Levon A. Avanov^b and Barbara L. Giles^b

^a Goddard Planetary Heliophysics Institute UMBC, Baltimore MD


^b NASA GSFC, Greenbelt, MD 20771

Fall AGU 2019. Updated for MMS Seminar (1.7.20 - GSFC)

2. Motivation and Applications:


- 1. Unique MMS observations of impulsive structures.
- 2. Plasma physics of impulsive structures inside the inner magnetosphere and ionosphere. Active experiments in space (AMPTE). Plasma clouds in the ionosphere (HANE).
- 3. Interaction between the moon's exosphere and planetary magnetospheres. Formation of the whistler and Alfv'en wing near the Parker Solar Probe.
- 4. Astrophysical explosions. Observations and modeling on the Large Plasma Device (UCLA). Magnetic reconnection modeling on the Large Plasma Device (UMBC).

3. Scheme of the Earth magnetosphere, radiation belts and interplanetary shock




4. MMS satellite position at time 2016-03-07 20:00:00 UTC

MMS Location for 2016-03-07 20:00:00 UTC




5. MMS observations of the impulsive structures at time 2016-03-07

20:00:00 UTC. Density, bulk velocities and magnetic field



6. MMS observations of the impulse at time 2016-03-07 20:15:00– 20:19:00 UTC. Density, bulk velocities and magnetic field




7. Questions to be answered with hybrid modeling:

- **1. Plasma Cloud Formation and ionospheric plasma? Percolation/Reconnection at the magnetopause**
(Akhavan-Tafti et al. [2018, 2019]).
- **2. Formation of the shock-like impulses, whistler/shear Alfvénic waves, collapsing diamagnetic cavern triggered by plasma cloud** (Recent Rev. by Winske, Huba, Niemann et al. [2019], Ed. by J.E. Borovsky).
- **3. Particle velocity distribution function dynamics and wave-particle interactions in plasma clouds environment.**

8. Hybrid Model

- *Kinetic - ions.*
- *Fluid - electrons. Scalar (tensor) electron pressure. Electron inertia.*
- *Interpenetrating flows*
- *Effects of finite ion gyroradius estimated with thermal and bulk velocities: $k_{\perp}\rho_{ci} \geq 1$, $k_{\parallel}\rho_{ci} \geq 1$ and $l \approx \rho_{ci}$; $\omega \approx \Omega_{ci}$*
- *Energetic ion gyroradii are about 10^4 km in the ring current and outer radiation belt*
- *The modeling tool: (a) **Standard PIC** [Harlow, 1957, LANL Report]; (b) **Shape Function Kinetics (SFK)** [Larson & Young 2015] is a logical extension of **Complex Particle Kinetics (CPK)** [Hewett 2003; Lipatov 2012] and **Finite Mass Method (FMM)** [Yserentant et al. 1996-2007]*
- *The SFK and CPK aim to bridge the gap between continuum and kinetic methods. This method may save a computational resources by factor more than 100 in compare with Standard PIC method*

9. Scheme of the plasma cloud expansion. From Winske [2014].



10. Early research for dense plasma cloud/debris expansion in low density ambient background plasma.




Figure 1: Dense plasma cloud expansion in low density ambient plasma at the initial stage. **Excitation of the right-hand-polarized whistler waves along the magnetic field and left-hand polarized whistler waves in the opposite direction.** 2-D cuts (left), 1-D cuts (right) of 3-D modeling from [Lipatov 1996; 2002]. 2.5-D in [Lipatov, Sharma, & Papadopoulos 1994]. **Shock formation by explosive debris** was observed in 2.5-D [Winske, Gary 2007], in 3-D Hewett, Larson, Brecht [2011]. See also Golubev et al. [1979]; Bashurin et al. [1983]; Antonov et al. [1985]; Brecht, Thomas [1987].

11. Set up the plasma cloud and ambient plasma configuration for our modeling:

Ambient upstream plasma:

density: $N_p = 2.5 \text{ cm}^{-3}$;

thermal velocity: $V_{th,p} = 100.0 \text{ km/s}$;

$V_{th,e} = 1000.0 \text{ km/s}$;

Alfvenic velocity: $V_A = 1217 \text{ km/s}$;

bulk U: $U = (80; 80; 20) \text{ km/s}$

B-field: $B = (40; 40; 20) \text{ nT}$




Plasma cloud:

Gaussian 3-D profile for density:

$N_i = 400.0 \text{ cm}^{-3}$;


$V_{th,i} \approx 120 \text{ km/s}; V_{th,e} \approx 2500 \text{ km/s}$;

bulk U: $U = (350.0; 250.0; 0.0) \text{ km/s}$




12. After the passage of the cloud. Perturbations in magnetic field and density



profiles. $U_0 = 100 \text{ km/s}$ at time $t=2.5 \text{ sec.}$




13. After the passage of the cloud. Perturbations in magnetic field and density profiles. $U_0 = 100 \text{ km/s}$ at time $t=3.75 \text{ sec.}$





14. Non-Maxwellian ion VDF's (a,b,c-ambient) along x axis at time $t=3.5$ sec.




15. Non-Maxwellian ion VDF's (a,c,d-cloud) along x axis at time $t=3.5$ sec.





16. Non-Maxwellian ion VDF's (a,b,c -ambient) along y axis at time $t=3.5$ sec.



17. Non-Maxwellian ion VDF's (a,c,d-cloud) along y axis at time $t=3.75$ sec.






18. 1-D cuts of profiles for velocity and magnetic field along x axis at time $t=3.5$ s.




19. 1-D cuts of profiles along x axis at time $t=3.5$ sec.



Quasi-overshoot and fluctuations in density profile





20. After the passage of the cloud. Perturbations in magnetic field and density profiles. $U_0 = 100$ km/s at time t=4.58 sec.






21. Comparison between MMS observation and modeling. Density profiles.



Quasi-overshoot and fluctuations in density profile



22. Ion VDFs. MMS observations (top) & modeling (bottom, left).



23. Summary

- 1. The interaction between plasma cloud and moving low-plasma beta ambient plasma produced a complicated 3-D structure of the plasma cloud.
- 2. The plasma cloud forms a strong whistler/shear Alfvén waves directed along the external magnetic field while a strong compression waves are formed across the external magnetic field.
- 3. A strong heating and acceleration of the ambient ions are observed in the regions at the interface between the plasma cloud and ambient plasma.
- 4. The modeling shows a strong depletion in the ambient plasma due to the cloud expansion. Such effect is a similar to a formation of the diamagnetic cavern observed in the active experiment for plasma injection (AMPTE) into the magnetosphere and plasma cloud generation in the ionosphere.
- 5. The anisotropy of the ion VDF's may trigger EMIC waves and other instabilities like the mirror-balloonning instability.

Research was supported by the NASA.