Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing <u>scholarworks-group@umbc.edu</u> and telling us what having access to this work means to you and why it's important to you. Thank you.

Second law of thermodynamics for quantum correlations

Akram Touil^{1,*}, Barış Çakmak², and Sebastian Deffner^{1,3}

¹Department of Physics, University of Maryland, Baltimore County, Baltimore, MD 21250, USA

 $^2 {\rm College}$ of Engineering and Natural Sciences, Bahçeşehir University, Beşiktaş, İstanbul 34353, Turkey

³Instituto de Física 'Gleb Wataghin', Universidade Estadual de Campinas, 13083-859, Campinas, São Paulo, Brazil

E-mail: *akramt1@umbc.edu

Abstract. It is an established fact that quantum coherences have thermodynamic value. The natural question arises, whether other genuine quantum properties such as entanglement can also be exploited to extract thermodynamic work. In the present analysis, we show that the ergotropy can be expressed as a function of the quantum mutual information, which demonstrates the contributions to the extractable work from classical and quantum correlations. More specifically, we analyze bipartite quantum systems with locally thermal states, such that the only contribution to the egrotropy originates in the correlations. Our findings are illustrated for a two-qubit system collectively coupled to a thermal bath.

1. Introduction

What is a resource in thermodynamics? From the inception of the theory, the question appears rather simple to answer – namely a thermodynamic resource is any energy that can be extracted from, e.g., a heat or particle reservoir, and transformed into work [1]. However, it has been debated since essentially the beginnings of thermodynamics if and to what extent *information* can also be considered a resource [2–5]. Remarkably, the *thermodynamics of information* [6, 7] was fully established only rather recently, which was spurred by developing the stochastic thermodynamics with feedback [8, 9] and by proposing the notion of information reservoirs [10].

Evidently, describing information in quantum systems is more subtle, and hence also the thermodynamics of quantum information requires more thorough analyses [11, 12]. This area of research has received a boost by the realization that quantum thermodynamics contributes profoundly to the development of new generation quantum technologies [13, 14]. Among these emerging technologies, in particular quantum thermal machines [15–20] and quantum information engines [21– 25], aka quantum computers [26] necessitate a comprehensive study of quantum information as a thermodynamic resource. In this context, it is important to realize that from a thermodynamic perspective quantum information as quantified by the von Neumann entropy is not the only notion of information to be considered. Rather, understanding the contribution of information encoded in marginals [27–29], and in particular the thermodynamic value of genuine quantum correlations [30–32] is instrumental.

Therefore, the present analysis specifically focuses on the role that correlations (quantum or classical) play in the maximum extractable work, i.e., the *ergotropy*. In the literature, this problem was partially addressed from different perspectives, by relating work extraction to either coherences [33-36] or correlations [37-45], for a wide variety of scenarios. Nevertheless, all previous studies either rely on correlations with an ancillary system together with a feedback mechanism [37-40], are restricted to specific dynamical models [35, 42], or consider certain classes of states [41]. To the best of our knowledge, a general mathematical relationship between the correlations among the constituents of a quantum system and its work content is still lacking.

To address this void in our understanding of the thermodynamics of correlations, we prove a direct relationship between the ergotropy and the quantum mutual information in a bipartite, and locally thermal, quantum state. The locally thermal states are a judicious choice such that the only resources available for work extraction are total bipartite correlations. Furthermore, we derive an equality relating the ergotropy to both the quantum mutual information and the bound ergotropy [46]. We illustrate and analyze these general results in the specific context of an array of qubits collectively coupled to a thermal bath at a finite temperature. Finally, we upper bound the average power that can be extracted from an arbitrary quantum state. Our results explicitly quantify the role of correlations in the process of work extraction by means of cyclic and unitary processes, during which the system of interest is isolated and the only available resources are in the form of bipartite correlations.

2. Ergotropy and the process of work extraction

Ergotropy is the maximum amount of work that can be extracted from a quantum system by means of cyclic and unitary operations [47]. From a thermodynamic standpoint, work is then simply given by the change of internal energy, since the corresponding process is unitary, or in other words thermally isolated. Therefore, we consider the dynamics of a system governed by the Hamiltonian $H_{total} = H + \Gamma(t)$, where H denotes the self-Hamiltonian of the system and $\Gamma(t)$ is a time-dependent coupling term responsible for the extraction of work during time τ . Note that by construction $\Gamma(t)$ fulfills $\Gamma(0) = 0$ and $\Gamma(\tau) = 0$, such that the operation is cyclic with respect to H.

Now, consider a quantum system described by $H = \sum_{i=1}^{d} \varepsilon_i |\varepsilon_i\rangle \langle \varepsilon_i|$ and quantum state $\rho = \sum_{j=1}^{d} r_j |r_j\rangle \langle r_j|$, such that $\varepsilon_i \leq \varepsilon_{i+1}$ and $r_j \geq r_{j+1}$. The ergotropy is then calculated by performing an optimization over all possible unitary operations to achieve a final state that has the minimum average energy with respect to H,

$$\mathcal{E}(\rho) = \operatorname{tr} \{H\rho\} - \min_{U} \left\{ \operatorname{tr} \{HU\rho U^{\dagger}\} \right\} = \operatorname{tr} \{H(\rho - P_{\rho})\}, \qquad (1)$$

where $P_{\rho} \equiv \sum_{k} r_{k} |\varepsilon_{k}\rangle \langle \varepsilon_{k}|$ is called the *passive state* [48]. By plugging the explicit form of P_{ρ} in the equation above, we obtain the well-known expression [47]

$$\mathcal{E}(\rho) = \sum_{i,j} r_j \varepsilon_i \left(|\langle r_j | |\varepsilon_i \rangle|^2 - \delta_{ij} \right).$$
⁽²⁾

The specific unitary U that takes an arbitrary state ρ to its corresponding passive state P_{ρ} is given by $U = \sum_{k} |\varepsilon_{k}\rangle \langle r_{k}|$. Hence, we have a general form for the potential



Figure 1. Sketch of the successive processes (i) and (ii), where the build up of correlations between the partitions A and B (process (i)) enables work extraction by means of unitary evolution (process (ii)).

 $\Gamma(t)$, that generates the desired unitary operation, such that [47]

$$\Gamma(t) = \dot{\varphi}(t) \exp\left(-iHt/\hbar\right) \Lambda \exp\left(iHt/\hbar\right),\tag{3}$$

where $\varphi(0) = \dot{\varphi}(0) = \dot{\varphi}(\tau) = 0$, $\varphi(\tau) = \tau$, and in the interaction picture $U_I(\tau) \equiv \exp(iH\tau/\hbar) U = \exp(-i\Lambda\tau/\hbar)$. Note that the freedom to choose φ implies that the potential $\Gamma(t)$ is not unique. Finally, it is interesting to note that with the exception of thermal states, it is possible to extract work from multiple copies of passive states by processing them collectively [46]. For this reason, thermal states are also referred to as *completely passive states* [48].

3. Extractable work from correlations

3.1. Ergotropy and the mutual information

For the following analysis, we consider a quantum system S, that can be separated into two partitions A and B. For such scenarios, we now prove a direct relationship between the ergotropy and the mutual information,

$$\mathcal{I}(A:B) \equiv S(\rho_A) + S(\rho_B) - S(\rho), \tag{4}$$

where $S(\rho_i) = -\text{tr} \{\rho_i \ln(\rho_i)\}$ denotes the von Neumann entropy, and ρ is the quantum state of \mathcal{S} .

Our goal is now to assess how much work can be extracted from the quantum correlations between A and B. To this end, we consider a two-stroke operation on S: (i) correlations are built-up and S is driven into thermal (passive) states in A and B; and (ii) work is extracted under a cyclic, unitary operation on S, for which we can compute the ergotropy.

More specifically, consider the following situation: (i) the quantum system $S = A \otimes B$ evolves from $t = -\tau_0$ to $t = 0^-$ under

$$H(t) = H_A \otimes \mathbb{I}_B + \mathbb{I}_A \otimes H_B + H_I(t), \tag{5}$$

where $H_I(t)$ contains interactions among the subsystems and their interactions with the environment. We assume that at t = 0 the interaction Hamiltonian is negligible compared to the self-Hamiltonians of A and B, i.e., $H(0) = H_A \otimes \mathbb{I}_B +$ $\mathbb{I}_A \otimes H_B \equiv H$. As a result, we evolve an initially uncorrelated density matrix $\rho(-\tau_0) = \rho_A(-\tau_0) \otimes \rho_B(-\tau_0)$ to a steady state density matrix ρ such that both $\rho_A = \operatorname{tr}_B \{\rho\} = \exp(-\beta H_A)/Z_A$ and $\rho_B = \operatorname{tr}_A \{\rho\} = \exp(-\beta H_B)/Z_B$ are thermal states at inverse temperature β .

Then, (ii) work is extracted during time τ . As outlined above, the total Hamiltonian reads,

$$H(t) = H_A \otimes \mathbb{I}_B + \mathbb{I}_A \otimes H_B + \Gamma(t), \tag{6}$$

where $\Gamma(0) = \Gamma(\tau) = 0$. It is then a simple exercise to show that the ergotropy $\mathcal{E}(\rho)$ can be written as

$$\beta \mathcal{E}(\rho) = \mathcal{I}(A:B) - D(P_{\rho} || \rho_A \otimes \rho_B), \tag{7}$$

where $D(\rho||\sigma) = \text{tr} \{\rho \ln(\rho) - \rho \ln(\sigma)\}$ is the relative entropy, between the states ρ and σ [26]. Note that ρ_A and ρ_B and $\mathcal{I}(A:B)$ are evaluated at t = 0 (see Fig. 1).

Equation (7) constitutes our main result and shows that there are two additive contributions to the ergotropy of locally thermal states. While the mutual information between A and B is the main resource for the finite ergotropy, the distance between P_{ρ} and the tensor product of local states, at the beginning of the work extraction process, reduces the amount of ergotropy, owing to the non-negativity of the relative entropy. This suggests that the *optimal* work extraction process erases all correlations (quantum and classical) between the subsystems at the end of the process. However, in general this is not possible due to the fact that the entropies of the states P_{ρ} and $\rho_A \otimes \rho_B$ can be different in magnitude.[‡].

It is also interesting to note that Eq. (7) implies an "inverse Landauer" inequality for correlation,

$$\beta \mathcal{E}(\rho) \le \mathcal{I}(A:B). \tag{9}$$

The maximum amount of work that can be extracted from a closed, bipartite quantum system with locally thermal states, is given by the mutual information between its subsystems. The bound is saturated if and only if $P_{\rho} = \rho_A \otimes \rho_B$.

Finally, it is interesting to see how the present findings relate to established work extraction schemes, for which S is in contact with a heat bath. In general, the nonunitary schemes outperform the unitary procedure since the presence of the bath lifts the constant entropy restriction on the system. In this case, the maximum amount of extractable work $\mathcal{E}_{\beta}(\rho)$ is determined by the difference in the non-equilibrium free energies of the state at hand and the thermal state, $\mathcal{E}_{\beta}(\sigma) = F_{\beta}(\sigma) - F_{\beta}(\sigma_{\beta})$, where $F_{\beta}(\sigma) = \operatorname{tr} \{H\rho\} - 1/\beta S(\sigma) \ [40, 49-51]$. For a general bipartite system, it is then possible to write [40]

$$\mathcal{E}_{\beta}(\rho) = F_{\beta}(\rho) - F_{\beta}(\rho_{A,\beta} \otimes \rho_{B,\beta}),$$

= $\mathcal{E}_{\beta}(\rho_A) + \mathcal{E}_{\beta}(\rho_B) + 1/\beta \mathcal{I}(A:B).$ (10)

In our case, we have $\mathcal{E}_{\beta}(\rho_A) = \mathcal{E}_{\beta}(\rho_B) = 0$. Comparing this with Eq. (7) we can immediately identify the difference between the maximum extractable work in unitary

‡ Equation (7) can be equivalently obtained using the fact that for any two different density matrices ρ_1 and ρ_2 , with $S(\rho_1) = S(\rho_2)$, we have

$$\beta \operatorname{tr} \{ (\rho_1 - \rho_2) H \} = D(\rho_1 || \rho_\beta) - D(\rho_2 || \rho_\beta),$$
(8)

where $\rho_{\beta} = \exp(-\beta H)/Z$ is the global thermal state.

and non-unitary approaches as $D(P_{\rho}||\rho_A \otimes \rho_B)$. Shortly, we will see the present work extraction scheme (for process (ii)) can be further improved, when we utilize multiple copies of the same quantum state.

3.2. Bound ergotropy and multipartite correlations

For a given entropy, the quantum state with the minimum average energy is the thermal state. For states P_{ρ} , that are passive, but not thermal, i.e., not completely passive, the remaining energy in P_{ρ} can be accessed by implementing a secondary process. It has been shown that this secondary process can be constructed through multiple copies of P_{ρ} , and the additionally extractable work has been dubbed *bound* ergotropy $\mathcal{E}_{b}(\rho)$ [46]. It can be written as

$$\mathcal{E}_b(\rho) = \operatorname{tr}\left\{ (P_\rho - P_\rho^{\operatorname{th}})H \right\} \ge 0, \tag{11}$$

where P_{ρ}^{th} is the thermal state corresponding to P_{ρ} such that $S(P_{\rho}) = S(P_{\rho}^{\text{th}})$. Exploiting Eqs. (7) and (8) we further have

$$\beta\left(\mathcal{E}(\rho) + \mathcal{E}_b(\rho)\right) = \mathcal{I}(A:B) - D(P_{\rho}^{\text{th}} || \rho_A \otimes \rho_B).$$
(12)

Again noting that the relative entropy is non-negative, we can write

$$\beta \left(\mathcal{E}(\rho) + \mathcal{E}_b(\rho) \right) \le \mathcal{I}(A:B). \tag{13}$$

Equation (13) is a tighter version of the inverse Landauer's principle (7), and it is in fact tight. Below in Sec. 4, we will show that the bound can be saturated for qubits collectively coupled to a thermal bath.

Moreover, it is interesting to note that the ergotropy is a non-extensive quantity. Namely, it is easy to see from the strong subadditivity of the von Neumann entropy that the global ergotropy, $\mathcal{E}_G(\rho)$, for N copies of ρ is greater or equal to $N\mathcal{E}(\rho)$. Hence, we have

$$\beta \mathcal{E}_G(\rho) \equiv \beta N \left(\mathcal{E}(\rho) + \mathcal{E}_b(\rho) \right) \le N \mathcal{I}(A:B) \,. \tag{14}$$

Note, however, that the bound ergotropy can be extracted by acting globally on the N copies of ρ (since P_{ρ} is not completely passive). Inequality (14) is saturated if and only if the thermal state can be expressed as follows $\rho_A \otimes \rho_B = P_{\rho}^{\text{th}}$.

We conclude this section by noting that Eqs. (7) and (14) can be readily generalized to multipartite correlations. Using Eq. (8), and assuming that S is composed of "k" thermal states correlated with each other, we obtain

$$\beta \mathcal{E}(\rho) \le \mathcal{I}(A_1 : A_2 : \dots : A_k), \tag{15}$$

and

$$\beta N \left(\mathcal{E}(\rho) + \mathcal{E}_b(\rho) \right) \le N \mathcal{I}(A_1 : A_2 : \dots : A_k), \tag{16}$$

where $\mathcal{I}(A_1 : A_2 : ... : A_k) = D(\rho_{A_1...A_k} || \rho_{A_1} \otimes ... \otimes \rho_{A_k})$ is the multipartite mutual information between the k partitions of \mathcal{S} [52].

3.3. Ergotropy and quantum discord

We continue with a closer analysis of the nature of the correlations. It has been established that the quantum mutual information quantifies the amount of total correlations between two parties A and B [53, 54]. In particular, this means that $\mathcal{I}(A:B)$ is comprised of classical *and* quantum correlations. An accepted measure for the purely quantum contribution is the *quantum discord* [54], which is typically written as

$$\mathfrak{D}(A:B) = \mathcal{I}(A:B) - J(A:B), \tag{17}$$

where J(A:B) is the Holevo information [26] that quantifies the maximum amount of classical information that is determined from optimized generalized measurements on *B*. Hence, J(A:B) quantifies the maximal, classical information that *B* can carry about *A*. Therefore, $\mathfrak{D}(A:B)$ is the genuinely quantum information. Using the definition of the quantum discord, Eq. (7) can be written as

$$\beta \mathcal{E}(\rho) = \mathfrak{D}(A:B) + J(A:B) - D(P_{\rho} || \rho_A \otimes \rho_B), \tag{18}$$

which highlights the interplay of classical and quantum correlations. It is then instructive to consider under what conditions and for what states ρ_0 the egrotropy vanishes, $\mathcal{E}(\rho_0) = 0$. For such ρ_0 we have

$$\mathfrak{D}(A:B) + J(A:B) = D(P_{\rho_0} || \rho_A \otimes \rho_B).$$
(19)

We immediately conclude that the existence of correlations (quantum or classical) is necessary, but not sufficient, for a non-zero ergotropy in locally thermal states.

3.4. Ergotropy and system-environment correlations

As alluded to above, $\mathcal{E}(\rho)$ is a lower bound on the work that can be extracted by non-unitary operations on \mathcal{S} ,

$$\mathcal{E}(\rho) \le \mathcal{E}_{\beta}(\rho). \tag{20}$$

which follows from exploiting correlations between S and its environment [37–40]. Additionally, Eq. (7) can also be written as [55],

$$\beta \mathcal{E}(\rho) = \Delta S_A + \Delta S_B - \mathcal{I}(\mathcal{S}:E) - \beta \langle Q \rangle - D(P_\rho || \rho_A \otimes \rho_B), \qquad (21)$$

where $\langle Q \rangle$ is the heat exchanged between system and environment, and $\mathcal{I}(S:E)$ quantifies the buildup of correlations. Equation (21) demonstrates that systemenvironment correlations and the dissipated heat diminish the amount of work that can be extracted by unitary processes. This insight is complementary to what has been shown in the literature [38, 56, 57], and emphasizes our different approach that focuses on work extraction by means of cyclic and unitary operations on a state that is locally completely passive.§

4. Illustrative case study: ergotropy from X-states

After having established the conceptual framework, the remainder of the analysis is dedicated to an instructive case study. In particular, we elucidate the conditions and physical mechanisms that lead to locally thermal states.

[§] As a side note, it is interesting to consider our result, in Eq. (21), in context of work extraction from information scrambling [55, 58-63]. However, a thorough analysis of extracting work from scrambled states, or even from quantum chaos, is beyond the scope of the present work, see also Refs. [64, 65].

4.1. Two-qubit systems and X-states

Consider now that S is comprised of two qubits, which are initially prepared in an arbitrary quantum state. Note that working with qubits makes the analysis particularly simple, since any diagonal qubit state can be described by a thermal state at an effective inverse temperature β . A straightforward, though not the most general, way to then have locally thermal states is if S relaxes into an X-state

$$\rho(t) = \begin{pmatrix}
\rho_{11} & 0 & 0 & \rho_{14} \\
0 & \rho_{22} & \rho_{23} & 0 \\
0 & \rho_{23}^* & \rho_{33} & 0 \\
\rho_{14}^* & 0 & 0 & \rho_{44}
\end{pmatrix}.$$
(22)

Such states have been widely studied in the literature [66], as they can be found in generalized Pauli channels [67] or in collective dephasing models for two-qubit systems [68]. Moreover, the ground state of one-dimensional spin chains that are invariant under translations, and parity transformations, i.e., exhibit \mathbb{Z}_2 symmetry, also have reduced bipartite density matrices in X-shape [69–71]. For a more detailed discussion of X-states we refer to the literature [66].

4.2. A physical model: Collective dissipation

We continue with an even more specific scenario, and now consider two interacting qubits, which are assumed to behave as point-like dipoles with identical dipole moments, and which are immersed in a finite-temperature environment composed of thermal photons resonant with the qubits' transition frequency. The master equation governing the dynamics is given by [72-77]

$$\frac{\partial \rho}{\partial t} = -\frac{i}{\hbar} [(H_0 + H_d), \rho] + \mathcal{D}_-(\rho) + \mathcal{D}_+(\rho) = \mathcal{L}(\rho), \qquad (23)$$

where $H_0 = \hbar \omega (\sigma_1^+ \sigma_1^- + \sigma_2^+ \sigma_2^-)$ and $H_d = \hbar f (\sigma_1^+ \sigma_2^- + \sigma_2^+ \sigma_1^-)$ are the self-Hamiltonian of the whole system and the interaction Hamiltonian between the qubits, respectively. As usual, $\sigma_i^+ = |e_i\rangle\langle g_i| \ (\sigma_i^- = (\sigma_i^+)^{\dagger})$ is the raising (lowering) operator with $i \in \{1, 2\}$ denoting the qubit label. The explicit form of the dissipators describing the interaction with the bath reads

$$\mathcal{D}_{-}(\rho) = \sum_{i,j=1}^{2} \gamma_{ij} (\bar{n}+1) (\sigma_{j}^{-} \rho \sigma_{i}^{+} - \frac{1}{2} \{\sigma_{i}^{+} \sigma_{j}^{-}, \rho\}),$$

$$\mathcal{D}_{+}(\rho) = \sum_{i,j=1}^{2} \gamma_{ij} \bar{n} (\sigma_{j}^{+} \rho \sigma_{i}^{-} - \frac{1}{2} \{\sigma_{i}^{-} \sigma_{j}^{+}, \rho\}).$$
(24)

Here, $\bar{n} = [\exp(\beta_e \omega) - 1]^{-1}$ is the mean number of photons at the temperature of the environment β_e , and γ_{ij} are the spontaneous decay rates. Without loss of generality, we assume $\gamma_{ij} = \gamma_{ji}$.

Depending on the spatial configuration of the system qubits, the model above has two distinct limits. If the qubits are well separated, i.e., the distance between them is much larger than the wavelength of the thermal photons in the environment, they individually couple to the bath and their steady-state is simply described by the Gibbs state, at the bath temperature, $\rho_{\beta_e} = \exp(-\beta_e H_0)/Z$. Naturally, such a



Figure 2. Ergotropy and bound ergotropy as a function of the initial state parameter c for $\beta_e = 0.01$ (a), $\beta_e = 1$ (b) and $\beta_e = 10$ (c).

steady-state is completely passive. On the other hand, when the qubits are closely packed, such that the separation between them is much smaller than the wavelength of the thermal photons, they collectively couple to the environment. In this regime, the qubits become indistinguishable due to the impossibility of resolving which qubit absorbed or emitted a photon. Thus, their steady-state admits coherences in the energy eigenbasis [72, 73] and has the following form [73]

$$\rho_{ss} = (1-c) |\psi_{-}\rangle \langle\psi_{-}|$$

$$+ cZ_{+}^{-1} (\beta_{e}) (\exp (-2\omega\beta_{e}) |\psi_{ee}\rangle \langle\psi_{ee}| + \exp (-\omega\beta_{e}) |\psi_{+}\rangle \langle\psi_{+}| + |\psi_{gg}\rangle \langle\psi_{gg}|).$$
(25)

Here, $|\psi_{gg}\rangle = |gg\rangle$, $|\psi_{ee}\rangle = |ee\rangle$, $|\psi_{\pm}\rangle = |ge\rangle \pm |eg\rangle /\sqrt{2}$, $c = \langle \psi_{gg} | \rho_0 | \psi_{gg} \rangle + \langle \psi_{ee} | \rho_0 | \psi_{ee} \rangle + \langle \psi_{+} | \rho_0 | \psi_{+} \rangle$, and $Z_{+} (\beta_e) = 1 + \exp(-\omega\beta_e) + \exp(-2\omega\beta_e)$.

Consequently, the steady-state preserves some information about the initial state of the dynamics through the parameter "c", and thus it is not unique. Only recently, it has been shown that such states possess a finite amount of ergotropy [35] for a wide range of initial states and environment temperatures. Further, Eq. (25) is also in the X-shape form (22) with $\rho_{14} = 0$. Note, however, that the local temperature of the qubits, β , is generally different from the bath temperature β_e . In fact, it is a simple exercise to show that

$$\beta = \frac{1}{\omega} \ln \left[\frac{1 + 2\cosh(\beta_e \omega) + 2c\sinh(\beta_e \omega)}{1 + 2\cosh(\beta_e \omega) - 2c\sinh(\beta_e \omega)} \right].$$
 (26)

As we have seen above (7), to maximize the ergotropy the target passive state needs to be as close as possible to the product state $\rho_A \otimes \rho_B$. However, achieving such a passive state through any unitary process is generally not possible since the states P_{ρ} and $\rho_A \otimes \rho_B$ have different values of entropy. Despite this fact, it is still possible to further process the passive state P_{ρ} provided that it is not a completely passive state, and we have access to, and ability to act on, multiple copies of it. To this end, we are interested in the bound ergotropy $\mathcal{E}_b(\rho)$ (11), which also can be determined analytically.

In Fig. 2, we plot $\mathcal{E}(\rho)$ and $\mathcal{E}_b(\rho)$ as a function of c, that is, for different initial states of the open system dynamics. Note that the ergotropy is evaluated using Eq. (7). In contrast to the ergotropy, which bears a finite value for nearly all values of c at high environment temperatures (low β_e), the overall magnitude of bound ergotropy is the largest at low environment temperatures (high β_e). However, the amount of the bound ergotropy does not represent a significant amount as compared to the ergotropy. A two-qubit system with a self-Hamiltonian of H_0 has an energy gap of $2\hbar\omega$, which



Figure 3. Total ergotropy $(\mathcal{E} + \mathcal{E}_b)$ and $\mathcal{I}(A : B)/\beta$ as a function of the initial state parameter c for $\beta_e = 0.01$ (a), $\beta_e = 1$ (b) and $\beta_e = 10$ (c).

corresponds to the highest amount of ergotropy that one can get out of this system if both qubits were to be in their excited states. Comparing \mathcal{E}_b with the maximum we observe for the steady-states of the considered model and with the absolute maximum of $2\hbar\omega$, we conclude that \mathcal{E}_b is reasonably smaller than both of these quantities, and the difference is more pronounced for the latter case. This result can be expected considering that the bound ergotropy is actually the amount of extractable work from a state that is already passive, and can only be accessed by acting on multiple copies.

The zero of the ergotropy corresponds to situations for which S is prepared in a thermal state at inverse temperature β_e . For such states it has actually been shown that the quantum system can not maintain any ergotropy in the long time limit [35]. Curiously, this behavior is observed for a range of values of c, cf. Fig. 2.

Finally, in Fig. 3, we depict the global ergotropy, $\mathcal{E}_G = \mathcal{E} + \mathcal{E}_b$, together with $\mathcal{I}(A : B)/\beta$. We observe that the bound is far from tight for high environment temperatures as displayed in Fig. 3 (a). The only point where we have equality $\beta \mathcal{E}_G = \mathcal{I}(A : B)$ is when both quantities vanish at c = 0.75. This value of the parameter c corresponds to that of a thermal initial state at the considered environment temperature and hence neither ergotropy nor any form of correlation is generated as a result of the interaction with the bath. As we lower the temperature in Fig. 3 (b) and (c), i.e., increase β_e , we observe that the bound is sharp for a wide range of initial states of the model. Therefore, it is interesting to observe that the ergotropy is aptly estimated by the mutual information, which is a lot simpler to compute for high-dimensional scenarios. Also note that the set of states for which we attain equality in Eq. (13) includes the ground initial state (c = 1), which does not contain any quantum or classical correlations before the open system evolution.

From a practical point of view, one problem that is critical in the extraction of the bound ergotropy is the preparation of multiple copies of a quantum state. In this sense, the present model for collective dissipation can be considered minimal in terms of the resources involved in the state preparation. An alternative approach, that is particularly useful in creating a number of copies of bipartite states, is a translationally invariant spin chain with \mathbb{Z}_2 symmetry [69–71]. As mentioned above, the ground states of such spin chains have reduced bipartite density matrices in the X-shape (especially see Sec. 3.3 of [70]), and are thus locally thermal. Although it is in general not an easy task to prepare a complex many-body system in its ground state, if achieved in a chain of N spins (with N as an even number), we end up with N/2 copies of the same X-shaped state.

5. Bound on the average extractable power

Finally, we would like to present few considerations on more practical matters. In all realistic scenarios, one is arguably more interested in the average power, rather than in the total amount of work that can be generated. Thus we now define

$$\mathcal{P}(\rho) \equiv \frac{\mathcal{E}(\rho)}{\tau} \ . \tag{27}$$

The natural question arises whether the subtleties of quantum dynamics, in particular the quantum speed limit [78], pose additional constraints. For unitary dynamics under driven Hamiltonians it has been shown that a generalized Mandelstam-Tamm bound takes the form [79]

$$\tau \ge \frac{\hbar}{\Delta E_{\tau}} \mathcal{L}\left(\rho, P_{\rho}\right),\tag{28}$$

where \mathcal{L} is the Bures angle [80, 81]

$$\mathcal{L}(\rho, P_{\rho}) = \arccos\left(\operatorname{tr}\left\{\sqrt{\sqrt{\rho}P_{\rho}\sqrt{\rho}}\right\}\right).$$
(29)

Further, ΔE_{τ} is the time-averaged variance of the Hamiltonian

$$\Delta E_{\tau} \equiv \frac{1}{\tau} \int_0^{\tau} dt \,\Delta H_t = \frac{1}{\tau} \int_0^{\tau} dt \,\sqrt{\langle H(t)^2 \rangle - \langle H(t) \rangle^2}.$$
(30)

While there are many formulations of the quantum speed limit, see Ref. [28, 78, 82, 83] and references therein, this version is particularly convenient for the present purposes. In particular, ΔE_{τ} is upper bounded by using the *finite energy constraint* [84, 85] and the expression of $\Gamma(t)$ in Eq. (3). We have,

$$(\exists \ \Omega > 0) \ (\forall t \in [0, \ \tau]); \ \mathrm{tr}\left\{H^2(t)\right\} \le \Omega^2, \tag{31}$$

which implies

$$(\exists \ \Omega > 0) \ (\forall t \in [0, \ \tau]); \ \mathrm{tr}\left\{\rho(t)H^2(t)\right\} \le \Omega^2.$$
(32)

The above conditions state that the energy bandwidth of the Hamiltonian is uniformly bounded, simply implying that the physical resources at our disposal are limited. Therefore, we can write

$$\mathcal{P}(\rho) \le \mathcal{I}(A:B) \frac{G(\Omega,\Lambda)}{\hbar\beta \mathcal{L}(\rho, P_{\rho})}.$$
(33)

Equation (33) gives an upper bound on the average extractable power that only depends on Ω (the upper bound on the bandwidth of the Hamiltonian H(t)), the time-independent potential Λ , the temperature T, the overlap between the initial state ρ and the passive state P_{ρ} , and the mutual information. Note that this result is independent of the process duration τ . We have,

$$G(\Omega, \Lambda) = \min\left\{\Omega, \sqrt{\left(\Omega + \langle\Lambda\rangle_0 + \operatorname{tr}\left\{H_{AB}\right\}\right)\left(\Omega - \langle\Lambda\rangle_0\right)}\right\},\tag{34}$$

such that $\langle \Lambda \rangle_0 = \text{tr} \{ \rho \Lambda \}$. Details of the derivation can be found in Appendix A.

6. Concluding remarks

We conclude the analysis with a few remarks on the direct implications of our main results. Interestingly, it is straightforward to define a notion of *efficiency* relating the amount of energy expended in process (i) to create correlations, and the useful work extracted during process (ii). The work of formation " W_f " quantifying the energetic cost associated with creating correlations was studied in Ref. [43], and it was found that it is directly related to the multipartite mutual information. Extending this result to arbitrarily correlated states we can define the upper bound on efficiency as $\eta \equiv \mathcal{E}(\rho)/W_f$, see also Ref. [86]. We leave the detailed analysis of this bound, and its relation to the mutual information, for future work. However, it is worth emphasizing already at this point that such an analysis will constitute an essential step towards a more complete thermodynamic description of quantum information processing.

In the present analysis, we derived a general equality relating ergotropy to the quantum mutual information between thermal partitions, A and B, of a bipartite quantum system S. Our conceptual considerations are based on two successive processes: we first create bipartite correlations as S is coupled to a thermal bath, then we decouple the system from the bath and extract the work from the resulting state of S, through a cyclic and unitary process. For such scenarios, we proved a relationship between ergotropy, bound ergotropy, and the quantum mutual information. The resulting inequalities were demonstrated for the experimentally relevant scenario of an array of qubits collectively coupled to a thermal bath at a finite temperature. Our results demonstrate that correlations shared within a quantum system in fact pertain thermodynamic value while being totally absent in its local states. Such a conclusion can prove to be useful in designing many-body quantum batteries, which are quantum systems utilized as a work reservoir, in a non-trivial setting.

Finally, we sketched an analysis of the maximal average power that can be extracted from quantum correlations. However, due to the intricacies of a more thorough study, we leave bounds on the instantaneous power for future work.

Acknowledgements

B. Ç. is supported by the BAGEP Award of the Science Academy and by the Research Fund of Bahçeşehir University (BAUBAP) under project no: BAP.2019.02.03. S.D. acknowledges support from the U.S. National Science Foundation under Grant No. DMR-2010127. This research was supported by grant number FQXi-RFP-1808 from the Foundational Questions Institute and Fetzer Franklin Fund, a donor advised fund of Silicon Valley Community Foundation (SD).

Appendix A. Bounding the average extractable power

In this appendix, we outline the derivation of Eq. (33). To this end, we consider Hamiltonians fulfilling the finite energy constraint. For such situations we have

$$\Delta E_{\tau} = \frac{1}{\tau} \int_{0}^{\tau} dt \sqrt{\langle H(t)^{2} \rangle - \langle H(t) \rangle^{2}},$$

$$\leq \frac{1}{\tau} \int_{0}^{\tau} dt \sqrt{\Omega^{2} - \langle H(t) \rangle^{2}},$$

$$\leq \frac{1}{\tau} \sqrt{\int_{0}^{\tau} dt \left(\Omega - \langle H(t) \rangle\right)} \sqrt{\int_{0}^{\tau} dt \left(\Omega + \langle H(t) \rangle\right)},$$
(A.1)

which follows from the Cauchy–Schwarz inequality. Using the fact that $0 \leq 1/\tau \int_0^\tau dt \operatorname{tr} \{\rho(t)H_{AB}\} \leq 1/\tau \int_0^\tau dt \operatorname{tr} \{H_{AB}\}$ when the self-Hamiltonian's eigenvalues are greater or equal to zero, we obtain

$$\Delta E_{\tau} \leq \sqrt{\Omega - \frac{1}{\tau} \int_{0}^{\tau} dt \, \langle \Gamma(t) \rangle} \, \sqrt{\Omega + \frac{1}{\tau} \int_{0}^{\tau} dt \, \langle \Gamma(t) \rangle + \operatorname{tr} \{H_{AB}\}}.$$
(A.2)

Furthermore, it can be shown that

$$\frac{1}{\tau} \int_0^\tau dt \, \langle \Gamma(t) \rangle = \operatorname{tr} \left\{ \rho \Lambda \right\} \equiv \langle \Lambda \rangle_0 \, . \tag{A.3}$$

In conclusion, the time-averaged variance is upper bounded by

$$\Delta E_{\tau} \le \sqrt{\left(\Omega + \langle \Lambda \rangle_0 + \operatorname{tr} \{H_{AB}\}\right) \left(\Omega - \langle \Lambda \rangle_0\right)}, \qquad (A.4)$$

from which we obtain Eq. (33).

References

- [1] Callen H B 1998 Thermodynamics and an Introduction to Thermostatistics (American Association of Physics Teachers)
- [2] Leff H S and Rex A F 2014 Maxwell's demon: entropy, information, computing (Princeton University Press)
- [3] Landauer R 1961 IBM Journal of Research and Development 5 183-191 URL https://doi. org/10.1147/rd.53.0183
- Bennett C H 1982 International Journal of Theoretical Physics 21 905-940 URL https: //doi.org/10.1007/BF02084158
- [5] Zurek W H 1989 Nature 341 119-124 URL https://doi.org/10.1038/341119a0
- [6] Parrondo J M R, Horowitz J M and Sagawa T 2015 Nat. Phys. 11 131 URL https://doi.org/ 10.1038/nphys3230
- [7] Wolpert D H 2019 J. Phys. A: Math. Theor. 52 193001 URL https://doi.org/10.1088/ 1751-8121/ab0850
- [8] Sagawa T and Ueda M 2008 Phys. Rev. Lett. 100 080403 URL https://link.aps.org/doi/10. 1103/PhysRevLett.100.080403
- [9] Sagawa T and Ueda M 2010 Phys. Rev. Lett. 104 090602 URL https://link.aps.org/doi/10. 1103/PhysRevLett.104.090602
- [10] Definer S and Jarzynski C 2013 Phys. Rev. X 3(4) 041003 URL https://link.aps.org/doi/ 10.1103/PhysRevX.3.041003

- [11] Goold J, Huber M, Riera A, del Rio L and Skrzypczyk P 2016 Journal of Physics A: Mathematical and Theoretical 49 143001 URL http://stacks.iop.org/1751-8121/49/i=14/ a=143001
- [12] Deffner S and Campbell S 2019 Quantum Thermodynamics (Morgan & Claypool Publishers) ISBN 978-1-64327-658-8 URL http://dx.doi.org/10.1088/2053-2571/ab21c6
- [13] Dowling J P and Milburn G J 2003 Philosophical Transactions of the Royal Society of London. Series A: Mathematical, Physical and Engineering Sciences 361 1655-1674 URL https://royalsocietypublishing.org/doi/abs/10.1098/rsta.2003.1227
- [14] Deutsch I H 2020 PRX Quantum 1(2) 020101 URL https://link.aps.org/doi/10.1103/ PRXQuantum.1.020101
- [15] Tuncer A and Müstecaphoğlu O E 2020 Turk. J. Phys. 44(5) 404 URL https://journals. tubitak.gov.tr/physics/abstract.htm?id=28103
- Bhattacharjee S and Dutta A 2020 arXiv preprint arXiv:2008.07889 URL https://arxiv.org/ abs/2008.07889
- [17] Mitchison M T 2019 Contemporary Physics 60 164-187 URL https://doi.org/10.1080/ 00107514.2019.1631555
- [18] Latune C, Sinayskiy I and Petruccione F 2020 arXiv preprint arXiv:2006.01166 URL https: //arxiv.org/abs/2006.01166
- [19] Mukherjee V and Divakaran U 2021 Many-body quantum technologies URL https://arxiv. org/abs/2102.08301
- [20] Ghosh A, Mukherjee V, Niedenzu W and Kurizki G 2019 The European Physical Journal Special Topics 227 2043-2051 ISSN 1951-6401 URL https://doi.org/10.1140/epjst/ e2019-800060-7
- [21] Quan H T, Wang Y D, Liu Y x, Sun C P and Nori F 2006 Phys. Rev. Lett. 97(18) 180402 URL https://link.aps.org/doi/10.1103/PhysRevLett.97.180402
- [22] Definer S 2013 Phys. Rev. E 88 062128 URL https://link.aps.org/doi/10.1103/PhysRevE. 88.062128
- [23] Strasberg P, Schaller G, Brandes T and Esposito M 2017 Phys. Rev. X 7 021003 URL https://link.aps.org/doi/10.1103/PhysRevX.7.021003
- [24] Stevens J and Definer S 2019 Phys. Rev. E 99 042129 URL https://link.aps.org/doi/10. 1103/PhysRevE.99.042129
- [25] Ashrafi M, Anza F and Crutchfield J P 2020 arXiv preprint arXiv:2010.14652 URL https: //arxiv.org/abs/2010.14652
- [26] Nielsen M A and Chuang I 2002 Quantum computation and quantum information (American Association of Physics Teachers)
- [27] Polkovnikov A 2011 Ann. Phys. 326 486 499 ISSN 0003-4916 URL https://doi.org/10.1016/ j.aop.2010.08.004
- [28] Deffner S 2020 Phys. Rev. Research 2 013161 URL https://link.aps.org/doi/10.1103/ PhysRevResearch.2.013161
- [29] Definer S 2021 arXiv preprint arXiv:2102.05118 URL https://arxiv.org/abs/2102.05118
- [30] Chapman A and Miyake A 2015 Phys. Rev. E 92(6) 062125 URL https://link.aps.org/doi/ 10.1103/PhysRevE.92.062125
- [31] Šafránek D and Deffner S 2018 Phys. Rev. A 98(3) 032308 URL https://link.aps.org/doi/ 10.1103/PhysRevA.98.032308
- [32] Dağ C B, Niedenzu W, Müstecaphoğlu Ö E and Kurizki G 2016 Entropy 18 244 ISSN 1099-4300 URL https://www.mdpi.com/1099-4300/18/7/244
- [33] Francica G, Binder F C, Guarnieri G, Mitchison M T, Goold J and Plastina F 2020 Phys. Rev. Lett. 125(18) 180603 URL https://link.aps.org/doi/10.1103/PhysRevLett.125.180603
- [34] Korzekwa K, Lostaglio M, Oppenheim J and Jennings D 2016 New Journal of Physics 18 023045 URL https://doi.org/10.1088%2F1367-2630%2F18%2F2%2F023045
- [35] Çakmak B 2020 Phys. Rev. E 102(4) 042111 URL https://link.aps.org/doi/10.1103/ PhysRevE.102.042111
- [36] Guarnieri G, Morrone D, Çakmak B, Plastina F and Campbell S 2020 Physics Letters A 126576 ISSN 0375-9601 URL http://www.sciencedirect.com/science/article/pii/ S0375960120304436
- [37] Funo K, Watanabe Y and Ueda M 2013 Phys. Rev. A 88(5) 052319 URL https://link.aps. org/doi/10.1103/PhysRevA.88.052319
- [38] Francica G, Goold J, Plastina F and Paternostro M 2017 npj Quantum Information 3(1) 12 URL https://doi.org/10.1038/s41534-017-0012-8
- [39] Bernards F, Kleinmann M, Gühne O and Paternostro M 2019 Entropy 21 ISSN 1099-4300 URL https://www.mdpi.com/1099-4300/21/8/771

- [40] Manzano G, Plastina F and Zambrini R 2018 Phys. Rev. Lett. 121(12) 120602 URL https: //link.aps.org/doi/10.1103/PhysRevLett.121.120602
- [41] Perarnau-Llobet M, Hovhannisyan K V, Huber M, Skrzypczyk P, Brunner N and Acín A 2015 Phys. Rev. X 5(4) 041011 URL https://link.aps.org/doi/10.1103/PhysRevX.5.041011
- [42] Fusco L, Paternostro M and De Chiara G 2016 Phys. Rev. E 94(5) 052122 URL https: //link.aps.org/doi/10.1103/PhysRevE.94.052122
- [43] Sapienza F, Cerisola F and Roncaglia A J 2019 Nature communications 10 1–7 URL https: //doi.org/10.1038/s41467-019-10572-8
- [44] Oppenheim J, Horodecki M, Horodecki P and Horodecki R 2002 Phys. Rev. Lett. 89(18) 180402 URL https://link.aps.org/doi/10.1103/PhysRevLett.89.180402
- [45] Jevtic S, Jennings D and Rudolph T 2012 Phys. Rev. Lett. 108(11) 110403 URL https: //link.aps.org/doi/10.1103/PhysRevLett.108.110403
- [46] Niedenzu W, Huber M and Boukobza E 2019 Quantum 3 195 ISSN 2521-327X URL https: //doi.org/10.22331/q-2019-10-14-195
- [47] Allahverdyan A E, Balian R and Nieuwenhuizen T M 2004 Europhysics Letters (EPL) 67 565– 571 URL https://doi.org/10.1209%2Fepl%2Fi2004-10101-2
- [48] Pusz W and Woronowicz S L 1978 Communications in Mathematical Physics 58 273-290 URL https://doi.org/10.1007/BF01614224
- [49] Bergmann P G and Lebowitz J L 1955 Phys. Rev. 99(2) 578-587 URL https://link.aps.org/ doi/10.1103/PhysRev.99.578
- [50] Esposito M and den Broeck C V 2011 EPL (Europhysics Letters) 95 40004 URL https: //doi.org/10.1209/0295-5075/95/40004
- [51] Parrondo J M R, Horowitz J M and Sagawa T 2015 Nature Physics 11 131–139 ISSN 1745-2481 URL https://doi.org/10.1038/nphys3230
- [52] Watanabe S 1960 IBM Journal of Research and Development 4 66-82 URL https://doi.org/ 10.1147/rd.41.0066
- [53] Henderson L and Vedral V 2001 Journal of Physics A: Mathematical and General 34 6899-6905 URL https://doi.org/10.1088/0305-4470/34/35/315
- [54] Ollivier H and Zurek W H 2001 Phys. Rev. Lett. 88(1) 017901 URL https://link.aps.org/ doi/10.1103/PhysRevLett.88.017901
- [55] Touil A and Definer S 2021 PRX Quantum 2 010306 URL https://doi.org/10.1103/ PRXQuantum.2.010306
- [56] Bernards F, Kleinmann M, Gühne O and Paternostro M 2019 Entropy 21 771 URL https: //www.mdpi.com/1099-4300/21/8/771
- [57] Manzano G, Plastina F and Zambrini R 2018 Phys. Rev. Lett. 121(12) 120602 URL https: //link.aps.org/doi/10.1103/PhysRevLett.121.120602
- [58] Campisi M and Goold J 2017 Phys. Rev. E 95(6) 062127 URL https://link.aps.org/doi/10. 1103/PhysRevE.95.062127
- [59] Yan B, Cincio L and Zurek W H 2020 Phys. Rev. Lett. 124(16) 160603 URL https://link. aps.org/doi/10.1103/PhysRevLett.124.160603
- [60] Nakamura S, Iyoda E, Deguchi T and Sagawa T 2019 Phys. Rev. B 99(22) 224305 URL https://link.aps.org/doi/10.1103/PhysRevB.99.224305
- [61] Iyoda E and Sagawa T 2018 Phys. Rev. A 97(4) 042330 URL https://link.aps.org/doi/10. 1103/PhysRevA.97.042330
- [62] Touil A and Definer S 2020 Quantum Science and Technology 5 035005 URL https://doi.org/ 10.1088%2F2058-9565%2Fab8ebb
- [63] Zanardi P and Anand N 2020 arXiv preprint arXiv:2012.13172 URL https://arxiv.org/abs/ 2012.13172
- [64] Rossini D, Andolina G M, Rosa D, Carrega M and Polini M 2020 Phys. Rev. Lett. 125(23) 236402 URL https://link.aps.org/doi/10.1103/PhysRevLett.125.236402
- [65] Rosa D, Rossini D, Andolina G M, Polini M and Carrega M 2020 Journal of High Energy Physics 2020 67 ISSN 1029-8479 URL https://doi.org/10.1007/JHEP11(2020)067
- [66] Quesada N, Al-Qasimi A and James D F 2012 Journal of Modern Optics 59 1322–1329 URL https://doi.org/10.1080/09500340.2012.713130
- [67] Cai J M, Zhou Z W and Guo G C 2005 Phys. Rev. A 72(2) 022312 URL https://link.aps. org/doi/10.1103/PhysRevA.72.022312
- [68] Carnio E G, Buchleitner A and Gessner M 2016 New Journal of Physics 18 073010 URL https://doi.org/10.1088/1367-2630/18/7/073010
- [69] Amico L, Fazio R, Osterloh A and Vedral V 2008 Rev. Mod. Phys. 80(2) 517-576 URL https://link.aps.org/doi/10.1103/RevModPhys.80.517
- [70] Sarandy M S, De Oliveira T R and Amico L 2013 International Journal of Modern Physics B

27 1345030 URL https://doi.org/10.1142/S0217979213450306

- [71] Çakmak B, Karpat G and Fanchini F F 2015 Entropy 17 790-817 ISSN 1099-4300 URL https://www.mdpi.com/1099-4300/17/2/790
- [72] Çakmak B, Manatuly A and Müstecaplıoğlu O E 2017 Phys. Rev. A 96(3) 032117 URL https://link.aps.org/doi/10.1103/PhysRevA.96.032117
- [73] Latune C L, Sinayskiy I and Petruccione F 2019 Phys. Rev. A 99(5) 052105 URL https: //link.aps.org/doi/10.1103/PhysRevA.99.052105
- [74] Gross M and Haroche S 1982 Physics Reports 93 301 396 ISSN 0370-1573 URL http: //www.sciencedirect.com/science/article/pii/0370157382901028
- [75] Stephen M J 1964 The Journal of Chemical Physics 40 669-673 URL https://doi.org/10. 1063/1.1725188
- [76] Lehmberg R H 1970 Phys. Rev. A 2(3) 883-888 URL https://link.aps.org/doi/10.1103/ PhysRevA.2.883
- [77] Damanet F and Martin J 2016 Journal of Physics B: Atomic, Molecular and Optical Physics 49 225501 URL https://doi.org/10.1088%2F0953-4075%2F49%2F22%2F225501
- [78] Definer S and Campbell S 2017 Journal of Physics A: Mathematical and Theoretical 50 453001 URL https://doi.org/10.1088/1751-8121/aa86c6
- [79] Definer S and Lutz E 2013 Journal of Physics A: Mathematical and Theoretical 46 335302 URL https://doi.org/10.1088/1751-8113/46/33/335302
- [80] Kakutani S 1948 Annals of Mathematics 214-224 URL https://www.jstor.org/stable/ 1969123?seq=1
- [81] Bures D 1969 Transactions of the American Mathematical Society 135 199-212 URL https: //www.jstor.org/stable/1995012?seq=1
- [82] Definer S 2017 New Journal of Physics 19 103018 URL https://doi.org/10.1088/1367-2630/ aa83dc
- [83] O'Connor E, Guarnieri G and Campbell S 2021 Phys. Rev. A 103(2) 022210 URL https: //link.aps.org/doi/10.1103/PhysRevA.103.022210
- [84] Allan D, Hörnedal N and Andersson O 2020 arXiv:2011.11963 URL https://arxiv.org/abs/ 2011.11963
- [85] Wakamura H and Koike T 2020 New Journal of Physics URL https://iopscience.iop.org/ article/10.1088/1367-2630/ab8ab3/meta
- [86] Niedenzu W, Mukherjee V, Ghosh A, Kofman A G and Kurizki G 2018 Nature Communications 9 165 URL https://doi.org/10.1038/s41467-017-01991-6