

Creative Commons Attribution 4.0 International (CC BY 4.0)

<https://creativecommons.org/licenses/by/4.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Temporal Changes in Epinephrine Dosing in Out-of-Hospital Cardiac Arrest: A Review of EMS Protocols across the United States

Eric Garfinkel, DO;¹  Katelyn Michelsen, MD;² Benjamin Johnson, MSc;¹ Asa Margolis, DO, MS, MPH;¹ Matthew Levy, DO, MSc¹ 

1. Department of Emergency Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland USA
2. SSM Health St. Anthony Hospital, Oklahoma City, Oklahoma USA

Correspondence:

Eric Garfinkel, DO
Johns Hopkins Hospital Emergency
Department
1800 Orleans St.
Baltimore, Maryland 21287 USA
E-mail: egarfin2@jhu.edu

Conflicts of interest: The author(s) declare none.

Keywords: ACLS; cardiac arrest; EMS; epinephrine; resuscitation

Abbreviations:

ACLS: Advanced Cardiac Life Support
CPR: cardiopulmonary resuscitation
EMS: Emergency Medical Services
ROSC: return of spontaneous circulation

Received: July 21, 2022

Revised: August 17, 2022

Accepted: September 7, 2022

doi:[10.1017/S1049023X22001418](https://doi.org/10.1017/S1049023X22001418)

© The Author(s), 2022. Published by Cambridge University Press on behalf of the World Association for Disaster and Emergency Medicine. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

Background: Administration of epinephrine has been associated with worse neurological outcomes for survivors of out-of-hospital cardiac arrest. The publication of the 2018 PARAMEDIC-2 trial, a randomized and double-blind study of epinephrine in out-of-hospital cardiac arrest, provides the strongest evidence to date that epinephrine increases return of spontaneous circulation (ROSC) but not neurologically intact survival. This study aims to determine if Emergency Medical Services (EMS) cardiac arrest protocols have changed since the publication of PARAMEDIC-2.

Methods: States in the US utilizing mandatory or model state-wide EMS protocols, including Washington DC, were included in this study. The nontraumatic cardiac arrest protocol as of January 1, 2018 was compared to the protocol in effect on January 1, 2021 to determine if there was a change in the administration of epinephrine. Protocols were downloaded from the relevant state EMS website. If a protocol could not be obtained, the state medical director was contacted.

Results: A 2021 state-wide protocol was found for 32/51 (62.7%) states. Data from 2018 were available for 21/51 (41.2%) states. Of the 11 states without data from 2018, all follow Advanced Cardiac Life Support (ACLS) guidelines in the 2021 protocol. Five (15.6%) of the states with a state-wide protocol made a change in the cardiac arrest protocols. Maximum cumulative epinephrine dose was limited to 4mg in Maryland and 3mg in Vermont. Rhode Island changed epinephrine in shockable rhythms to be administered after three cycles of cardiopulmonary resuscitation (CPR) and an anti-arrhythmic. Rhode Island also added an epinephrine infusion as an option. No states removed epinephrine administration from their cardiac arrest protocol. Simple statistical analysis was performed with Microsoft Excel.

Conclusion: Several states have adjusted cardiac arrest protocols since 2018. The most frequent change was limiting the maximum cumulative dosage of epinephrine. One state changed timing of epinephrine dosing depending on the rhythm and also provided an option of an epinephrine infusion in place of bolus dosing. While the sample size is small, these changes may reflect the future direction of prehospital cardiac arrest protocols. Significant limitations apply, including the exclusion of local and regional protocols which are more capable of quickly adjusting to new research. Additionally, this study is only focused on EMS in the United States.

Garfinkel E, Michelsen K, Johnson B, Margolis A, Levy M. Temporal changes in epinephrine dosing in out-of-hospital cardiac arrest: a review of EMS protocols across the United States. *Prehosp Disaster Med.* 2022;37(6):832–835.

Introduction

The administration of epinephrine has been a cornerstone of cardiac arrest resuscitation for decades.¹ Stimulation of alpha receptors increases myocardial and cerebral blood flow and leads to increased rates of return of spontaneous circulation (ROSC), while beta stimulation increases myocardial oxygen demand and arrhythmogenicity.^{2,3} Since 1974, the American Heart Association (AHA; Dallas, Texas USA) has published Advanced Cardiac Life Support (ACLS) guidelines on cardiac arrest. While a variety of adrenergic and antiarrhythmic medications have come and gone, epinephrine remains the only drug that continues to be recommended for all nontraumatic cardiac arrests.



Evidence for the use of epinephrine in cardiac arrest continues to evolve, and while recent studies continue to affirm increased rates of ROSC associated with epinephrine use, there is variability in the most critical outcome: neurologically intact survival.⁴⁻⁷ The 2018 PARAMEDIC-2 trial is the largest trial of epinephrine in out-of-hospital cardiac arrest and demonstrated an increase in the primary outcome of survival at 30 days (3.2% versus 2.4%) but no statistically significant difference in survival to hospital discharge with a favorable neurologic status.

Treating a patient in cardiac arrest is a fundamental role of Emergency Medical Services (EMS). It is essential for EMS systems to ensure that treatment protocols and resuscitation practices are informed by scientific evidence and include new advances in the understanding of resuscitative medicine. This project sought to describe the current state of administration of epinephrine within prehospital cardiac arrest protocols across the United States.

Methods

An internet search engine was used during the period of July 1, 2021 through December 31, 2021 to access publicly available state EMS agency websites in all 50 US States and Washington, DC. The EMS treatment protocols in place as of January 1, 2018 were compared to the protocols in place as of January 1, 2021. Any changes in epinephrine administration in cardiac arrest management protocols were recorded including dosage, frequency, and difference between shockable or non-shockable rhythm. For any states in which the 2021 protocol was not found online, the state medical director's office was contacted for further information. Information for states unobtainable despite these efforts, as well as those states that do not have state-wide protocols, were excluded. Summary and descriptive statistics were performed in Microsoft Excel (Microsoft Corp.; Redmond, Washington USA). Two reviewers, KM and BJ, verified the protocols independently. If there was an inconsistency, a tie-breaking decision was made by EG. This project was reviewed and approved by the Johns Hopkins School of Medicine (Baltimore, Maryland USA) Institutional Review Board, protocol number 00298128.

Results

Of the 50 states and the District of Columbia, 21 (41.2%) yielded complete data from both 2018 and 2021. In 11 states (21.6%), the 2018 protocols were unable to be accessed. However, all 11 states had 2021 protocols consistent with current ACLS guidelines and thus it was assumed that there was no change from 2018. A further 19 (37.3%) states were confirmed to have no state-wide protocols. Table 1 lists which states utilized state-wide protocols and the data availability.

Of the 32 (62.7%) states with state-wide protocols, five (15.6%) recorded a change between 2018 and 2021 in epinephrine administration during cardiac arrest and 27 (84.4%) had no change. The five states with recorded changes were Arizona, Maryland, Rhode Island, Utah, and Vermont (Figure 1). The differences are listed in Table 2. Arizona and Utah have model protocols to guide local protocols, while Maryland, Rhode Island, and Vermont have mandatory state-wide EMS protocols.

Arizona limits epinephrine to a maximum of three total doses. Maryland limits epinephrine to a maximum of four doses of epinephrine, plus an additional two doses if the patient rearrests following ROSC. Utah recommends considering limiting

States with State-Wide Protocols and Complete Data	States with State-Wide Protocols Missing 2018 Data	States with Local or Regional Protocols
Alabama	Georgia	Alaska
Arizona	Idaho	California
Arkansas	Maine	Colorado
Connecticut	Massachusetts	Florida
Delaware	Montana	Illinois
Hawaii	New Jersey	Indiana
Iowa	New Mexico	Kansas
Kentucky	North Carolina	Louisiana
Maryland	Ohio	Minnesota
Michigan	Oklahoma	Mississippi
Nebraska	West Virginia	Missouri
New Hampshire		Nevada
New York		Oregon
North Dakota		South Dakota
Pennsylvania		Texas
Rhode Island		Virginia
South Carolina		Washington
Tennessee		Wisconsin
Utah		Wyoming
Vermont		
Washington DC (District)		
21 Total	11 Total	19 Total

Garfinkel © 2022 Prehospital and Disaster Medicine

Table 1. Protocol Data Availability for Each State and the District of Columbia

epinephrine to three doses unless there is a response. Vermont limits epinephrine to three doses.

Rhode Island changed the protocol from administering epinephrine every three-to-five minutes in all rhythms to administering epinephrine after three cycles of cardiopulmonary resuscitation (CPR), electrical, therapy, and one dose of antiarrhythmic in shockable rhythms. In addition, Rhode Island is the only state which added an epinephrine infusion as an option in place of frequent boluses.


In all state protocols in force at January 1, 2018, except for New Jersey, epinephrine dosing and frequency was consistent with the ACLS guidelines. New Jersey limited epinephrine to a maximum of three doses. No states added or eliminated epinephrine from their protocol during the studied period.

Discussion

This study showed that only a limited number of states changed protocols to reflect the recent literature regarding epinephrine in cardiac arrest. Amongst these states, there appears to be an emerging trend towards limiting epinephrine dosage. The most frequent dose limitation was to a total of three milligrams, except for Maryland which limits the dosage to four milligrams. No study has conclusively looked at the ideal amount of epinephrine to be administered during cardiac arrest and international guidelines do not recommend a maximum epinephrine dose.⁸ Fothergill, et al demonstrated a significant drop in the adjusted odds ratio for survival to hospital discharge once the third dose of epinephrine

State	System Type	2018 Protocol	2021 Changes
Arizona	Model	ACLS	Maximum three doses of epinephrine
Maryland	Mandatory	ACLS	Maximum four doses of epinephrine, two more if rearrest
Rhode Island	Mandatory	ACLS	For shockable rhythms, administer epinephrine after three cycles of CPR, defibrillation, and an antiarrhythmic
			Consider epinephrine infusion in place of bolus
Utah	Model	ACLS	Recommends maximum three doses of epinephrine, unless a clear response is observed
Vermont	Mandatory	ACLS	Maximum three doses of epinephrine

Garfinkel © 2022 Prehospital and Disaster Medicine
Table 2. States with Changes in Epinephrine Use in Out-of-Hospital Cardiac Arrest from 2018 through 2021
Note: ACLS protocol is consistent with the American Heart Association’s Advanced Cardiac Life Support algorithm.
Abbreviations: ACLS, Advanced Cardiac Life Support; CPR, cardiopulmonary resuscitation.



Garfinkel © 2022 Prehospital and Disaster Medicine
Figure 1. Map of States with State-Wide Protocols which had Changes in Protocol during the Study Period.
Abbreviation: EMS, Emergency Medical Services.

was administered, from 0.7 for two doses of epinephrine to 0.15 for three or greater doses.⁹ Higher epinephrine dosing is associated with a longer resuscitation period, which is clearly associated with a worse outcome, thus identifying the optimal dosing of epinephrine is a challenge. The Fothergill study suggests that a cut off of 3mg is reasonable, however, further research is required.

The early administration of epinephrine is associated with higher rates of ROSC for all forms of cardiac arrest.^{10,11} Both ACLS and the majority of the state protocols reviewed recommend defibrillation as soon as possible and epinephrine delivery after the second shock, followed by amiodarone or lidocaine. Rhode Island changed its protocol to administer epinephrine in shockable rhythms after three cycles of CPR, defibrillation, and an antiarrhythmic. Epinephrine administration after an antiarrhythmic

drug is unique to the states surveyed and of unclear significance. Although less robustly studied, amiodarone has also been associated with increased survival with earlier administration and thus this is an area of clinical equipoise.¹²

Rhode Island allows for an epinephrine infusion in place of epinephrine bolus dosing. Studies to support this are lacking, but there are several theoretical benefits such as simplifying the resuscitation process and producing a more consistent serum epinephrine level. In the case of ROSC, the epinephrine infusion is already available and can be rapidly titrated to avoid hypotension. The downside is potentially increased dosing errors and time commitment to mix the infusion.

Studies have suggested that it takes on average 17 years for basic research to change clinical practice.¹³ It is thus not surprising that

only a limited number of states have changed epinephrine use in nontraumatic out-of-hospital cardiac arrest. The changes in these states, however, may reflect the future of cardiac arrest resuscitation and provide an important framework for future research.

Limitations

This study focused on prehospital cardiac arrest resuscitation in the United States only, so it may not be generalizable to other countries. Additionally, a further limitation was the exclusion of states that did not have state-wide protocols. The states that were included did have approximately 49.4% of the 2020 population of the United States, which suggests that the studied states reflected a significant portion of the country.¹⁴ While inclusion of local and regional protocols would have allowed for a better understanding of the changes that have been made, it would have also complicated the generalizability of these findings, as jurisdictions of smaller size are able to adapt to research findings at a faster rate than that of protocols that are dictated on a state-wide level.

Additionally, the authors were unable to find the 2018 protocols for several states. Since the 2021 protocols were consistent with ACLS, it was assumed that there was no change. However, it is possible that the states with missing data could have made changes between 2018 and 2021 to become congruent with ACLS guidelines. Finally, the limited number of states that instituted a change in their protocol limits drawing any definitive conclusions but does suggest potential future directions.

Conclusion

Five states have changed their cardiac arrest protocols to alter epinephrine administration from 2018 through 2021. The most frequent change was limiting the total number of epinephrine administered to either three or four milligrams. This may represent the future direction of epinephrine use for out-of-hospital nontraumatic cardiac arrest, however, conclusions are limited by a small sample size and focus on a single country's EMS system.

References

1. Paradis NA, Koscove EM. Epinephrine in cardiac arrest: a critical review. *Ann Emerg Med.* 1990;19(11):1288–1301.
2. Yakaitis RW, Otto CW, Blitt CD. Relative importance of α and β adrenergic receptors during resuscitation. *Crit Care Med.* 1979;7(7):293–296.
3. Ditchey RV, Lindenfeld J. Failure of epinephrine to improve the balance between myocardial oxygen supply and demand during closed-chest resuscitation in dogs. *Circulation.* 1988;78(2):382–389.
4. Lin S, Callaway CW, Shah PS, et al. Adrenaline for out-of-hospital cardiac arrest resuscitation: a systematic review and meta-analysis of randomized controlled trials. *Resuscitation.* 2014;85(6):732–740.
5. Loomba RS, Nijhawan K, Aggarwal S, et al. Increased return of spontaneous circulation at the expense of neurologic outcomes: is prehospital epinephrine for out-of-hospital cardiac arrest really worth it? *J Crit Care.* 2015;30(6):1376–1381.
6. Nakahara S, Tomio J, Takahashi H, et al. Evaluation of pre-hospital administration of adrenaline (epinephrine) by emergency medical services for patients with out of hospital cardiac arrest in Japan: controlled propensity matched retrospective cohort study. *BMJ.* 2013;347:f6829.
7. Perkins GD, Ji C, Deakin CD, et al. A randomized trial of epinephrine in out-of-hospital cardiac arrest. *N Engl J Med.* 2018;379(8):711–721.
8. Merchant RM, Topjian AA, Panchal AR, et al. Part 1: Executive Summary: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation.* 2020;142(16_suppl_2):S337–S357.
9. Fothergill RT, Emmerson AC, Iyer R, et al. Repeated adrenaline doses and survival from an out-of-hospital cardiac arrest. *Resuscitation.* 2019;138:316–321.
10. Holmberg MJ, Issa MS, Moskowitz A, et al. Vasopressors during adult cardiac arrest: a systematic review and meta-analysis. *Resuscitation.* 2019;139:106–121.
11. Okubo M, Komukai S, Callaway CW, et al. Association of timing of epinephrine administration with outcomes in adults with out-of-hospital cardiac arrest. *JAMA Netw Open.* 2021;4(8):e2120176.
12. Wissa J, Schultz BV, Wilson D, et al. Time to amiodarone administration and survival outcomes in refractory ventricular fibrillation. *Emerg Med Australas.* 2021;33(6):1088–1094.
13. Morris ZS, Wooding S, Grant J. The answer is 17 years, what is the question: understanding time lags in translational research. *J R Soc Med.* 2011;104(12):510–520.
14. 2020 Population and Housing State Data. United States Census Bureau. <https://www.census.gov/library/visualizations/interactive/2020-population-and-housing-state-data.html>. Accessed July 13, 2020.