

This item is likely protected under Title 17 of the U.S. Copyright Law. Unless on a Creative Commons license, for uses protected by Copyright Law, contact the copyright holder or the author.

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.



NEWS

Community

Policy & Society

Science & Tech

Arts & Culture

Science & Tech

New UMBC/Los Alamos research on megafire smoke plumes clarifies what they contain, how they move, and their potential impacts

Published: Apr 21, 2023



By: [Adriana Fraser](#)



Active flame front of the Zaca fire, the single largest wildfire of the 2007 California wildfire season. (John Newman of the U.S. Forest Service/Wikimedia Commons)

In recent years, large, intense wildfires, known as megafires, have increasingly caused severe damage to forests, homes, and crops. In addition to megafires fatally impacting humans and wildlife alike, they may also be impacting climate change. New research led by UMBC's **Stephen Guimond** provides insight into how the large smoke plumes produced by megafires can be more accurately modeled and characterized to improve our understanding of how they might impact the earth.

Guimond, an associate research professor of physics, collaborated with scientists at the [Los Alamos National Laboratory](#) to determine the long-term effects of smoke plumes from megafires. Their findings, recently published in the [Journal of Advances in Modeling Earth Systems](#), demonstrates how previous research utilized a model grid spacing that does not sample smoke plumes accurately. These

inaccuracies in defining the dynamics of the problem lead to errors in interpretation of the smoke's properties, vertical and horizontal movement of the plume, and potential climatic effects.

Tracking how smoke rises

The smoke plumes from megafires are voluminous and can rise very high into the upper atmosphere. Initially, the plumes get transported upwards by convective cells and travel into the stratosphere, explains Guimond, who is also a scientist at UMBC's [Goddard Earth Sciences Technology and Research \(GESTAR\) II](#) (previously known as the Joint Center for Earth Systems Technology).

"Once it gets up into the stratosphere, the smoke can stay around for many months, even up to a year or more. The fact that it can stay up there so long means that you can get effects on the solar radiation reaching the surface," says Guimond. "If you have a big, dark-colored blanket of smoke up there, it's going to absorb most of the sunlight, which will lead to less sunlight reaching the surface of the earth. Because of this, you could get, over a long period of time, a cooling that happens on the surface of the earth," among other impacts.

For three years, scientists at Los Alamos studied the chemical properties of smoke plumes by burning objects like trees in a controlled setting to determine the percentages of carbon that the smoke emitted. The scientists evaluated atmospheric particulates, or aerosols, which have a major effect on climate. Guimond used a NASA climate model to determine the carbon characteristics of the smoke plumes, how they rise into the atmosphere, and the underlying causes of rotation within the plumes.

"The measurements we looked at included particle types, the spectrum of the particles, and their sizes," says Guimond. "We also looked at the contributions of



Stephen Guimond. (Photo courtesy of Stephen Guimond)

different chemical species such as black carbon, organic carbon, and other chemical compounds that come off of burning materials.”

Assessing previous smoke plume research

The color of the smoke is an important factor, Guimond notes, as different types of smoke have different radiative properties. White smoke is composed mostly of organic carbon: brightly-colored aerosol particles that in large part reflect radiation back into the atmosphere. Black smoke is composed mostly of black carbon: dark-colored aerosol particles that absorb radiation.

The researchers determined that previous models didn't accurately sample the types of carbon within smoke plumes, leading to miscalculations or incorrect assumptions about the percentage of black carbon the plumes contained.

As the black smoke absorbs solar radiation, the smoke heats up, which can create a lofting effect that pushes the smoke higher into the atmosphere. The higher the smoke rises, the longer it stays in the stratosphere. The longer it stays, the more time it has to impact the surface of the earth. This means that inaccurate characterization of the percentage of black carbon in wildfire smoke can lead to inaccurate calculations of the lofting effect, height of the plume and stratospheric lifetime, as well as climatic effects.




Figure 1 (left): The horizontal structure of the smoke plume at 6.2 days by vertically integrating the total smoke mixing ratio over the entire model atmosphere into the simulations for varying resolutions: (a) 2.0° (b) 1.0° (c) 0.25° and (d) 7 km.

Figure 2 (right): The same as figure 1, except at 16.2 days. The horizontal structure of the smoke plume is significantly different between the various resolution simulations. (Images courtesy of Stephen Guimond and the *Journal of Advances in Modeling Earth Systems*)

Limits in how previous research represented the atmosphere also made for less accurate smoke plume simulations, Guimond said. He notes that prior smoke plume research used “coarse representation of the smoke plume in the model calculations, which has significant downstream effects on all other components of this problem, including the conclusions drawn from the research.”

Guimond hopes that his research can improve understanding of the dynamics of this problem: tracking of atmospheric motion and forces in three-dimensions, and how phenomena like rotating smoke plumes form and decay.

“Scientists need to accurately simulate the dynamics in order to get more accurate answers about aerosol properties inside the smoke plumes,” Guimond says, “such as how much of the smoke is black carbon, how long it is going to last in the stratosphere, how high it rises, and its effects on the radiation of the earth.”

With more accurate models and simulations, future research will be able to better inform climate policy and megafire response.

Tags: [CNMS](#), [GESTAR](#), [JCET](#), [Physics](#), [rca-2](#), [Research](#)

Share this Story

Related Posts

Class of 2023

Class of 2023



Leading through service: Meet the community builders in UMBC's Class of 2023

Community



UMBC's vibrant learning community helps students discover careers to fit their passions



UMBC softball nabs 4th consecutive America East title, advances to NCAA tournament

Want more UMBC news?

Get top stories delivered to your inbox.

Sign up for our weekly UMBC Top Stories email:

Join our list

Contact Us



Share a story idea and learn more about the news team.

More about UMBC News



University of Maryland, Baltimore County
1000 Hilltop Circle, Baltimore, MD 21250

Directions & Parking Information



Resources

[Alumni](#)

[Career Center](#)

[Events](#)

[Get Help](#)

[News](#)

[Visit Campus](#)

[Work at UMBC](#)

Important Contacts

[Online Directory](#)

[Contact UMBC](#)

[Request Support](#)

Emergency Info

UMBC Police: 410-455-5555

[Sign Up for Text Alerts](#)

REQUEST INFO

APPLY

Top Stories of the Week

Enter email to subscribe

Email

GO

© UMBC: A University System of Maryland
Member Institution

[Accreditation](#)

[Equal Opportunity](#)

[Privacy](#)

[Title IX](#)

[Web Accessibility](#)