




This work is on a Creative Commons Attribution 4.0 International (CC BY 4.0) license, <https://creativecommons.org/licenses/by/4.0/>. Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

Article

Quantum Euler Relation for Local Measurements

Akram Touil ^{1,*} , Kevin Weber ^{1,2}  and Sebastian Deffner ^{1,3} 
¹ Department of Physics, University of Maryland, Baltimore County, Baltimore, MD 21250, USA; kweber12@umd.edu (K.W.); deffner@umbc.edu (S.D.)

² Institute for Physical Science and Technology, University of Maryland, College Park, MD 20742, USA

³ Instituto de Física ‘Gleb Wataghin’, Universidade Estadual de Campinas, Campinas 13083-859, SP, Brazil

* Correspondence: akramt1@umbc.edu

Abstract: In classical thermodynamics the Euler relation is an expression for the internal energy as a sum of the products of canonical pairs of extensive and intensive variables. For quantum systems the situation is more intricate, since one has to account for the effects of the measurement back action. To this end, we derive a quantum analog of the Euler relation, which is governed by the information retrieved by local quantum measurements. The validity of the relation is demonstrated for the collective dissipation model, where we find that thermodynamic behavior is exhibited in the weak-coupling regime.

Keywords: information gain; holevo bound; quantum discord; local measurements; ergotropy



Citation: Touil, A.; Weber, K.; Deffner, S. Quantum Euler Relation for Local Measurements. *Entropy* **2021**, *23*, 889. <https://doi.org/10.3390/e23070889>

Academic Editor: Rosario Lo Franco

Received: 11 June 2021

Accepted: 8 July 2021

Published: 13 July 2021

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Thermodynamics is a phenomenological theory studying the average behavior of heat and work [1,2] originally developed to optimize heat engines [3]. Remarkably, this theory has been extended far beyond its origins and led to profound fundamental statements about nature, such as the second law linking the monotonicity of entropy to the arrow of time [2], and the seminal role of information in physics [4,5]. Building on the success of this theory, quantum thermodynamics [6,7] emerged as generalization of stochastic thermodynamics [6,8–10] to the quantum realm.

Quantum thermodynamics aims to ground our understanding of the universal laws and statements on a fundamental and genuine quantum level [6], which is set to have a direct impact on the development of new generation quantum technologies [11,12]. Therefore, many recent studies focused on probing the role of quantum information as a thermodynamic resource [13–26], and deriving second law statements for quantum systems [6,26,27]. However, to the best of our knowledge, fundamental statements such as quantum generalizations of the Euler relation are still lacking.

In classical thermodynamics, the Euler relation is a fundamental statement expressing the internal energy U as a sum of the products of canonical pairs of extensive X_i and intensive Y_i variables [1],

$$U = \sum_i Y_i X_i. \quad (1)$$

In its simplest form, this fundamental statement relates the internal energy and the Clausius entropy S , with the volume V and the particle number N , such that $U = TS - PV + \mu N$, and the intensive variables are the temperature T , pressure P , and chemical potential μ . This relation is derived using Euler’s homogeneous function theorem [1], hence the name “Euler relation”. The main goal of the present analysis is to derive a quantum analog of the Euler relation, that we dub the quantum Euler relation. To this end, we have to relate the internal energy to thermodynamic quantities as well as the information output of quantum measurements. Consequently, our new relation connects the information gained from quantum measurements to inherent thermodynamic properties of a system.

In the following, we focus on an established measure that quantifies the output information of a quantum measurement and its effects on the system we are probing. This measure was first introduced in the seventies [28], when Groenewold proposed entropy reduction as a straightforward quantifier of information gain. Hereafter, we will simply refer to this quantifier as the information gain “ \mathcal{I}_g ”. Note that the information gain is generally different from the Holevo information or the Holevo bound “ χ ” [29,30], which is widely used in the literature to quantify the classical information gained from measurements of bipartite quantum states. The discrepancy between these two information theoretic quantities will be made clear in our analysis.

To obtain the quantum Euler relation, we consider the general setup of a quantum system \mathcal{S} with two arbitrary partitions A and B , in a Hilbert space $\mathcal{H}_{\mathcal{S}} = \mathcal{H}_A \otimes \mathcal{H}_B$. We start by exploring the connection between the information gain and bipartite correlations, classical and quantum, in the case of local measurements (cf. Figure 1). In particular, we show that correlations between the two partitions of \mathcal{S} directly hinder the amount of information gain accessible through measurements. Then, in Section 3, we present our main results by first separating the information gain into classical and quantum contributions, and by deriving upper bounds on \mathcal{I}_g as a function of the Holevo bound and thermodynamic quantities.

We illustrate the tightness of the latter bounds in a collective dissipation model, where two qubits are collectively coupled to a thermal bath. Remarkably, by combining the aforementioned results and beyond our illustrative case study, we arrive at fundamental statements. The first, constitutes the quantum Euler relation involving the maximum work that can be extracted from the state of \mathcal{S} (through unitary and cyclic operations) and the quantum contribution of the information gain \mathcal{I}_g . The second statement is a trade-off relation involving quantum correlations within the state of \mathcal{S} , and the entanglement that one of its partitions (A or B) shares with the surrounding environment.




Figure 1. Sketch illustrating fundamentally different ways one can carry out quantum measurements: either (i) globally or (ii) locally. For global measurements, we are usually interested in quantities such as the information gain \mathcal{I}_g and the Holevo bound $\chi_{\mathcal{S}}$. For local measurements, the direct quantities we measure are the local counterparts of \mathcal{I}_g and $\chi_{\mathcal{S}}$: $(\mathcal{I}_g^A, \mathcal{I}_g^B, \chi_A, \chi_B)$.

2. The Information Gain

We start by introducing notions and notations before we will derive upper bounds on \mathcal{I}_g . This will clarify the effect of bipartite correlations (classical and quantum) on the amount of \mathcal{I}_g that is accessible through measurements.

2.1. Positive Operator-Valued Measures

General quantum measurements [30–32] are described by a set of positive semi-definite operators, $\mathcal{M} = \{M_n\}_{n \in [1, d]}$, where “ n ” labels the measurement outcome. These operators M_n are called positive operator-valued measures or POVMs for short. They live on a C^* -algebra over the Hilbert space of the system being measured [30], and satisfy the completeness relation

$$\sum_{n=1}^d M_n^\dagger M_n = \mathbb{I}. \quad (2)$$

For a general mixed state, characterized by a density matrix ρ , the n th measurement outcome occurs with a probability $p(n)$,

$$p(n) = \text{tr}\{M_n \rho M_n^\dagger\}, \quad (3)$$

and the post-measurement state (after applying M_n) reads

$$\rho_n = \frac{M_n \rho M_n^\dagger}{p(n)}. \quad (4)$$

Additionally, by considering the set of measurements \mathcal{M} , the end state of the measurement process reads $\mathcal{M}(\rho) = \sum_{n=1}^d p(n) \rho_n$, which simplifies to $\mathcal{M}(\rho) = \sum_{n=1}^d M_n \rho M_n^\dagger$. Here, we are implicitly considering efficient measurements where each measurement M_n is represented by a single Kraus operator [28,33–38], as opposed to the case of weak [39] or inefficient measurements $M_n(\rho) = \sum_i M_{n,i} \rho M_{n,i}^\dagger$. Previous studies [33–37,40] have shown that the information gain \mathcal{I}_g is positive for all efficient measurements.

2.2. Maximal Information Gain

To probe the effect of correlations on the accessible information gain, we now derive two general upper bounds on \mathcal{I}_g . We begin by considering arbitrary POVMs, applied on a bipartite quantum system \mathcal{S} living on the Hilbert space $\mathcal{H}_\mathcal{S} = \mathcal{H}_A \otimes \mathcal{H}_B$, with partitions A and B , and dimension $d_\mathcal{S} = d_A d_B$. The information gain \mathcal{I}_g can be written as [28]

$$\mathcal{I}_g = S(\rho_{AB}) - \sum_n p_n S(\rho_n), \quad (5)$$

where $S(\rho) = -\text{tr}\{\rho \ln(\rho)\}$ is the von Neumann entropy, and the density matrices ρ_{AB} and ρ_n represent the pre-measurement and post-measurement (with probability p_n) states, respectively.

From the definition of the quantum mutual information between the two partitions A and B ,

$$\mathcal{I}(A : B) = S(\rho_A) + S(\rho_B) - S(\rho_{AB}), \quad (6)$$

and for any choice of POVMs, we arrive at a trivial bound by recognizing that the second term in the expression of the information gain (cf. Equation (5)) is negative, hence

$$\mathcal{I}_g \leq \ln(d_\mathcal{S}) - \mathcal{I}(A : B). \quad (7)$$

The bound in Equation (7) shows that the information contained in the correlations between A and B directly hinders the amount of accessible information gain. In other words, the higher the value of the mutual information $\mathcal{I}(A : B)$ the less information gain we can access through measurements.

Now, considering general, local POVMs applied on ρ_{AB} , of the form $M_n = M_{n_A} \otimes M_{n_B}$ (cf. Figure 1), we get

$$\begin{aligned} \mathcal{I}_g &= S(\rho_A) + S(\rho_B) - \mathcal{I}(A : B) - \sum_n p_n (S(\rho_A^n) + S(\rho_B^n) - \mathcal{I}^n(A : B)), \\ &= \left(S(\rho_A) - \sum_n p_n S(\rho_A^n) \right) + \left(S(\rho_B) - \sum_n p_n S(\rho_B^n) \right) - \mathcal{I}(A : B) + \sum_n p_n \mathcal{I}^n(A : B), \end{aligned} \quad (8)$$

hence,

$$\mathcal{I}_g = \mathcal{I}_g^A + \mathcal{I}_g^B - \mathcal{I}(A : B) + \sum_n p_n \mathcal{I}^n(A : B), \quad (9)$$

where \mathcal{I}_g^A and \mathcal{I}_g^B refer to the local information gain on each partition. We have [41,42]

$$\sum_n p_n \mathcal{I}^n(A : B) \leq \mathcal{I}(A : B), \quad (10)$$

which directly implies the inequality

$$\mathcal{I}_g \leq \mathcal{I}_g^A + \mathcal{I}_g^B, \quad (11)$$

that is reminiscent of the subadditivity of the von Neumann entropy [30]. In fact, from the above inequality we infer that, due to correlations between A and B , the total information gain we can access locally (from each partition) is greater or equal to the information gain of the composite system, and the more correlations between the partitions the less information gain we can access.

It is noteworthy that the type of measurements considered above (i.e., general local POVMs) describe more realistic settings [30,38,39,43], where one usually considers a quantum system interacting with an environment/ancilla. In such scenarios, measurements of a physical observable are either applied on the system (cf. [39], $M_n = M_n^A \otimes \mathbb{I}_B$) or simultaneously on the system and environment (cf. [43], $M_n = M_{n_A} \otimes M_{n_B}$). More specifically, to evaluate the work applied on the system or the heat exchanged with the environment, these measurements take the form of local projective energy measurements. The latter will be the main focus of the following section where we explore the thermodynamic significance of the information gain.

In addition it is interesting to note that from the expression in Equation (9) we can define a new quantity $\mathcal{I}_l = \mathcal{I}(A : B) - \sum_n p_n \mathcal{I}^n(A : B)$, which represents the amount of total correlations lost as a consequence of measuring \mathcal{S} . Note that, following Equation (10), the quantity \mathcal{I}_l is positive for local measurements. This quantity might prove useful in quantifying the invasiveness of quantum measurements, i.e., their effect on the correlations in a bipartite state. Finally, note that since the quantum mutual information does not increase under general local measurements we consider that the post measurement states have support on orthogonal subspaces [30]. In particular we forgo classical communication between the two partitions, and hence through general local measurements alone the quantum mutual information does not increase [44].

Illustrative Example:

Consider an arbitrary bipartite state described by a density matrix ρ_{AB} , and von Neumann measurements [30] $M_n = \Pi_{n_A} \otimes \Pi_{n_B}$ such that Π_{n_A} (Π_{n_B}) are rank-one projectors acting on subsystem A (B). In this scenario, Equation (11) becomes

$$S(\rho_{AB}) \leq S(\rho_A) + S(\rho_B). \quad (12)$$

This is nothing but the subadditivity of the von Neumann entropy [30], and it reflects the fact that the quantum mutual information is positive $\mathcal{I}(A : B) \geq 0$. Additionally, the inequality is saturated if and only if the partitions of the composite system are uncorrelated, i.e., $\rho_{AB} = \rho_A \otimes \rho_B$.

It is worth emphasizing that the subadditivity of the information gain Equation (11), is valid for general local measurements such that the post-measurement states have support on orthogonal subspaces [30]. In scenarios where these local measurements are represented by rank-one projectors we recover the subadditivity of the von Neumann entropy.

2.3. The Holevo Bound

However, before we continue with local measurements on a single partition and derive our main results, it is crucial to consider the subtle difference between the information gain and the Holevo bound, first. The Groenewold information gain is directly related to the Holevo bound χ_S (defined through measurement \mathcal{M} on the composite system S) via [38],

$$\mathcal{I}_g + \Delta = \chi_S, \quad (13)$$

where $\Delta = S(\mathcal{M}(\rho_{AB})) - S(\rho_{AB})$ represents the change in entropy caused by the measurement. This quantity is also referred to as the entropy cost of a measurement [38], and, as the name suggests, it quantifies the change in entropy between the pre- and post-measurement states. Depending on the type of measurements we carry out, the quantity Δ can be either positive or negative. For instance, it is straightforward to infer that for all projective measurements $\Delta \geq 0$ [30].

Illustrative Example:

To illustrate the discrepancy between these quantities we consider the same example we studied above. Namely, for von Neumann measurements $M_n = \Pi_{n_A} \otimes \Pi_{n_B}$ composed of rank-one projectors on each subsystem. Now, we further assume that the arbitrary state ρ_{AB} has coherences in the basis determined by the latter projectors. Thus, the Holevo bound becomes

$$\chi_S = S(\mathcal{M}(\rho_{AB})) - \sum_n p_n S(\rho_n), \quad (14)$$

which implies

$$\chi_S = S(\mathcal{M}(\rho_{AB})) - 0 = S(\rho_{AB}^{\text{diag}}), \quad (15)$$

where ρ_{AB}^{diag} is the incoherent state of S , generated by removing the off-diagonal elements of ρ_{AB} . On the other hand, the information gain is equal to the von Neumann entropy of S , $\mathcal{I}_g = S(\rho_{AB})$. Therefore, it is straightforward to get the strict inequality

$$\mathcal{I}_g < \chi_S, \quad (16)$$

since the difference between the two quantities ($\chi_S - \mathcal{I}_g$) is exactly the relative entropy of coherence [45] in the basis determined by the rank-one projectors.

3. Projective Local Measurements

We are now positioned to relate the information gain to thermodynamic quantities, and derive fundamental statements in quantum information theory as well as in quantum thermodynamics. To this end, we focus on the case of local projective energy measurements [43,46] such that

$$\mathcal{M} = \{\mathbb{I}_A \otimes |E_B^n\rangle\langle E_B^n|\}_{n \in [1, d_B]} = \{M_n\}_{n \in [1, d_B]}, \quad (17)$$

where $|E_B^n\rangle$ is the n th energy eigenstate of the self Hamiltonian of subsystem B , and d_B is the dimension of its Hilbert space. Moreover, we assume the state ρ_{AB} to be locally thermal (on both partitions) with inverse temperature β , i.e., $\rho_A = \exp(-\beta H_A)/Z_A$ and $\rho_B = \exp(-\beta H_B)/Z_B$, while A and B share correlations such that the global state ρ_{AB} is not necessarily thermal. Similar scenarios were discussed in the literature [21,26] to assign thermodynamic value to quantum correlations. More specifically, we have shown in Ref. [26] that the maximum work that can be extracted from a bipartite quantum state

" ρ_{AB} ", under unitary and cyclic operations, is directly related to the quantum mutual information " $\mathcal{I}(A : B)$ ", which will be crucial later on in our analysis.

It is a simple exercise to show that we have

$$(\forall n \in \llbracket 1, d_B \rrbracket); S(\rho_B^n) = \mathcal{I}^n(A : B) = 0. \quad (18)$$

Therefore, Equation (9) simplifies to

$$\begin{aligned} \mathcal{I}_g &= \mathcal{I}_g^A + S(\rho_B) - \mathcal{I}(A : B), \\ &= \chi_B + S(\rho_B) - \mathcal{I}(A : B), \end{aligned} \quad (19)$$

where χ_B is the Holevo information obtained by projective measurements on B (cf. Equation (17)), and it reflects the information we gain about the partition A by measuring subsystem B . In this case, we have $\mathcal{I}_g^A = \chi_B$ since we apply local measurements on B . The derivation directly follows from the cyclic property of the partial trace for the local measurements we are considering (cf. Equation (17)). More specifically, we have

$$\chi_B = S(\text{tr}_B\{\mathcal{M}(\rho_{AB})\}) - \sum_n p_n S(\rho_A^n) = S(\rho_A) - \sum_n p_n S(\rho_A^n), \quad (20)$$

which implies $\mathcal{I}_g^A = \chi_B$.

From the equality in Equation (19) we can separate the information gain into classical and quantum contributions, and we can relate \mathcal{I}_g to the maximum work extractable from ρ_{AB} . As a consequence of combining both results we will be able to derive fundamental statements for \mathcal{I}_g .

3.1. Quantum and Classical Contributions

Now exploiting the Koashi–Winter relation [47], Equation (19) can be written as

$$\mathcal{I}_g = \chi_B + \chi_A^{\max} + E(B : C) - \mathcal{I}(A : B), \quad (21)$$

for a subsystem " C " that purifies the state of $\mathcal{S} = A \cup B$. This subsystem can be regarded as the environment for the system of interest \mathcal{S} , such that the state of the quantum universe " ABC " is pure. We emphasize that we are implicitly adopting the realistic setting of a system of interest \mathcal{S} open to an environment " C ", with which it can share quantum as well as classical correlations. Moreover, χ_A^{\max} refers to the Holevo bound under optimal measurements on A , and " $E(B : C)$ " is the entanglement of formation [48] which represents a generalization of entanglement entropy to mixed states. To elaborate, the entanglement of formation " $E(B : C)$ " quantifies the entanglement in the mixed state ρ_{BC} , since the entanglement entropy is no longer a viable measure of entanglement for such states [30].

The expression in Equation (21) is further simplified by noting that the quantum mutual information can be decomposed into classical and quantum contributions [49,50]. In fact, the quantum contribution is known as quantum discord which is defined as the difference between the mutual information and the Holevo bound [49]

$$\mathcal{D}_A = \mathcal{I}(A : B) - \chi_A^{\max}, \quad (22)$$

hence it quantifies purely the quantum correlations in a bipartite state. From this definition, we get the equality

$$\mathcal{I}_g = \chi_B + (E(B : C) - \mathcal{D}_A), \quad (23)$$

which expresses the information gain, on the composite state of \mathcal{S} , in terms of classical and quantum contributions. The classical contribution is the Holevo information χ_B , and the quantum contribution is represented by the difference between the entanglement of formation $E(B : C)$ and quantum discord \mathcal{D}_A , which we will refer to as the quantum information gain " \mathcal{I}_g^Q " where $\mathcal{I}_g^Q = E(B : C) - \mathcal{D}_A$. This is a crucial result that will lead

to interesting conclusions, as we will explore in the remainder of our analysis. Namely, once we look at the problem from a thermodynamics perspective through the notion of ergotropy [51] (aka the maximum extractable work under unitary and cyclic operations).

3.2. Maximum Extractable Work

The maximum amount of work that can be extracted from the composite state ρ_{AB} , through unitary and cyclic operations, is referred to as the ergotropy “ \mathcal{E} ” [26,51,52]. Consider a quantum system with Hamiltonian $H = \sum_{i=1}^d \varepsilon_i |\varepsilon_i\rangle\langle\varepsilon_i|$ and quantum state $\rho = \sum_{j=1}^d r_j |r_j\rangle\langle r_j|$, such that $\varepsilon_i \leq \varepsilon_{i+1}$ and $r_j \geq r_{j+1}$. The ergotropy is calculated by performing an optimization over all possible unitary operations to achieve a final state that has the minimum average energy with respect to H ,

$$\mathcal{E}(\rho) = \text{tr}\{H\rho\} - \min_U \left\{ \text{tr}\{HU\rho U^\dagger\} \right\} = \text{tr}\{H(\rho - P_\rho)\}, \quad (24)$$

where $P_\rho \equiv \sum_k r_k |\varepsilon_k\rangle\langle\varepsilon_k|$ is called the passive state. An equivalent expression reads

$$\mathcal{E}(\rho) = \sum_{i,j} r_j \varepsilon_i \left(|\langle r_j | \varepsilon_i \rangle|^2 - \delta_{ij} \right). \quad (25)$$

Assuming the passive state P_ρ is not thermal, we can still extract work out of it, if we have access to, and ability to globally act upon, multiple copies of this state. This is captured by the bound ergotropy \mathcal{E}_b [26,53],

$$\mathcal{E}_b(\rho) = \text{tr}\left\{ (P_\rho - P_\rho^{\text{th}})H \right\}, \quad (26)$$

where P_ρ^{th} is a thermal state such that $S(P_\rho) = S(P_\rho^{\text{th}})$. In other words, P_ρ^{th} is the thermal state associated with the passive state P_ρ .

From the definition of the ergotropy and using the result of Ref. [26], namely the relationship between the quantum mutual information and the ergotropy, $\beta\mathcal{E} \leq \mathcal{I}(A : B)$, and Equation (19), we obtain

$$\mathcal{I}_g \leq \chi_B + \beta(\langle H_B \rangle - \mathcal{E} - F_B). \quad (27)$$

In the above inequality, $F_B = -(1/\beta) \log(Z_B)$ is the Helmholtz free energy. Furthermore, considering the bound ergotropy \mathcal{E}_b , where $\mathcal{E}_G = \mathcal{E} + \mathcal{E}_b$ is the global ergotropy, and using the general inequality $\beta\mathcal{E}_G \leq \mathcal{I}(A : B)$ from Ref. [26], we arrive at the tighter bound

$$\mathcal{I}_g \leq \chi_B + \beta(\langle H_B \rangle - \mathcal{E}_G - F_B). \quad (28)$$

In fact, the above inequality is saturated for a large class of states. This will be exemplified in the following section through a collective dissipation model. For the class of states where the above inequality is saturated, and using Equation (23), we get

$$E(B : C) - \mathcal{D}_A = \beta(\langle H_B \rangle - \mathcal{E}_G - F_B). \quad (29)$$

This result can be interpreted as follows. First, it represents a trade-off relation between the quantum correlations present in \mathcal{S} (quantified by \mathcal{D}_A) and the entanglement between a partition of \mathcal{S} (B in this case) and the surrounding environment (C). This is complementary to the Koashi–Winter relation, which represents a trade-off between classical information and entanglement. Second, simply by rearranging the terms, we obtain a quantum Euler relation involving ergotropy, the free energy, and the quantum correlations within \mathcal{S} as well as the entanglement that \mathcal{S} shares with its surroundings, namely

$$\langle H_B \rangle = \mathcal{E}_G + F_B + \mathcal{I}_g^Q / \beta. \quad (30)$$

Equation (30) constitutes our main result. It relates the quantum information gained from local measurements to inherent thermodynamic properties of a bipartite system.

4. Collective Dissipation

After having established the conceptual framework and the main results, this part of the analysis is dedicated to an illustrative case study. In particular, we illustrate the tightness of the bounds derived in Equations (27) and (28). To this end, we consider two-qubit X-shape density matrices of the form

$$\rho(t) = \begin{pmatrix} \rho_{11} & 0 & 0 & \rho_{14} \\ 0 & \rho_{22} & \rho_{23} & 0 \\ 0 & \rho_{23}^* & \rho_{33} & 0 \\ \rho_{14}^* & 0 & 0 & \rho_{44} \end{pmatrix}. \quad (31)$$

For such matrices, taking a partial trace on either qubits results in a thermal state with inverse temperature β . Here, the composite system \mathcal{S} is described by the density matrix $\rho(t)$. Therefore, the partitions A and B are single qubit. These X-shape density matrices are ubiquitous in the literature [54], as they can be found in Pauli channels [55], collective dephasing models for two-qubit systems [56], or in 1-D spin chains that exhibit \mathbb{Z}_2 symmetry [57–59].

Such X-states are characteristic for the collective dissipation model where a two-qubit system is collectively coupled to a thermal bath with inverse temperature β_e . The master equation of the dynamics reads [60–65]

$$\frac{\partial \rho}{\partial t} = -\frac{i}{\hbar}[(H_0 + H_d), \rho] + D_-(\rho) + D_+(\rho), \quad (32)$$

where $H_0 = \hbar\omega(\sigma_1^+ \sigma_1^- + \sigma_2^+ \sigma_2^-)$ and $H_d = \hbar f(\sigma_1^+ \sigma_2^- + \sigma_2^+ \sigma_1^-)$ are the self-Hamiltonian of the whole system and the interaction Hamiltonian between the qubits, respectively. The σ_i^+ and σ_i^- are the usual Pauli raising and lowering operators, respectively. Additionally,

$$\begin{aligned} D_-(\rho) &= \sum_{i,j=1}^2 \gamma_{ij} (\bar{n} + 1) (\sigma_j^- \rho \sigma_i^+ - \frac{1}{2} \{\sigma_i^+ \sigma_j^-, \rho\}), \\ D_+(\rho) &= \sum_{i,j=1}^2 \gamma_{ij} \bar{n} (\sigma_j^+ \rho \sigma_i^- - \frac{1}{2} \{\sigma_i^- \sigma_j^+, \rho\}). \end{aligned} \quad (33)$$

Here, $\bar{n} = [\exp(\beta_e \omega) - 1]^{-1}$ is the mean number of photons at the temperature of the environment β_e , and γ_{ij} are the spontaneous decay rates. One can show that there is no unique fixed point for this model [66,67] and the steady state ρ_{ss} depends on the initial state ρ_0 . In fact, the analytic expression for the X-shape steady states [61] is

$$\begin{aligned} \rho_{ss} &= (1 - c) |\psi_-\rangle \langle \psi_-| \\ &+ c Z_+^{-1}(\beta_e) (\exp(-2\omega\beta_e) |\psi_{ee}\rangle \langle \psi_{ee}| + \exp(-\omega\beta_e) |\psi_+\rangle \langle \psi_+| + |\psi_{gg}\rangle \langle \psi_{gg}|), \end{aligned} \quad (34)$$

where, $|\psi_{gg}\rangle = |gg\rangle$, $|\psi_{ee}\rangle = |ee\rangle$, $|\psi_{\pm}\rangle = |ge\rangle \pm |eg\rangle / \sqrt{2}$, $c = \langle \psi_{gg} | \rho_0 | \psi_{gg} \rangle + \langle \psi_{ee} | \rho_0 | \psi_{ee} \rangle + \langle \psi_+ | \rho_0 | \psi_+ \rangle$, and $Z_+(\beta_e) = 1 + \exp(-\omega\beta_e) + \exp(-2\omega\beta_e)$. It is instructive to realize that the parameter c plays the role of an effective coupling constant. For $c = 1$ the stationary state is thermal and the usual thermodynamic behavior is recovered. In particular, in this case the stationary state is independent of the initial preparation from within the set of states that result in $c = 1$, or in other words, the stationary state carries no memory of the initial state from within the invariant subspace with $c = 1$. In stark contrast, for $c = 0$ the stationary state is strongly dependent on the initial preparation. Thus, one would expect thermodynamic statements to be only valid for $c \simeq 1$, whereas only weaker statements can hold for arbitrary c .

Additionally, note that the inverse temperature β of each partition/qubit is directly related to the bath temperature and we have

$$\beta = \frac{1}{\omega} \ln \left[\frac{1 + 2 \cosh(\beta_e \omega) + 2c \sinh(\beta_e \omega)}{1 + 2 \cosh(\beta_e \omega) - 2c \sinh(\beta_e \omega)} \right]. \quad (35)$$

For more technical details on the collective dissipation model, interested readers can refer to Refs. [15,26].

Here we continue by illustrating Equation (27). To this end, we plot the right and left hand sides of the inequality for $\beta_e = 10$ and $\omega = 1$. In the case of $\omega = 1$ and in the limit of low temperatures (namely, $\beta_e = 10$), the ergotropy can be expressed with a simple analytic formula [15]

$$\begin{aligned} & \left(\forall c \in [0, \frac{1}{2}] \right); \quad \mathcal{E} = 1 - 2c, \\ & \left(\forall c \in [\frac{1}{2}, 1] \right); \quad \mathcal{E} = 0. \end{aligned} \quad (36)$$

From Figure 2, we observe that the upper bound is not a tight bound for almost all values of c , this is due to the fact that the mutual information is not a sharp upper bound to the ergotropy as observed in Ref. [26]. Additionally, the kink in the red curve at the value $c = 0.5$ is a direct consequence of the functional form of the ergotropy, cf. Equation (36). For a tighter bound we illustrate Equation (28) in Figure 3, and we observe that the upper bound is saturated for all steady states where roughly $c \in [0.7, 1]$.

The collective dissipation model we use in this section is a perfect example of an experimentally relevant scenario [57–59] where the inequality is presented in Equation (28) is saturated. Consequently, this model shows the existence of a class of states where the statements in Equations (29) and (30) are valid, which can be qualitatively explained as follows. Recall that c quantifies the information the steady state carries about the initial state of the dynamics (cf. Equation (34)). Therefore, as c gets closer to one we recover the limit where thermodynamic expressions such as the Euler relation (i.e., Equation (30)) are valid. Beyond this limit, we have the general inequality

$$\langle H_B \rangle \geq \mathcal{E}_G + F_B + \mathcal{I}_g^Q / \beta. \quad (37)$$




Figure 2. Plots of the information gain \mathcal{I}_g and the right hand side of Equation (27) as a function of c .




Figure 3. Plots of the information gain \mathcal{I}_g and the right hand side of Equation (28) as a function of c .

5. Concluding Remarks

In the present analysis, we studied the information gain \mathcal{I}_g , originally proposed by Groenewold, to derive the quantum analog of the Euler relation. More specifically, we studied the relationship between \mathcal{I}_g and bipartite total correlations (quantum and classical) which led to the quantum Euler relation as well as a general trade-off statement. First, we started by showing that total correlations hinder the amount of information gain \mathcal{I}_g that can be accessed through measurements. We then presented the direct connection between the Holevo bound and \mathcal{I}_g . In Section 3, we showcased our main results, where we considered bipartite states that are locally thermal. In particular, we proved that the information gain can be separated into quantum and classical contributions. Additionally, we derived upper bounds on \mathcal{I}_g as a function of the Holevo bound, the average and free energies, as well as the ergotropy. The aforementioned inequalities were illustrated in the experimentally relevant example of a two-qubit system collectively coupled to a thermal bath.

Remarkably, combining the separation of terms (quantum and classical) and the thermodynamic bounds on \mathcal{I}_g , we arrived at a fundamental identity that is valid for a large class of states. From an information theoretic perspective, we observed that the identity represents a general trade-off relation between genuine quantum correlations within a bipartite state (\mathcal{S}) and the entanglement that one of its partitions (A or B) shares with the surrounding environment (C). Interestingly, from a thermodynamics perspective, the identity represents a novel quantum formulation of the Euler relation. In this relation, the internal energy is expressed as a function of the ergotropy and the quantum information gain, directly displaying the role of quantum correlations as a thermodynamic resource, and the thermodynamic significance of the quantum information gained through local measurements. A table summarizing the statements derived throughout the paper can be found in Figure 4.

Our results highlight the importance of the Groenewold information gain \mathcal{I}_g as a quantifier for the information output from a quantum measurement and its effects on the state we are probing. The main inequalities we presented in this work can be further examined in various scenarios to determine the general class of states for which the quantum Euler relation and the trade-off expression are valid. We leave this analysis for future work on the topic.

Interestingly, the quantum Euler relation can be used in order to estimate the global ergotropy (the ergotropy plus the bound ergotropy) without performing an optimization over all possible unitary operations (cf. Equation (24)), and without acting globally on many copies of the state of \mathcal{S} (to evaluate \mathcal{E}_b). In such cases, we only have to apply local projective energy measurements, on a single partition of \mathcal{S} , in order to compute the global ergotropy for the state of \mathcal{S} , which can be extremely practical in experimental settings that rely on work extraction from correlations (i.e., work extraction from quantum batteries, through cyclic and unitary operations). In fact, IBM Quantum Experience [68–71] is an adequate platform to experimentally probe this idea, where projective energy measurements can be implemented

locally to extract all the relevant, thermodynamic and information theoretic, quantities in the quantum Euler relation. Furthermore, only recently, a novel method of implementing two point measurement schemes was introduced in Ref. [72], and implemented in IBM quantum processors, which has the potential to further increase the scope of our main result. More specifically, implementing the two point measurement scheme (applying projective measurements at two instants; $t = 0$ and $t = \tau$) to our conceptual framework would lead to quantum Euler relations defined at $t = 0$ and $t = \tau$. The difference between the two equalities is nothing but the first law of thermodynamics, which expresses the change in internal energy as a function of the change in the ergotropy and the change in the quantum information gain. Further probing this novel formulation of the first law of quantum thermodynamics and its limitations, in various settings, would constitute a significant step in understanding the intimate connection between quantum information and inherent thermodynamic properties of a system. We leave such a task for future endeavors on the topic.

	Results	Measurements
Quantum Information Theory	<i>Subadditivity of the information gain:</i> $\mathcal{I}_g \leq \mathcal{I}_g^A + \mathcal{I}_g^B$	$M_{n_A} \otimes M_{n_B}$
	<i>Separation of terms:</i> $\mathcal{I}_g = \chi_B + (E(B : C) - \mathcal{D}_A) \equiv \chi_B + \mathcal{I}_g^Q$	$\mathbb{I}_A \otimes \Pi_n$
	<i>Trade-off:</i> $(\exists \rho_{AB}); E(B : C) - \mathcal{D}_A = \beta (\langle H_B \rangle - \mathcal{E}_G - F_B)$	$\mathbb{I}_A \otimes E_B^n\rangle\langle E_B^n $
Quantum Thermodynamics	<i>Quantum Euler relation:</i> $(\exists \rho_{AB}); \langle H_B \rangle = \mathcal{E}_G + F_B + \frac{\mathcal{I}_g^Q}{\beta}$	$\mathbb{I}_A \otimes E_B^n\rangle\langle E_B^n $
	<i>General statement:</i> $(\forall \rho_{AB}); \langle H_B \rangle \geq \mathcal{E}_G + F_B + \frac{\mathcal{I}_g^Q}{\beta}$	$\mathbb{I}_A \otimes E_B^n\rangle\langle E_B^n $

Figure 4. Table summarizing the results of the present manuscript, with the main result shown in red.

Author Contributions: Conceptualization, A.T. and S.D.; formal analysis, A.T. and K.W.; writing—original draft preparation, A.T. and S.D.; writing—review and editing, A.T. and S.D.; supervision, S.D.; funding acquisition, S.D. All authors have read and agreed to the published version of the manuscript.

Funding: S.D. acknowledges support from the U.S. National Science Foundation under Grant No. DMR-2010127. This research was supported by grant number FQXi-RFP-1808 from the Foundational Questions Institute and Fetzer Franklin Fund, a donor advised fund of Silicon Valley Community Foundation (SD).

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: This work was conducted as part of the Undergraduate Research Program (K.W.) in the Department of Physics at UMBC.

Conflicts of Interest: The authors declare no conflict of interest.

References

- Callen, H.B. *Thermodynamics and an Introduction to Thermostatistics*; American Association of Physics Teachers: College Park, MD, USA, 1998.
- Huang, K. *Introduction to Statistical Physics*; CRC Press: Boca Raton, FL, USA, 2009.
- Carnot, S. *Réflexions Sur la Puissance Motrice du Feu*; Number 26; Vrin: Paris, France, 1978.
- Landauer, R. Irreversibility and Heat Generation in the Computing Process. *IBM J. Res. Dev.* **1961**, *5*, 183–191. [[CrossRef](#)]
- Bennett, C.H. The thermodynamics of computation—A review. *Int. J. Theor. Phys.* **1982**, *21*, 905–940. [[CrossRef](#)]
- Deffner, S.; Campbell, S. *Quantum Thermodynamics*; Morgan & Claypool Publishers: San Rafael, CA, USA, 2019; pp. 2053–2571. [[CrossRef](#)]
- Goold, J.; Huber, M.; Riera, A.; del Rio, L.; Skrzypczyk, P. The role of quantum information in thermodynamics—A topical review. *J. Phys. A Math. Theor.* **2016**, *49*, 143001. [[CrossRef](#)]
- Jarzynski, C. Nonequilibrium Equality for Free Energy Differences. *Phys. Rev. Lett.* **1997**, *78*, 2690–2693. [[CrossRef](#)]
- Crooks, G.E. Entropy production fluctuation theorem and the nonequilibrium work relation for free energy differences. *Phys. Rev. E* **1999**, *60*, 2721–2726. [[CrossRef](#)]
- Sagawa, T.; Ueda, M. Generalized Jarzynski Equality under Nonequilibrium Feedback Control. *Phys. Rev. Lett.* **2010**, *104*, 090602. [[CrossRef](#)]
- Deutsch, I.H. Harnessing the Power of the Second Quantum Revolution. *PRX Quantum* **2020**, *1*, 020101. [[CrossRef](#)]
- Dowling, J.P.; Milburn, G.J. Quantum technology: The second quantum revolution. *Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci.* **2003**, *361*, 1655–1674. [[CrossRef](#)]
- Francica, G.; Binder, F.C.; Guarnieri, G.; Mitchison, M.T.; Goold, J.; Plastina, F. Quantum Coherence and Ergotropy. *Phys. Rev. Lett.* **2020**, *125*, 180603. [[CrossRef](#)] [[PubMed](#)]
- Korzekwa, K.; Lostaglio, M.; Oppenheim, J.; Jennings, D. The extraction of work from quantum coherence. *New J. Phys.* **2016**, *18*, 023045. [[CrossRef](#)]
- Çakmak, B. Ergotropy from coherences in an open quantum system. *Phys. Rev. E* **2020**, *102*, 042111. [[CrossRef](#)]
- Guarnieri, G.; Morrone, D.; Çakmak, B.; Plastina, F.; Campbell, S. Non-equilibrium steady-states of memoryless quantum collision models. *Phys. Lett. A* **2020**, 126576. [[CrossRef](#)]
- Funo, K.; Watanabe, Y.; Ueda, M. Thermodynamic work gain from entanglement. *Phys. Rev. A* **2013**, *88*, 052319. [[CrossRef](#)]
- Francica, G.; Goold, J.; Plastina, F.; Paternostro, M. Daemonic ergotropy: Enhanced work extraction from quantum correlations. *NPJ Quantum Inf.* **2017**, *3*, 12. [[CrossRef](#)]
- Bernards, F.; Kleinmann, M.; Gühne, O.; Paternostro, M. Daemonic Ergotropy: Generalised Measurements and Multipartite Settings. *Entropy* **2019**, *21*, 771. [[CrossRef](#)] [[PubMed](#)]
- Manzano, G.; Plastina, F.; Zambrini, R. Optimal Work Extraction and Thermodynamics of Quantum Measurements and Correlations. *Phys. Rev. Lett.* **2018**, *121*, 120602. [[CrossRef](#)] [[PubMed](#)]
- Perarnau-Llobet, M.; Hovhannisyann, K.V.; Huber, M.; Skrzypczyk, P.; Brunner, N.; Acín, A. Extractable Work from Correlations. *Phys. Rev. X* **2015**, *5*, 041011. [[CrossRef](#)]
- Fusco, L.; Paternostro, M.; De Chiara, G. Work extraction and energy storage in the Dicke model. *Phys. Rev. E* **2016**, *94*, 052122. [[CrossRef](#)]
- Sapienza, F.; Cerisola, F.; Roncaglia, A.J. Correlations as a resource in quantum thermodynamics. *Nat. Commun.* **2019**, *10*, 1–7. [[CrossRef](#)]
- Oppenheim, J.; Horodecki, M.; Horodecki, P.; Horodecki, R. Thermodynamical Approach to Quantifying Quantum Correlations. *Phys. Rev. Lett.* **2002**, *89*, 180402. [[CrossRef](#)] [[PubMed](#)]
- Jevtic, S.; Jennings, D.; Rudolph, T. Maximally and Minimally Correlated States Attainable within a Closed Evolving System. *Phys. Rev. Lett.* **2012**, *108*, 110403. [[CrossRef](#)]
- Touil, A.; Çakmak, B.; Deffner, S. Second law of thermodynamics for quantum correlations. *arXiv* **2021**, arXiv:2102.13606.
- Sagawa, T.; Ueda, M. Second Law of Thermodynamics with Discrete Quantum Feedback Control. *Phys. Rev. Lett.* **2008**, *100*, 080403. [[CrossRef](#)]
- Groenewold, H.J. A problem of information gain by quantal measurements. *Int. J. Theo. Phys.* **1971**, *4*, 327. [[CrossRef](#)]
- Holevo, A.S. Bounds for the quantity of information transmitted by a quantum communication channel. *Probl. Peredachi Informatsii* **1973**, *9*, 3–11.
- Nielsen, M.A.; Chuang, I. Quantum computation and quantum information. *Am. J. Phys.* **2002**, *70*. [[CrossRef](#)]
- Von Neumann, J. *Mathematische Grundlagen der Quantenmechanik*; Springer: Berlin/Heidelberg, Germany, 2013; Volume 38.
- Thingna, J.; Talkner, P. Quantum measurements of sums. *Phys. Rev. A* **2020**, *102*, 012213. [[CrossRef](#)]
- Luo, S. Information conservation and entropy change in quantum measurements. *Phys. Rev. A* **2010**, *82*, 052103. [[CrossRef](#)]
- Buscemi, F.; Hayashi, M.; Horodecki, M. Global Information Balance in Quantum Measurements. *Phys. Rev. Lett.* **2008**, *100*, 210504. [[CrossRef](#)]

35. Ozawa, M. On information gain by quantum measurements of continuous observables. *J. Math. Phys.* **1986**, *27*, 759–763. [\[CrossRef\]](#)
36. Lindblad, G. An entropy inequality for quantum measurements. *Commun. Math. Phys.* **1972**, *28*, 245–249. [\[CrossRef\]](#)
37. Lindblad, G. Entropy, information and quantum measurements. *Commun. Math. Phys.* **1973**, *33*, 305–322. [\[CrossRef\]](#)
38. Xi, Z. Information gain and information leak in quantum measurements. *Phys. Rev. A* **2016**, *93*, 052308. [\[CrossRef\]](#)
39. Naghiloo, M.; Alonso, J.J.; Romito, A.; Lutz, E.; Murch, K.W. Information Gain and Loss for a Quantum Maxwell’s Demon. *Phys. Rev. Lett.* **2018**, *121*, 030604. [\[CrossRef\]](#)
40. Fuchs, C.A.; Jacobs, K. Information-tradeoff relations for finite-strength quantum measurements. *Phys. Rev. A* **2001**, *63*, 062305. [\[CrossRef\]](#)
41. Lindblad, G. Completely positive maps and entropy inequalities. *Commun. Math. Phys.* **1975**, *40*, 147–151. [\[CrossRef\]](#)
42. Vedral, V.; Rippin, M.; Plenio, M. Quantum correlations, local interactions and error correction. *J. Mod. Opt.* **1997**, *44*, 2185–2205. [\[CrossRef\]](#)
43. Deffner, S.; Lutz, E. Nonequilibrium Entropy Production for Open Quantum Systems. *Phys. Rev. Lett.* **2011**, *107*, 140404. [\[CrossRef\]](#)
44. Vedral, V.; Plenio, M.B.; Rippin, M.A.; Knight, P.L. Quantifying Entanglement. *Phys. Rev. Lett.* **1997**, *78*, 2275–2279. [\[CrossRef\]](#)
45. Baumgratz, T.; Cramer, M.; Plenio, M.B. Quantifying Coherence. *Phys. Rev. Lett.* **2014**, *113*, 140401. [\[CrossRef\]](#) [\[PubMed\]](#)
46. Kafri, D.; Deffner, S. Holevo’s bound from a general quantum fluctuation theorem. *Phys. Rev. A* **2012**, *86*, 044302. [\[CrossRef\]](#)
47. Koashi, M.; Winter, A. Monogamy of quantum entanglement and other correlations. *Phys. Rev. A* **2004**, *69*, 022309. [\[CrossRef\]](#)
48. Bennett, C.H.; DiVincenzo, D.P.; Smolin, J.A.; Wootters, W.K. Mixed-state entanglement and quantum error correction. *Phys. Rev. A* **1996**, *54*, 3824–3851. [\[CrossRef\]](#) [\[PubMed\]](#)
49. Ollivier, H.; Zurek, W.H. Quantum Discord: A Measure of the Quantumness of Correlations. *Phys. Rev. Lett.* **2001**, *88*, 017901. [\[CrossRef\]](#) [\[PubMed\]](#)
50. Henderson, L.; Vedral, V. Classical, quantum and total correlations. *J. Phys. A Math. Gen.* **2001**, *34*, 6899. [\[CrossRef\]](#)
51. Allahverdyan, A.E.; Balian, R.; Nieuwenhuizen, T.M. Maximal work extraction from finite quantum systems. *EPL* **2004**, *67*, 565. [\[CrossRef\]](#)
52. Sone, A.; Deffner, S. Quantum and classical ergotropy from relative entropies. *arXiv* **2021**, arXiv:2103.10850.
53. Niedenzu, W.; Huber, M.; Boukobza, E. Concepts of work in autonomous quantum heat engines. *Quantum* **2019**, *3*, 195. [\[CrossRef\]](#)
54. Quesada, N.; Al-Qasimi, A.; James, D.F. Quantum properties and dynamics of X states. *J. Mod. Opt.* **2012**, *59*, 1322–1329. [\[CrossRef\]](#)
55. Cai, J.M.; Zhou, Z.W.; Guo, G.C. Stability of pairwise entanglement in a decoherent environment. *Phys. Rev. A* **2005**, *72*, 022312. [\[CrossRef\]](#)
56. Carnio, E.G.; Buchleitner, A.; Gessner, M. Generating and protecting correlated quantum states under collective dephasing. *New J. Phys.* **2016**, *18*, 073010. [\[CrossRef\]](#)
57. Amico, L.; Fazio, R.; Osterloh, A.; Vedral, V. Entanglement in many-body systems. *Rev. Mod. Phys.* **2008**, *80*, 517–576. [\[CrossRef\]](#)
58. Sarandy, M.S.; De Oliveira, T.R.; Amico, L. Quantum discord in the ground state of spin chains. *Int. J. Mod. Phys. B* **2013**, *27*, 1345030. [\[CrossRef\]](#)
59. Çakmak, B.; Karpát, G.; Fanchini, F.F. Factorization and Criticality in the Anisotropic XY Chain via Correlations. *Entropy* **2015**, *17*, 790–817. [\[CrossRef\]](#)
60. Çakmak, B.; Manatuly, A.; Müstecaplıoğlu, O.E. Thermal production, protection, and heat exchange of quantum coherences. *Phys. Rev. A* **2017**, *96*, 032117. [\[CrossRef\]](#)
61. Latune, C.L.; Sinayskiy, I.; Petruccione, F. Energetic and entropic effects of bath-induced coherences. *Phys. Rev. A* **2019**, *99*, 052105. [\[CrossRef\]](#)
62. Gross, M.; Haroche, S. Superradiance: An essay on the theory of collective spontaneous emission. *Phys. Rep.* **1982**, *93*, 301–396. [\[CrossRef\]](#)
63. Stephen, M.J. First-Order Dispersion Forces. *J. Chem. Phys.* **1964**, *40*, 669–673. [\[CrossRef\]](#)
64. Lehmberg, R.H. Radiation from an *N*-Atom System. I. General Formalism. *Phys. Rev. A* **1970**, *2*, 883–888. [\[CrossRef\]](#)
65. Damanet, F.; Martin, J. Competition between finite-size effects and dipole–dipole interactions in few-atom systems. *J. Phys. B At. Mol. Opt. Phys.* **2016**, *49*, 225501. [\[CrossRef\]](#)
66. Thingna, J.; Manzano, D.; Cao, J. Dynamical signatures of molecular symmetries in nonequilibrium quantum transport. *Sci. Rep.* **2016**, *6*, 1–11. [\[CrossRef\]](#)
67. Thingna, J.; Manzano, D.; Cao, J. Magnetic field induced symmetry breaking in nonequilibrium quantum networks. *New J. Phys.* **2020**, *22*, 083026. [\[CrossRef\]](#)
68. Deffner, S. Demonstration of entanglement assisted invariance on IBM’s quantum experience. *Heliyon* **2017**, *3*, e00444. [\[CrossRef\]](#)
69. Berta, M.; Wehner, S.; Wilde, M.M. Entropic uncertainty and measurement reversibility. *New J. Phys.* **2016**, *18*, 073004. [\[CrossRef\]](#)
70. Devitt, S.J. Performing quantum computing experiments in the cloud. *Phys. Rev. A* **2016**, *94*, 032329. [\[CrossRef\]](#)
71. Rundle, R.P.; Mills, P.W.; Tilma, T.; Samson, J.H.; Everitt, M.J. Simple procedure for phase-space measurement and entanglement validation. *Phys. Rev. A* **2017**, *96*, 022117. [\[CrossRef\]](#)
72. Solfanelli, A.; Santini, A.; Campisi, M. Experimental verification of fluctuation relations with a quantum computer. *arXiv* **2021**, arXiv:2106.04388.