

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

**Please provide feedback**

Please support the ScholarWorks@UMBC repository by emailing [scholarworks-group@umbc.edu](mailto:scholarworks-group@umbc.edu) and telling us what having access to this work means to you and why it's important to you. Thank you.

# Tectonically controlled subglacial lakes on the flanks of the Gamburtsev Subglacial Mountains, East Antarctica

Robin E. Bell,<sup>1</sup> Michael Studinger,<sup>1</sup> Mark A. Fahnestock,<sup>2</sup> and Christopher A. Shuman<sup>3</sup>

Received 11 November 2005; revised 6 December 2005; accepted 12 December 2005; published 25 January 2006.

[1] The morphology of surface lakes strongly influences their ecology and limnology (Wetzel, 2001). This morphology is a result of both the geologic processes that produce topographic basins and the regional climatic and local hydrologic processes that control water depth and sediment infilling (Carroll and Bohacs, 1999). Although basin forming processes range from glacial scour to meteorite impacts (Cohen, 2003), the deepest, oldest surface lakes are tectonically controlled (Meybeck, 1995) and contain diverse exotic ecosystems (Rossiter and Kawanabe, 2000). Subglacial lakes are also thought to be ancient systems that may contain exotic biota (Bulat et al., 2004; Karl et al., 1999; Priscu et al., 1999). Here we present evidence for the scale and configuration of 2 large subglacial lakes in East Antarctica that together with Lake Vostok define a province of major lakes on the flanks of the Gamburtsev Subglacial Mountains. Spatially-defined in the new Moderate Resolution Imaging Spectroradiometer (MODIS) imagery of Antarctica (T. Scambos et al., A MODIS-based mosaic of Antarctica: MOA, submitted to *Remote Sensing of Environment*, 2005, hereinafter referred to as Scambos et al., submitted manuscript, 2005), these lakes are aligned parallel to Lake Vostok. Other data shows that they are distinguished by distinct gravity lows, flat ice surface slopes and have estimated water depths of at least 900 m. Surface elevation data indicates that large deep subglacial lakes have a profound influence on the regional ice sheet topography and probably ice sheet flow. These deep subglacial lakes with elongate, rectilinear morphology are tectonically controlled features. Unlike the shallow lakes in West Antarctica and beneath Dome Concordia, these deep subglacial lakes remained stable environments through many glacial cycles since their origin 10–35 Ma enabling the development of novel ecosystems. **Citation:** Bell, R. E., M. Studinger, M. A. Fahnestock, and C. A. Shuman (2006), Tectonically controlled subglacial lakes on the flanks of the Gamburtsev Subglacial Mountains, East Antarctica, *Geophys. Res. Lett.*, 33, L02504, doi:10.1029/2005GL025207.

## 1. Subglacial Lakes Distribution

[2] Since the first evidence for subglacial lakes was advanced in 1970 [Robin et al., 1970] over 145 subglacial lakes have been identified in Antarctica ranging in length

from 1 to 280 km [Bogorodsky and Sheremet'yev, 1981; Kapitsa et al., 1996; Siegert et al., 2005]. While many small and shallow lakes have been found, no lakes similar in area or volume to Lake Vostok (14,000 km<sup>2</sup>, volume of 5,400 km<sup>3</sup>) [Kapitsa et al., 1996; Studinger et al., 2004] have been identified. Subglacial lakes are concentrated along the ice divides of Dome Concordia and Ridge B where the ice thickness is close to 4000 m [Siegert et al., 2005]. In the Dome Concordia lake province, the subglacial lakes are generally small (<20 km<sup>2</sup>) with the exception of the 45 km long Lake Concordia (800 km<sup>2</sup>, volume of 160 km<sup>3</sup>) [Tabacco et al., 2003; Tikku et al., 2005]. The Ridge B province includes Lake Vostok and smaller lakes to the west (Figures 1 and 2).

## 2. Description of the 90°E and Sovetskaya Lakes

[3] Here we describe in detail, a large deep subglacial lake aligned along 90°E and a lake beneath Sovetskaya Station. These lakes have been identified in earlier inventories but their size, depth and origin have not been investigated [Robin et al., 1970; Siegert et al., 2005; Studinger et al., 2003b]. Our analysis is based on five data sets including a multiple-image-composite mosaic using MODIS satellite imagery (Scambos et al., submitted manuscript, 2005), aerogeophysical data from the 2000–2001 survey of Lake Vostok [Studinger et al., 2003a], laser-based ice surface altimetry from NASA's Ice, Cloud and Land Elevation Satellite (ICESat), ground traverse data from the 3rd Soviet Antarctic Expedition (SAE, 1958–59) [Sorokhtin et al., 1964], and radar-based satellite ice surface elevation data from ERS-1 [Liu et al., 2001].


[4] The aerogeophysical profile over the 90°E lake extends from Lake Vostok to the foothills of the Gamburtsev Subglacial Mountains, a region thought to be a foreland basin [Studinger et al., 2003b]. Between Lake Vostok and 90°E, the subglacial topography is very rugged, ranging from –300 m to +2400 m asl. The western 7 km of the aerogeophysical ice-penetrating radar data line contains a very bright horizontal basal reflector (Figure 3), characteristic of water contained within subglacial lakes [Robin et al., 1970]. The ice-penetrating radar data over the Sovetskaya lake documents an ice thickness of 4200 m over the lake.

[5] The MODIS mosaic in this region (Figures 1 and 2) reveals the ice surface topography largely as a function of the slope-controlled patterns of brightness with illumination from solar azimuths between 0–90° East (upper right in Figures 1 and 2). In the mosaic, the 90°E lake resembles Lake Vostok; it is characterized by a 123 km long flat featureless ice surface elongate in the north-south orientation and 20 km wide in the center tapering to 7 km wide at the ends. To the southwest of the 90°E lake, the Sovetskaya

<sup>1</sup>Lamont-Doherty Earth Observatory of Columbia University, Palisades, New York, USA.

<sup>2</sup>Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, New Hampshire, USA.

<sup>3</sup>NASA Goddard Space Flight Center, Greenbelt, Maryland, USA.



**Figure 1.** MODIS mosaic of Ridge B region including the 90°E and Sovetskaya lakes as well as Lake Vostok. Geographic north is to the right. The inset map of Antarctica indicates the image location, the Gamburtsev Subglacial Mountains (GSM) and Dome C (DC). Easting and northing are based on a polar stereographic projection with a latitude of true scale at 71°S. The dashed box indicates the location of Figure 2 and the solid white lines locate the profiles in Figure 4.

lake appears as a rectilinear flat featureless region 35 km wide in the east-west orientation that extends for 75 km in a north-south direction.

[6] Over the eastern grounding line of the 90°E lake, the airborne laser altimeter data and the ERS-1 ice surface altimetry (Figure 3a) both indicate a 14-m-deep and 10-km-wide trough in the ice surface coincident with the limit of the bright basal reflector in the ice-penetrating radar and the eastern limit of the flat featureless texture in the MODIS imagery. The ice surface trough at the upstream lake margin (Figure 2) is similar in size and shape to the 5-m-deep and 5-km-wide trough along the upstream margin of Lake Vostok (Figure 4) [Studinger *et al.*, 2003a]. Based on the grounding line trough and the bright horizontal reflector coincident with the flat featureless ice surface, we interpret this structure as a large ( $\sim 2000 \text{ km}^2$ ) subglacial lake, aligned approximately north-south along 90°E. This 90°E lake, similar in area to the state of Rhode Island, is second only Lake Vostok in surface area for a subglacial lake. The closed contour along the eastern edge of the Sovetskaya lake from the ERS-1 data is associated with an 8 m deep grounding line trough resolved by the ICESat altimetry. The estimated surface area of the Sovetskaya lake is  $\sim 1600 \text{ km}^2$ .


### 3. Identification of the Tilted Ceilings: Implications for Circulation

[7] Based on the gradient of the ERS-1 ice surface (Figure 2), the primary ice flow over the 90°E lake is from Ridge B in a generally westerly orientation although a southwesterly component is evident at the southern end of the lake. The ERS-1 ice surface over the 90°E lake [Liu *et al.*, 2001] slopes up 30 m to the north from 3660 m in south to 3690 m in north. In an east-west orientation, the ice surface slopes up 15 m from 3665 m at the eastern shoreline to 3680 m at the western shoreline (Figure 3a). The ice surface gradient indicates the ice flow over the Sovetskaya


lake is from southeast to northwest and the ice surface slopes 20 m down to the north from 3670 m in the south to 3650 m in the north. A bedrock ridge bisects the Sovetskaya lake producing a relatively complex lake morphology evidenced in the MODIS imagery and the ice surface elevation.

[8] Using the ice thickness measurements along the ice-penetrating radar profile and the ice surface elevation, we can estimate the ice thickness over the lake assuming the ice sheet is in hydrostatic equilibrium and the lake water is fresh. To test the method, we predicted the ice thickness along the 7 km flight line. The standard deviation between the observed and predicted ice thickness is 9 m, which is below the  $\sim 15 \text{ m}$  resolution of the radar data (Figure 3b). Along this profile the observed ice thickness is 3980 m at the eastern ground line and the estimated ice thickness is 4032 m at the western shoreline. The estimated ice thickness along a north-south profile over the lake changes from 3900 m in the south to 4156 m in the north. Over the Sovetskaya lake the 20 m change in surface elevation observed in the ERS-1 ice surface (Figure 2) requires a 200 m thinning of the ice sheet from the observed 4200 m in the south [Robin *et al.*, 1970] to an estimated ice thickness of 4000 m in the north.

[9] Melting and freezing have been documented in both Lake Vostok and Lake Concordia based on accretion ice reflectors [Bell *et al.*, 2002], basal ice chemistry [Jouzel *et al.*, 1999] and ice thicknesses [Tikku *et al.*, 2004, 2005]. Above both the 90°E and Sovetskaya lakes, the overlying ice sheet forms a tilted ceiling similar to the ice sheet above Vostok. The tilted ceilings will introduce lake circulation as



**Figure 2.** Subglacial lakes near 90°E. The base map is the MODIS mosaic (Scambos *et al.*, submitted manuscript, 2005). 90°E is the zero line along the y axis (km northing). The 10 m ice surface elevation contours are from the ERS-1 data [Liu *et al.*, 2001]. The triangles are the lake locations [Siebert *et al.*, 2005]. The white line shows the location of Figure 3 with the solid line indicating the extent of the radar data shown and the dashed line indicating the profile where the ice thickness was estimated. The colored points indicate free-air gravity measurements from the 1958 SAE in mGal [Sorokhtin *et al.*, 1964].



**Figure 3.** Stacked profiles along profile over 90°E Lake. Location is shown in Figure 2. Subglacial lake is from 15–35 km across as indicated by double-ended arrows. (a) Airborne laser altimetry [Studinger *et al.*, 2003b] and ERS-1 ice surface [Liu *et al.*, 2001] measurements along profile. (b) Ice-penetrating radar and predicted ice thickness (dashed line) along profile. Accreted ice basal reflector is located at 32 km along profile.

well as melting and freezing of the ice sheet [Mayer *et al.*, 2003; Wüest and Carmack, 2000]. Over the 90°E lake, the ice thickness increases rapidly adjacent to the eastern shoreline and a weak basal reflector (Figure 3b, 32 km) indicates that accretion of lake water as ice may occur along the grounding line as the lake dynamically interacts with the overlying ice sheet.

#### 4. Depth Estimate: Support for Tectonic Origin of Lakes

[10] Gravity data can be used to estimate water depth of subglacial lakes [Studinger *et al.*, 2004]. The 3rd SAE traverse conducted geophysical studies across this region. While no seismic measurements were acquired over the lake [Sorokhtin *et al.*, 1964], gravity data was collected every 11 km (Figure 2). A distinct 80 mGal free-air gravity low is centered over the 90°E lake and a 70 mGal free-air gravity low exists over the Sovetskaya lake. Simple models of the gravity field can be used to constrain the water depths in the 90°E lake using the calculated ice thickness tied to the radar estimates. For models with a bedrock density of 2670 kg/m<sup>3</sup>, a minimum water depth of 900 m is estimated based on the topography observed along the airborne radar profile. The estimated volume of the 90°E lake is ~1800 km<sup>3</sup>. The estimated water depths are similar to the maximum water depths resolved with both seismic and gravity inversions over Lake Vostok [Masolov *et al.*, 1999, 2001; Studinger *et al.*, 2004] and are also similar to other tectonically controlled lakes such as fault-bounded lakes including Tahoe, USA (501 m) and Issyk-kul, Kyrgyzstan (668 m) as well as rift lakes including Tanganyika, Africa (1479 m), Malawi, Africa (706 m) and Baikal, Siberia (1637 m) [Herdendorf, 1982]. With the exception of the Great Slave Lake, Canada (624 m) glacially


scoured lakes tend to have maximum water depths of less than 420 m [Herdendorf, 1982]. While the majority of surface lakes are glacial in origin (75%) [Meybeck, 1995], the majority (85%) of the deep lakes (>500 m) are tectonic in origin [Herdendorf, 1982; Cohen, 2003]. We interpret the steep, rectilinear morphology of these subglacial lakes to indicate a tectonic origin. While tectonic control of these basins is not indicative of active tectonics or elevated geothermal heat flow, the basin bounding faults should provide conduits of active fluid flow rich in dissolved minerals into the lakes. These deep elongate basins probably pre-date the onset of Antarctic glaciation and likely contained surface lakes prior to becoming encased in ice.

#### 5. Influence of Large Subglacial Lakes on Ice Sheet Geometry

[11] Subglacial lakes are often assumed to have little influence on ice sheet dynamics. Ice sheet elevation profiles across the 90°E lake and Lake Vostok demonstrate that large subglacial lakes have a profound effect on the surface topography of the ice sheet, and by inference, on ice flow. The ice sheet elevation profiles (Figure 4) show the parabolic ice sheet profile with 5–10 m undulations characteristic of grounded ice flow in the interior, until the lakes are reached. At the upflow grounding line, the ice surface slope steepens, producing a marginal trough 5 to 20 meters deep. The profiles show a characteristic gentle downward slope over both lakes, and a ridge at the downstream grounding line. This pattern in the topographic profiles reflects the strong transitions in basal boundary conditions that control the rate that ice enters and leaves the lakes.

#### 6. Summary and Implications

[12] It is striking that these newly defined lakes and the eastern shoreline of Lake Vostok are all aligned parallel to 90°E and that Lake Vostok and the 90°E lake have a significant impact on the ice sheet geometry. Studinger *et al.* [2003b] demonstrated that Lake Vostok formed along the



**Figure 4.** Ice surface elevation over 90°E lake (blue) and Lake Vostok (red). The profiles over the 90°E lake are from ICESat's Laser 2a operations period and are somewhat oblique to ice flow. These GLAS12 Release 21 elevations have a spacing along track of ~172 m and vertical accuracy of a few decimeters based on analysis of crossover residuals (C. A. Shuman *et al.*, ICESat Antarctic elevation data: Preliminary precision and accuracy assessment, submitted to *Geophysical Research Letters*, 2005). Profile locations are shown in Figure 1.



eastern edge of a foreland basin that extends to the west beneath Ridge B toward the Gamburtsev Subglacial Mountains and that the linear eastern shoreline of Lake Vostok is a fault controlled tectonic boundary. The subrectangular morphology of the Sovetskaya and 90°E lakes and their position along the western edge of this foreland basin indicate these features may be similarly fault controlled. The tectonic fabric of this foreland basin or pre-existing structures [Studinger *et al.*, 2003b] provides the template for the elongate fault-bounded topographic depressions necessary to form this province of large, deep subglacial lakes [Wetzel, 2001; Meybeck, 1995]. The tectonically controlled depth of these lakes should provide consistent water depths through changing climatic conditions over the past 10–35 My [Carroll and Bohacs, 1999]. These deep subglacial lakes are likely to have been stable environments through many glacial cycles and may have developed novel ecosystems [Rossiter and Kawanabe, 2000; Bulat *et al.*, 2004; Karl *et al.*, 1999; Priscu *et al.*, 1999], in contrast to many of the shallow lakes in West Antarctica [Siebert *et al.*, 2004] and beneath Dome Concordia [Tabacco *et al.*, 2003; Tikku *et al.*, 2005].

[13] **Acknowledgments.** Bell and Studinger were supported by the Doherty Endowment of LDEO and the Palisades Geophysical Institute. Fahnestock and Shuman were supported by NASA's Cryospheric program for work on mosaic production and altimetry. We thank NASA's Science Mission Directorate, Cryospheric Sciences Program, and the ICESat Mission for both support and data. The acquisition of the radar data has been supported by NSF as part of the study of Lake Vostok. We thank the NSIDC and Ted Scambos for providing data. LDEO contribution 6851.

## References

- Bell, R. E., *et al.* (2002), Origin and fate of Lake Vostok water frozen to the base of the East Antarctic ice sheet, *Nature*, **416**, 307–310.
- Bogorodsky, V. V., and A. N. Sheremet'yev (1981), Subglacial lakes of Antarctica (Podlednikovye ozera Antarktidy), *Priroda*, **12**, 49–51.
- Bulat, S. A., *et al.* (2004), DNA signature of thermophilic bacteria from the aged accretion ice of Lake Vostok, Antarctica: Implications for searching for life in extreme ice environments, *Int. J. Astrobiol.*, **3**, 1–12.
- Carroll, R. A., and K. M. Bohacs (1999), Stratigraphic classification of ancient lakes: Balancing tectonic and climatic controls, *Geology*, **27**, 99–102.
- Cohen, A. S. (2003), *Paleolimnology: The History and Evolution of Lake Systems*, 500 pp., Oxford Univ. Press, New York.
- Herdendorf, C. E. (1982), Large lakes of the world, *J. Great Lakes Res.*, **8**, 379–412.
- Jouzel, J., *et al.* (1999), More than 200 meters of lake ice above subglacial Lake Vostok, Antarctica, *Science*, **286**, 2138–2141.
- Kapitsa, A. P., J. K. Ridley, G. D. Robin, M. J. Siebert, and I. A. Zotikov (1996), A large deep freshwater lake beneath the ice of central East Antarctica, *Nature*, **381**, 684–686.
- Karl, D. M., *et al.* (1999), Microorganisms in the accreted ice of Lake Vostok, Antarctica, *Science*, **286**, 2144–2147.
- Liu, H. X., K. C. Jezek, B. Li, and Z. Zaho (2001), Radarsat Antarctic Mapping Project digital elevation model version 2, <http://nsidc.org/data/nsidc-0082.html>, *Natl. Snow and Ice Data Cent.*, Boulder, Colo.
- Masolov, V. N., *et al.* (1999), Earth science studies in the Lake Vostok region: Existing data and proposals for future research, paper presented at International Workshop on Subglacial Lake Exploration, Sci. Comm. on Antarct. Res., Cambridge, U. K.
- Masolov, V. N., V. V. Lukin, A. N. Sheremetiev, and S. V. Popov (2001), Geophysical investigations of the subglacial Lake Vostok in eastern Antarctica, *Dokl. Earth Sci.*, **379**, 734–738.
- Mayer, C., K. Grosfeld, and M. J. Siebert (2003), Salinity impact on water flow and lake ice in Lake Vostok, Antarctica, *Geophys. Res. Lett.*, **30**(14), 1767, doi:10.1029/2003GL017380.
- Meybeck, M. (1995), Global distribution of lakes, in *Physics and Chemistry of Lakes*, edited by A. Lerman *et al.*, pp. 1–35, Springer, New York.
- Priscu, J. C., *et al.* (1999), Geomicrobiology of subglacial ice above Lake Vostok, Antarctica, *Science*, **286**, 2141–2144.
- Robin, G. d. Q., C. W. M. Swithinbank, and B. M. E. Smith (1970), Radio echo exploration of the Antarctic ice sheet, paper presented at International Symposium on Antarctic Glaciological Exploration (ISAGE), Sci. Comm. on Antarct. Res., Hanover, N. H., 3–7 Sept.
- Rossiter, A., and H. Kawanabe (Ed.) (2000), *Ancient Lakes: Biodiversity, Ecology and Evolution*, 624 pp., Elsevier, New York.
- Siebert, M. J., *et al.* (2000), Subglacial Lake Ellsworth: A candidate for in situ exploration in West Antarctica, *Geophys. Res. Lett.*, **31**, L23403, doi:10.1029/2000GL014777.
- Siebert, M. J., S. Carter, I. Tabacco, S. Popov, and D. Blankenship (2005), A revised inventory of Antarctic subglacial lakes, *Antarct. Sci.*, **17**, 453–460.
- Sorokhtin, O. G., Y. Avsyuk, and V. I. Koptev (1964), Determination of the thickness of the ice cap in East Antarctica, in *Soviet Antarctic Expedition: Information Bulletin*, pp. 4–7, Elsevier, New York.
- Studinger, M., *et al.* (2003a), Ice cover, landscape setting, and geological framework of Lake Vostok, East Antarctica, *Earth Planet. Sci. Lett.*, **205**, 195–210.
- Studinger, M., *et al.* (2003b), Geophysical models for the tectonic framework of the Lake Vostok region, East Antarctica, *Earth Planet. Sci. Lett.*, **216**, 663–677.
- Studinger, M., R. E. Bell, and A. A. Tikku (2004), Estimating the depth and shape of subglacial Lake Vostok's water cavity from aerogravity data, *Geophys. Res. Lett.*, **31**, L12401, doi:10.1029/2004GL019801.
- Tabacco, I. E., *et al.* (2003), Evidence of 14 new subglacial lakes in the Dome C-Vostok area, *Terra Antarct. Rep.*, **8**, 175–179.
- Tikku, A. A., R. E. Bell, M. Studinger, and G. K. C. Clarke (2004), Ice flow over field Lake Vostok, East Antarctica, inferred by structure tracking, *Earth Planet. Sci. Lett.*, **227**, 249–261.
- Tikku, A. A., *et al.* (2005), Influx of meltwater to subglacial Lake Concordia, East Antarctica, *J. Glaciol.*, **51**(172), 96–104.
- Wetzel, R. G. (2001), *Limnology: Lake and River Ecosystems*, 3rd ed., 1006 pp., Elsevier, New York.
- Wüest, A., and E. Carmack (2000), A priori estimates of mixing and circulation in the hard-to-reach water body of Lake Vostok, *Ocean Modell.*, **2**, 29–43.

R. E. Bell and M. Studinger, Lamont-Doherty Earth Observatory of Columbia University, Palisades, NY 10964, USA. (robinb@ldeo.columbia.edu)

M. A. Fahnestock, Institute for the Study of Earth, Oceans, and Space, University of New Hampshire, Durham, NH 03824, USA.

C. A. Shuman, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA.