

This work was written as part of one of the author's official duties as an Employee of the United States Government and is therefore a work of the United States Government. In accordance with 17 U.S.C. 105, no copyright protection is available for such works under U.S. Law.

Public Domain Mark 1.0

<https://creativecommons.org/publicdomain/mark/1.0/>

Access to this work was provided by the University of Maryland, Baltimore County (UMBC) ScholarWorks@UMBC digital repository on the Maryland Shared Open Access (MD-SOAR) platform.

Please provide feedback

Please support the ScholarWorks@UMBC repository by emailing scholarworks-group@umbc.edu and telling us what having access to this work means to you and why it's important to you. Thank you.

EGU23-10536, updated on 03 Apr 2023

<https://doi.org/10.5194/egusphere-egu23-10536>

EGU General Assembly 2023

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



MAVEN Observations of Steepened Ultra-Low Frequency Waves in the Upstream Martian Foreshock Region

Gangkai Poh^{1,2}, Jared Espley¹, Shaosui Xu³, Guan Le¹, Norberto Romanelli^{1,4}, Jasper Halekas⁵, Gina DiBraccio¹, and Jacob Gruesbeck¹

¹Solar System Exploration Division, NASA Goddard Space Flight Center, Greenbelt, Maryland, USA

²Center for Research and Exploration in Space Sciences and Technology II, Catholic University of America, Washington DC, USA

³Space Sciences Laboratory, University of California, Berkeley, Berkeley, CA, USA,

⁴Center for Research and Exploration in Space Sciences and Technology II, University of Maryland Baltimore County, Baltimore, MD, USA

⁵Department of Physics and Astronomy, University of Iowa, Iowa City, IA, USA

In this study, we present the analysis of steepened ultra-low frequency (ULF) waves in the foreshock region upstream of Mars' bow shock observed by the Mars Atmosphere and Volatile EvolutionN (MAVEN) spacecraft at Mars. A survey of MAVEN magnetic field and plasma measurements shows quasi-periodic gradual increases followed by a sharp decrease in the magnetic field magnitude (B_{total}). Higher frequency waves were also commonly, but not always, observed at the trailing edge of the large-amplitude increase in B_{total} . These observations are consistent with the signatures of shocklets observed in the solar wind region upstream of Earth's and planetary bow shocks. Shocklets are believed to be formed as a result of the steepening of fast magnetosonic waves generated by reflected ions in the quasi-parallel foreshock region. We also performed the minimum variance analysis (MVA) and statistical analysis technique to determine the wave properties (e.g. polarization, wave propagation, amplitude and frequency) of the shocklets and higher frequency waves observed in its trailing edge. Our results showed that these shocklets are left-handed polarized in the spacecraft frame, with mean amplitude $\delta B/B$ of ~ 3.5 and time separation between adjacent shocklet events of ~ 40 s. We also analyzed measurements (ions and electrons) from MAVEN's plasma instruments to investigate the energization process of the particles during the observations of shocklets. We will discuss the possible generation mechanisms for these steepened ultra-low frequency waves at Mars, and any implications for the martian plasma environment downstream of Mars' bow shock.